

## **Common genetic and clinical risk factors: Association with fatal prostate cancer in the Cohort of Swedish Men**

Minh-Phuong Huynh-Le, MD, MAS<sup>1,2</sup>,  
Roshan Karunamuni, PhD<sup>2,3</sup>,  
Chun Chieh Fan, MD, PhD<sup>3</sup>,  
Wesley K. Thompson, PhD<sup>4,5</sup>,  
Kenneth Muir, PhD<sup>6,7</sup>,  
Artitaya Lophatananon, PhD<sup>6</sup>,  
Karen Tye, MD, MS<sup>2</sup>,  
Alicja Wolk, DMSc<sup>8,9</sup>,  
Niclas Håkansson, PhD<sup>8</sup>,  
Ian G. Mills, PhD<sup>10</sup>,  
Ole A. Andreassen, MD, PhD<sup>11</sup>,  
Anders M. Dale, PhD<sup>3,12</sup>,  
Tyler M. Seibert, MD, PhD<sup>2,3,12,13</sup>, for the PRACTICAL Consortium

<sup>1</sup> Division of Radiation Oncology, George Washington University, Washington, DC, USA

<sup>2</sup> Department of Radiation Medicine and Applied Sciences, University of California San Diego, La Jolla, CA, USA

<sup>3</sup> Center for Multimodal Imaging and Genetics, University of California, San Diego, La Jolla, CA, USA

<sup>4</sup> Division of Biostatistics and Halicioğlu Data Science Institute, University of California San Diego, La Jolla, CA, USA

<sup>5</sup> Department of Family Medicine and Public Health, University of California San Diego

<sup>6</sup> Division of Population Health, Health Services Research and Primary Care, University of Manchester, Oxford Road, Manchester, M13 9PL, UK

<sup>7</sup> Warwick Medical School, University of Warwick, Coventry, UK

<sup>8</sup> Unit of Cardiovascular and Nutritional Epidemiology, Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden

<sup>9</sup> Department of Surgical Sciences, Uppsala University, Uppsala, Sweden

<sup>10</sup> Nuffield Department of Surgical Sciences, University of Oxford, Oxford, UK

<sup>11</sup> NORMENT, KG Jebsen Centre, Oslo University Hospital and University of Oslo, Oslo, Norway

<sup>12</sup> Department of Radiology, University of California San Diego, La Jolla, CA, USA

<sup>13</sup> Department of Bioengineering, University of California San Diego, La Jolla, CA, USA

**Short running title:** Genetics/clinical risk factors for prostate cancer

**Key words:** prostate cancer; genetics; precision medicine; survival; risk factors

### **Corresponding Author:**

Tyler M. Seibert, MD, PhD

Assistant Professor

Department of Radiation Medicine and Applied Sciences

Department of Bioengineering

Department of Radiology  
University of California San Diego  
9500 Gilman Dr. Mail Code 0861  
La Jolla, CA 92093-0861  
[tseibert@ucsd.edu](mailto:tseibert@ucsd.edu)

### **Funding**

This study was funded in part by a grant from the United States National Institute of Health/National Institute of Biomedical Imaging and Bioengineering (#K08EB026503), United States Department of Defense (#W81XWH-13-1-0391), the Research Council of Norway (#223273), KG Jebsen Stiftelsen, and South East Norway Health Authority. The content is solely the responsibility of the authors and does not necessarily represent the official views of any of the funding agencies, who had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

### **Conflicts of Interest**

TM Seibert reports honoraria from Multimodal Imaging Services Corporation for imaging segmentation and honoraria from Varian Medical Systems and WebMD, Inc. for educational content. AM Dale has additional disclosures outside the present work: founder, equity holder, and advisory board member for CorTechs Labs, Inc.; advisory board member of Human Longevity, Inc.; recipient of nonfinancial research support from General Electric Healthcare. CC Fan is a scientific consultant for CorTechs Labs, Inc.

### **Data & Code Availability**

We obtained our dataset for this work from the Prostate Cancer Association Group to Investigate Cancer Associated Alterations in the Genome (PRACTICAL) consortium, which consists of a collaborative group of researchers, each of whom retains ownership of their contributed data. Members of the consortium can use pooled data via proposals that are reviewed by the Data Access Committee. Readers interested in participating in the PRACTICAL consortium and gaining access to member data may find information about application at: <http://practical.icr.ac.uk>. Code used for this work are stored in an institutional repository and will be shared upon request to the corresponding author.

## **Abstract**

Background: Clinical variables—age, family history, genetics—are used for prostate cancer risk stratification. Recently, polygenic hazard scores (PHS46, PHS166) were validated as associated with age at prostate cancer diagnosis. While polygenic scores are associated with all prostate cancer (not specific for fatal cancers), PHS46 was also associated with age at prostate cancer death. We evaluated if adding PHS to clinical variables improves associations with prostate cancer death.

Methods: Genotype/phenotype data were obtained from a nested case-control Cohort of Swedish Men (n=3,279; 2,163 with prostate cancer, 278 prostate cancer deaths). PHS and clinical variables (family history, alcohol intake, smoking, heart disease, hypertension, diabetes, body mass index) were tested via univariable Cox proportional hazards models for association with age at prostate cancer death. Multivariable Cox models with/without PHS were compared with log-likelihood tests.

Results: Median age at last follow-up/prostate cancer death were 78.0 (IQR: 72.3-84.1) and 81.4 (75.4-86.3) years, respectively. On univariable analysis, PHS46 (HR 3.41 [95%CI 2.78-4.17]), family history (HR 1.72 [1.46-2.03]), alcohol (HR 1.74 [1.40-2.15]), diabetes (HR 0.53 [0.37-0.75]) were each associated with prostate cancer death. On multivariable analysis, PHS46 (HR 2.45 [1.99-2.97]), family history (HR 1.73 [1.48-2.03]), alcohol (HR 1.45 [1.19-1.76]), diabetes (HR 0.62 [0.42-0.90]) all remained associated with fatal disease. Including PHS46 or PHS166 improved multivariable models for fatal prostate cancer ( $p < 10^{-15}$ ).

Conclusions: PHS had the most robust association with fatal prostate cancer in a multivariable model with common risk factors, including family history. Adding PHS to clinical variables may improve prostate cancer risk stratification strategies.

## Introduction

Prostate cancer is a common malignancy that causes significant morbidity and mortality in men worldwide<sup>1</sup>. Risk-stratified screening has shown promise for identification of men most likely to develop prostate cancer<sup>2–5</sup>. Currently, estimates of a man's risk may include clinical variables associated with prostate cancer, including family history, age, and race/ethnicity<sup>6–11</sup>. Other risk factors have also been reported<sup>12,13</sup>. Genetic factors may provide additional analytical and objective information<sup>2,14,15</sup>, possibly improving risk stratification beyond clinical variables alone.

A polygenic hazard score (PHS)—the weighted sum of approximately 50 single-nucleotide polymorphisms (SNPs)—has been developed and validated as independently associated with age at prostate cancer diagnosis. Though the association was not specific for fatal cancers, PHS was also associated with age at prostate cancer death<sup>2,16,17</sup>. The goal of the present work was to determine whether PHS improves stratification of risk for prostate cancer death beyond that achieved with clinical variables available in the Cohort of Swedish Men (COSM)<sup>18</sup>. COSM is a prospective, longitudinal, population-based study with participant-level demographic and clinical information. Data available include the development (or not) of prostate cancer and over two decades of mature follow-up data and death information<sup>18</sup>, affording an opportunity for comparison of genetic risk and clinical variables for associations with prostate cancer death.

## Methods

### The COSM Study

COSM is a longitudinal, population-based study designed to study associations of lifestyle with morbidity and mortality<sup>18</sup>. Men were surveyed at study onset in 1997, and in 2008

and 2009. Between 2005-2008, all the men still alive in the cohort were asked to give a saliva sample for genotyping through a self-administered postal collection<sup>19</sup>. A nested case-control study identified prostate cancer cases and controls. Prostate cancer cases included all the men with genotype data available and a diagnosis of prostate cancer made by the end of 2011. Men without prostate cancer at last follow-up were selected as controls, matched to the age distribution of the cases. There were 3,279 men (2,163 prostate cancer cases, 1,116 controls) in this COSM dataset. Prior work suggested this dataset would have adequate power for the present analyses<sup>20</sup>.

We identified the following clinical variables available in the COSM cohort: family history of prostate cancer (yes or no), alcohol intake (yes or no), smoking status (never smoker, current daily smoker, and ex-smoker), history of heart disease (yes or no), history of hypertension (yes or no), history of diabetes (clinically diagnosed by doctor; yes or no), and body mass index (BMI; kg/m<sup>2</sup>). Family history was taken as presence or absence of a first degree relative with a prostate cancer diagnosis. The number of relatives has also been associated with prostate cancer risk<sup>6,10,21</sup>, but there were not enough available data in the COSM dataset to include this variable in the present analysis.

### Polygenic Hazard Score (PHS)

A version of the original PHS with 46 SNPs<sup>2,16</sup> (PHS46) was previously shown to be independently associated with any, aggressive, and fatal prostate cancers in men of European, African, and Asian genetic ancestry in a heterogeneous, multi-ethnic dataset, which included the COSM data analyzed here. The combination of PHS46 and family history in multivariable models performed better than family history alone for all endpoints<sup>16</sup>.

Here, we tested the performance of the adapted PHS for association with age at prostate cancer death in the COSM dataset, specifically, taking advantage of the rich clinical information, long follow-up, and population-based design of the COSM study. PHS46 was calculated for all patients in COSM and used as the sole covariate in Cox proportional hazards regressions for age at prostate cancer death. Secondly, we also evaluated PHS46 association with age at any and aggressive prostate cancer diagnosis. As in prior reports<sup>2,16</sup>, aggressive prostate cancer was defined as disease typically requiring treatment: Gleason score  $\geq 7$ , clinical stage T3-T4, PSA concentration  $\geq 10\text{ng/mL}$ , or nodal/distant metastasis. Separate Cox proportional hazards regressions were performed for age at diagnosis of any or aggressive prostate cancer. All *p*-values reported were truncated at  $<10^{-16}$ <sup>2</sup>. As Cox proportional hazards results may be biased by a higher number of cases in the dataset compared to the general population, sample weight corrections were applied to all Cox proportional hazards models in this study using Swedish population data, as described previously<sup>2,16,22</sup>.

Recognizing that more SNPs associated with prostate cancer have been reported since the development of the original and adapted PHS<sup>23</sup>, a PHS model has also recently been developed that incorporates 166 total SNPs (PHS166)<sup>24</sup>, drawing from the SNPs in PHS46 and SNPs reported in a meta-analysis of genome-wide association studies<sup>23</sup>. PHS166 improved the hazard ratios (HRs) for prostate cancer over PHS46. The COSM data were included in the SNP selection and model weight estimation of PHS166, so there may be some within-sample overestimation of performance. With that caveat, we additionally evaluated PHS166 here for association with fatal prostate cancer in COSM in models accounting for clinical variables.

#### Clinical variables and PHS – association with fatal prostate cancer

Univariable Cox proportional hazards models were used to determine whether individual clinical variables were independently associated with age at prostate cancer death. If a participant did not have the clinical variable of interest available, they were not included in those specific Cox models. Those clinical variables with  $p < 0.05$  for univariable prediction of age of diagnosis of any prostate cancer were combined in a multivariable model. To evaluate whether addition of genetic risk improved prediction over clinical variables alone, a log-likelihood test compared the multivariable model with and without PHS46 as a predictor. Finally, the multivariable model with PHS46 was compared to the univariable model with PHS46 as the sole predictor of age of diagnosis of any prostate cancer, again by log-likelihood test. Significance for comparison of multivariable models was set at  $\alpha = 0.05$ .

The above multivariable analyses were repeated, with PHS166 used instead of PHS46.

#### Code Availability

The code used for this work are stored in an institutional repository and will be shared upon request to the corresponding author.

## Results

### COSM participants

**Table 1** describes the cohort. There were 278 deaths from prostate cancer, as well as 1,403 cases meeting clinical criteria for aggressive disease. The median age at last follow-up was 78 years (IQR: 72.3, 84.1), and the median age at prostate cancer death was 81.4 years (IQR: 75.4, 86.3).



## PHS in COSM

PHS46 was associated with age at prostate cancer death in the COSM dataset (HR 3.41 [95% CI 2.78, 4.17],  $z=12$ ,  $p<10^{-16}$ ). PHS46 was also associated with any (HR 5.35 [4.97, 5.76],  $z=45$ ,  $p<10^{-16}$ ) and aggressive (HR 3.78 [3.45, 4.13],  $z=29$ ,  $p<10^{-16}$ ) prostate cancer, consistent with prior findings that SNP associations are not specific for aggressive forms of cancer.

PHS166 was also associated with age at prostate cancer death (HR 7.15 [95% CI 5.95, 8.58],  $z=21$ ,  $p<10^{-15}$ ).

## Clinical variables and PHS – Association with fatal prostate cancer

On univariable analysis, family history of prostate cancer (HR 1.72 [95% CI 1.46, 2.03],  $z=6.5$ ,  $p<10^{-9}$ ), alcohol intake (HR 1.74 [1.40, 2.15],  $z=5.1$ ,  $p<10^{-6}$ ), and diabetes history (HR 0.53 [0.37, 0.75],  $z=-3.6$ ,  $p=0.003$ ) were each associated with prostate cancer death; these were included in the multivariable models. Smoking history (HR 0.96 [0.84, 1.10],  $z=-0.6$ ,  $p=0.56$  for ex-smokers; HR 0.92 [0.79, 1.08],  $z=-1.0$ ,  $p=0.34$  for current daily smokers), heart disease (HR 0.97 [0.83, 1.13],  $z=-0.4$ ,  $p=0.68$ ), hypertension (HR 1.06 [0.93, 1.21],  $z=0.9$ ,  $p=0.30$ ), and BMI (HR 0.99 [0.97, 1.01],  $z=-0.8$ ,  $p=0.45$ ) were not associated with prostate cancer death.

In the multivariable model with all clinical variables (with univariable significance), each variable remained independently associated with prostate cancer death. Family history of prostate cancer (HR 1.70 [1.44, 2.01],  $z=6.3$ ,  $p<10^{-9}$ ) and history of alcohol intake (HR 1.44 [1.17, 1.78],  $z=3.5$ ,  $p=0.0005$ ) were associated with increased risk of prostate cancer death. Diabetes history (HR 0.63 [0.43, 0.92],  $z=-2.4$ ,  $p=0.014$ ) was associated with decreased risk of prostate cancer death.

Adding PHS46 improved multivariable model performance. PHS46, family history, alcohol intake, and diabetes history each remained independently significant in the multivariable model, with PHS (HR 2.45 [1.99, 2.97],  $z=8.6$ ,  $p<10^{-15}$ ) and family history (HR 1.73 [1.48, 2.03],  $z=6.5$ ,  $p<10^{-10}$ ) having the strongest associations with prostate cancer death (**Table 2**). A log-likelihood test confirmed model improvement with inclusion of genetic risk, compared to only clinical variables ( $p<10^{-15}$ ). Similarly, the full multivariable model was significantly better than PHS alone (log-likelihood test,  $p<10^{-10}$ ).

Multivariable models with PHS166 were also improved with addition of PHS166. PHS166, family history, and alcohol intake each remained independently significant in the multivariable model, with PHS166 (HR 5.48 [4.54, 6.61],  $z=17.9$ ,  $p<10^{-15}$ ) having the strongest association with prostate cancer death (**Table 3**). Diabetes history was not independently significant in the multivariable model using PHS166. A log-likelihood test confirmed model improvement with inclusion of genetic risk, compared to only clinical variables ( $p<10^{-15}$ ). Similarly, the full multivariable model was significantly better than PHS alone (log-likelihood test,  $p<10^{-7}$ ).

## Discussion

In a nested case-control study from a large, population-based, longitudinal Swedish cohort<sup>18</sup>, we found that a multivariable model accounting for both clinical and genetic factors best stratified men for risk of fatal prostate cancer. Specifically, the addition of PHS significantly improved associations for age at prostate cancer death over that of the clinical variables alone. PHS46 had the most robust effect in the multivariable model, after controlling for other clinical variables in the COSM dataset. Incorporating more SNPs, PHS166 improved on the performance

of PHS46 and yielded a similar pattern of genetics as an important risk factor for fatal prostate cancer. Our work suggests that a multifactorial approach combining genetics and medical history might be useful for prostate cancer risk stratification.

In the present analysis, multiple clinical variables were associated with fatal prostate cancer. We found that family history of prostate cancer, alcohol intake, and a lack of diabetes were associated with increased likelihood of prostate cancer death. These results are consistent with previous investigations into prostate cancer risk factors. Family history of prostate cancer was strongly and independently associated with prostate cancer death. This is consistent with prior reports and with the current use of family history in clinical risk stratification<sup>6,7,10,11,21,25,26</sup>. In fact, on multivariable analysis with PHS46, family history yielded the second highest hazard ratio (1.73) for prostate cancer death, second only to that of PHS46 (2.45). Prior epidemiological studies have also reported associations between positive family history and fatal prostate cancer<sup>27-29</sup>, though this association has not been universal across studies<sup>29</sup>, and the use of family history in screening and treatment decisions may complicate interpretation. It is possible that there are hereditary components of prostate cancer risk transmitted through family trees that lead to strong associations with fatal disease; age of the affected first-degree relative may also impact this risk<sup>27,28</sup>. These components may have led to us finding associations between family history and fatal prostate cancer using data from the COSM study. However, family history does not appear to be an adequate surrogate for an individual's genetic risk, as PHS46 was also independently associated with prostate cancer death, indicating PHS and family history offer complementary risk information.

The other clinical variables associated with fatal prostate cancer identified in the present work were alcohol use and diabetes history. Alcohol intake was associated with increased risk of

fatal disease, while diabetes was correlated with decreased risk of fatal disease. Alcohol consumption has been linked with prostate cancer risk in a recent meta-analysis<sup>12</sup>, but other studies have not found such a relationship<sup>30,31</sup>. Suggested mechanisms of alcohol contributing to carcinogenesis include cellular DNA repair changes, oxidative stress/damage, and changes in metabolism<sup>32</sup>. It is possible that the associations we found between alcohol use and fatal prostate cancer reflect some amount of generalized alcohol-induced carcinogenesis. It is also possible that uncontrolled confounding factors have led to the finding. Diabetes history, on the other hand, may reduce a man's risk of prostate cancer<sup>33-35</sup>. This might possibly be related to the use of anti-diabetic medications such as metformin, which has been hypothesized to have a protective effect, though the present work does not address causality<sup>36</sup>.

While our present findings are consistent with previous epidemiological studies of prostate cancer risk, this is the first study to specifically assess for association with age at prostate cancer death<sup>12,25,33</sup>. Age is a critical consideration in prostate cancer decision-making<sup>37</sup>, especially as medical co-morbidities and competing risk of death increase with aging<sup>38</sup>. Prostate cancer incidence also rises exponentially with age and is typically more aggressive when diagnosed at an older age<sup>39,40</sup>. Interpreting prostate cancer risk stratification in the context of age is therefore important for clinical translation.

The present work is limited to clinical variables available in the COSM dataset. Other prostate cancer risk factors have been reported previously, including the number of affected family members in those men with a family history of prostate cancer<sup>6,10,21</sup>. Additionally, this is a study of a single population. While the PHS has been validated in a multi-ethnic dataset for association with fatal prostate cancer<sup>16</sup>, the relationships reported here between clinical variables and PHS may not generalize to other populations and warrant further study<sup>41</sup>. The focus of this

study was the important endpoint of prostate cancer death, but, consistent with prior work, germline SNP associations here are not specific to fatal or aggressive prostate cancer. Finally, the performance of PHS166 may have been influenced by incorporation of COSM data in the prior development of the PHS166 model (done via a LASSO-regularized Cox regression framework)<sup>24</sup>; however, potential bias is likely low, given that COSM data represent less than 4% of the data used in that study. Despite these limitations, our results indicate that polygenic scores like PHS may significantly improve strategies to identify men at highest (or lowest) risk of dying from prostate cancer.

## **Conclusions**

PHS and clinical variables afford complementary information regarding risk of fatal prostate cancer. In a multivariable model, PHS, family history, alcohol intake, and history of diabetes were each independently associated with prostate cancer death. PHS had the largest hazard ratio for fatal prostate cancer among these variables. Combining genetic and clinical factors should be studied as a strategy to improve risk-stratification for prostate cancer screening.

## **Ethics Statement**

All contributing studies were approved by the relevant ethics committees and performed in accordance with the Declaration of Helsinki; written informed consent was obtained from the study participants. The present analyses used de-identified data from the PRACTICAL consortium and have been approved by the review board at the corresponding authors' institution.

## References

1. Torre LA, Bray F, Siegel RL, Ferlay J, Lortet-Tieulent J, Jemal A. Global cancer statistics, 2012. *CA Cancer J Clin*. 2015;65(2):87-108. doi:10.3322/caac.21262
2. Seibert TM, Fan CC, Wang Y, et al. Polygenic hazard score to guide screening for aggressive prostate cancer: Development and validation in large scale cohorts. *BMJ*. 2018;360:1-7. doi:10.1136/bmj.j5757
3. Shoag JE, Nyame YA, Gulati R, Etzioni R, Hu JC. Reconsidering the trade-offs of prostate cancer screening. *N Engl J Med*. 2020;382(25):2465-2468. doi:10.1056/NEJMs2000250
4. Loeb S, Lilja H, Vickers A. Beyond prostate-specific antigen: Utilizing novel strategies to screen men for prostate cancer. *Curr Opin Urol*. 2016;26(5):459-465. doi:10.1097/MOU.0000000000000316
5. Ström P, Nordström T, Aly M, Egevad L, Grönberg H, Eklund M. The Stockholm-3 Model for Prostate Cancer Detection: Algorithm Update, Biomarker Contribution, and Reflex Test Potential. *Eur Urol*. 2018;74(2):204-210. doi:10.1016/j.eururo.2017.12.028
6. Steinberg GD, Carter BS, Beaty TH, Childs B, Walsh PC. Family history and the risk of prostate cancer. *Prostate*. 1990;17(4):337-347. doi:10.1002/pros.2990170409
7. Bratt O, Drevin L, Akre O, Garmo H, Stattin P. Family History and Probability of Prostate Cancer, Differentiated by Risk Category: A Nationwide Population-Based Study. *J Natl Cancer Inst*. 2016;108(10):djw110. doi:10.1093/jnci/djw110
8. Rawla P. Epidemiology of Prostate Cancer. *World J Oncol*. 2019;10(2):63-89. doi:10.14740/wjon1191
9. Baquet CR, Horm JW, Gibbs T, Greenwald P. Socioeconomic factors and cancer

- incidence among blacks and whites. *J Natl Cancer Inst.* 1991;83(8):551-557.  
doi:10.1093/jnci/83.8.551
10. Albright F, Stephenson RA, Agarwal N, et al. Prostate cancer risk prediction based on complete prostate cancer family history. *Prostate.* 2015;75(4):390-398.  
doi:10.1002/pros.22925
  11. Giri VN, Beebe-Dimmer JL. Familial prostate cancer. *Semin Oncol.* 2016;43(5):560-565.  
doi:10.1053/j.seminoncol.2016.08.001
  12. Zhao J, Stockwell T, Roemer A, Chikritzhs T. Is alcohol consumption a risk factor for prostate cancer? A systematic review and meta-analysis. *BMC Cancer.* 2016;16(1):845.  
doi:10.1186/s12885-016-2891-z
  13. Giovannucci E, Liu Y, Platz EA, Stampfer MJ, Willett WC. Risk factors for prostate cancer incidence and progression in the health professionals follow-up study. *Int J Cancer.* 2007;121(7):1571-1578. doi:10.1002/ijc.22788
  14. Witte JS. Personalized prostate cancer screening: Improving PSA tests with genomic information. *Sci Transl Med.* 2010;2(62):62ps55. doi:10.1126/scitranslmed.3001861
  15. Taylor RA, Fraser M, Rebello RJ, et al. The influence of BRCA2 mutation on localized prostate cancer. *Nat Rev Urol.* 2019;16(5):281-290. doi:10.1038/s41585-019-0164-8
  16. Huynh-Le M-P, Fan CC, Karunamuni R, et al. Polygenic hazard score is associated with prostate cancer in multi-ethnic populations. *medRxiv.* 2019:19012237.  
doi:10.1101/19012237
  17. Karunamuni R, Huynh-Le M-P, Fan C, et al. African-specific improvement of a polygenic hazard score for age at diagnosis of prostate cancer. *Int J Cancer.* 2020.  
doi:10.1002/ijc.33282



18. Discacciati A, Orsini N, Andersson S-O, et al. Coffee consumption and risk of localized, advanced and fatal prostate cancer: a population-based prospective study. *Ann Oncol*. 2013;24(7):1912-1918. doi:10.1093/annonc/mdt105
19. Rylander-Rudqvist T, Håkansson N, Tybring G, Wolk A. Quality and quantity of saliva DNA obtained from the self-administrated oragene method - A pilot study on the cohort of Swedish men. *Cancer Epidemiol Biomarkers Prev*. 2006;15(9):1742-1745. doi:10.1158/1055-9965.EPI-05-0706
20. Karunamuni RA, Huynh-Le M-P, Fan CC, et al. The effect of sample size on polygenic hazard models for prostate cancer. *Eur J Hum Genet*. June 2020. doi:10.1038/s41431-020-0664-2
21. Abdel-Rahman O. Prostate Cancer Incidence and Mortality in Relationship to Family History of Prostate Cancer; Findings From The PLCO Trial. *Clin Genitourin Cancer*. 2019. doi:10.1016/j.clgc.2019.05.015
22. Therneau TM, Li H. Computing the Cox Model for Case Cohort Designs. *Lifetime Data Anal*. 1999;5(2):99-112. doi:10.1023/A:1009691327335
23. Schumacher FR, Al Olama AA, Berndt SI, et al. Association analyses of more than 140,000 men identify 63 new prostate cancer susceptibility loci. *Nat Genet*. 2018;50(7):928-936. doi:10.1038/s41588-018-0142-8
24. Karunamuni RA, Huynh-Le M-P, Fan CC, et al. Additional SNPs improve the performance of a polygenic hazard score for prostate cancer. *medRxiv*. 2020. doi:10.1101/2020.09.11.20188383
25. Wolf AMD, Wender RC, Etzioni RB, et al. American Cancer Society Guideline for the Early Detection of Prostate Cancer: Update 2010. *CA Cancer J Clin*. 2010;60(2):70-98.

doi:10.3322/caac.20066

26. Goldgar DE, Easton DF, Cannon-albright LA, Skolnick MH. Systematic population-based assessment of cancer risk in first-degree relatives of cancer probands. *J Natl Cancer Inst.* 1994;86(21):1600-1608. doi:10.1093/jnci/86.21.1600
27. Rodriguez C, Calle EE, Miracle-McMahill HL, et al. Family history and risk of fatal prostate cancer. *Epidemiology.* 1997;8(6):653-657. doi:10.1097/00001648-199710000-00007
28. Hemminki K. Familial risk and familial survival in prostate cancer. *World J Urol.* 2012;30(2):143-148. doi:10.1007/s00345-011-0801-1
29. Barber L, Gerke T, Markt SC, et al. Family history of breast or prostate cancer and prostate cancer risk. *Clin Cancer Res.* 2018;24(23):5910-5917. doi:10.1158/1078-0432.CCR-18-0370
30. Adami HO, McLaughlin JK, Hsing AW, et al. Alcoholism and cancer risk: a population-based cohort study. *Cancer Causes Control.* 1992;3(5):419-425. doi:10.1007/BF00051354
31. Brunner C, Davies NM, Martin RM, et al. Alcohol consumption and prostate cancer incidence and progression: A Mendelian randomisation study. *Int J Cancer.* 2017;140(1):75-85. doi:10.1002/ijc.30436
32. Ratna A, Mandrekar P. Alcohol and cancer: Mechanisms and therapies. *Biomolecules.* 2017;7(3). doi:10.3390/biom7030061
33. Fall K, Garmo H, Gudbjörnsdottir S, Stattin P, Zethelius B. Diabetes mellitus and prostate cancer risk; A nationwide case-control study within PCBaSe Sweden. *Cancer Epidemiol Biomarkers Prev.* 2013;22(6):1102-1109. doi:10.1158/1055-9965.EPI-12-1046
34. Bonovas S, Filioussi K, Tsantes A. Diabetes mellitus and risk of prostate cancer: A meta-

- analysis. *Diabetologia*. 2004;47(6):1071-1078. doi:10.1007/s00125-004-1415-6
35. Beckmann K, Crawley D, Nordström T, et al. Association Between Antidiabetic Medications and Prostate-Specific Antigen Levels and Biopsy Results. *JAMA Netw open*. 2019;2(11):e1914689. doi:10.1001/jamanetworkopen.2019.14689
  36. Hankinson SJ, Fam M, Patel NN. A review for clinicians: Prostate cancer and the antineoplastic properties of metformin. *Urol Oncol Semin Orig Investig*. 2017;35(1):21-29. doi:10.1016/j.urolonc.2016.10.009
  37. NCCN Clinical Practice Guidelines in Oncology. Prostate Cancer. Version 1.2019.
  38. *NCCN Guidelines Version 1.2019 Older Adult Oncology*.; 2019.
  39. Huynh-Le M-P, Fan CC, Karunamuni R, et al. A Genetic Risk Score to Personalize Prostate Cancer Screening, Applied to Population Data. *Cancer Epidemiol Biomarkers Prev*. 2020;29(9):1731-1738. doi:10.1158/1055-9965.epi-19-1527
  40. Huynh-Le MP, Myklebust TÅ, Feng CH, et al. Age dependence of modern clinical risk groups for localized prostate cancer—A population-based study. *Cancer*. 2020;126(8):1691-1699. doi:10.1002/cncr.32702
  41. DeSantis CE, Siegel RL, Sauer AG, et al. Cancer statistics for African Americans, 2016: Progress and opportunities in reducing racial disparities. *CA Cancer J Clin*. 2016;66(4):290-308. doi:10.3322/caac.21340

**Table 1.** Participant characteristics, n=3,279.

<b><u>Participants</u></b>	<b><u>Number</u></b>
Controls	1,116
Prostate cancer cases	2,163
Aggressive prostate cancer cases <sup>a</sup>	1,403
Fatal prostate cancer cases	278
<b><u>Age Demographics</u></b>	<b><u>Median (IQR)</u></b> <sup>b</sup>
Age at prostate cancer diagnosis	72.4 (66.2, 78.5)
Age at last follow up	78.0 (72.3, 84.1)
Age at prostate cancer death	81.4 (75.4, 86.3)
<b><u>Polygenic Hazard Scores</u></b>	<b><u>Median (IQR)</u></b>
PHS46 overall	0.4863 (0.2976, 0.6949)
PHS46 in fatal prostate cancer cases	0.5035 (0.3306, 0.7051)
PHS46 in non-fatal prostate cancer cases	0.4850 (0.2968, 0.6935)
PHS166 overall	0.0148 (-0.2306, 0.2508)
PHS166 in fatal prostate cancer cases	0.0770 (-0.1548, 0.2957)
PHS166 in non-fatal prostate cancer cases	0.0042 (-0.2380, 0.2472)
<b><u>Participants with Known Clinical Variables Information</u></b>	<b><u>Number</u></b>
Family history of prostate cancer	2,453 355 with positive family history
History of heart disease	3,279 382 with history of heart disease
History of hypertension	3,279 2,163 with history of hypertension
Alcohol intake	3,268 3,026 with reported alcohol intake
Current smoking status	3,237 1,310 never smokers 1,304 ex-smokers 623 current daily smokers
History of diabetes (clinically diagnosed by doctor)	3,279 120 with history of diabetes
Body mass index	3,132

<sup>a</sup> Aggressive prostate cancer defined as Gleason score  $\geq 7$ , PSA  $\geq 10$  ng/mL, T3-T4 stage, nodal metastases, or distant metastases.

<sup>b</sup> IQR: interquartile range

**Table 2.** Multivariable Cox proportional hazards model with both the polygenic hazard score (PHS46) and all clinical variables with significant univariable association with prostate cancer death.

<b>Clinical variable</b>	<b><math>\beta</math></b>	<b>HR (95% CI)</b>	<b>z-score</b>	<b><i>p</i>-value</b>
PHS46	0.90	2.45 (1.99, 2.97)	8.7	$<10^{-15}$
Family history of prostate cancer	0.55	1.73 (1.48, 2.03)	6.5	$<10^{-10}$
Alcohol intake	0.37	1.45 (1.19, 1.76)	3.5	0.0004
Diabetes	-0.48	0.62 (0.42, 0.90)	-2.5	0.012

**Table 3.** Multivariable Cox proportional hazards model with PHS166 and all clinical variables with significant univariable association with prostate cancer death.

<b>Clinical variable</b>	<b><math>\beta</math></b>	<b>HR (95% CI)</b>	<b>z-score</b>	<b><i>p</i>-value</b>
PHS166	1.70	5.48 (4.54, 6.61)	17.9	$<10^{-15}$
Family history of prostate cancer	0.39	1.48 (1.25, 1.75)	4.5	$<10^{-5}$
Alcohol intake	0.42	1.52 (1.22, 1.88)	3.9	$<10^{-4}$
Diabetes	-0.31	0.73 (0.49, 1.07)	-1.6	0.11