

Utility values for childhood obesity interventions: a systematic review and meta-analysis of the evidence for use in economic evaluation

Abbreviations:

CUA: cost-utility analysis. QALY: quality-adjusted life year. HRQoL: health-related quality of life. MAUIs: multi-attribute utility instruments. PRISMA: Preferred Reporting Items for Systematic Reviews and Meta-Analyses. PROSPERO: International Prospective Register of Systematic Reviews. PEDE: Pediatric Economic Database Evaluation. BMI: body mass index. LFK: Luis Furuya-Kanamori. WMD: weighted mean difference. 95% UI: 95% uncertainty interval. CHEERS: Consolidated Health Economic Evaluation Reporting Standards. AQoL-6D: Assessment of Quality of Life 6D. CHU-9D: child health utility index 9D. EQ-5D-Y: Euroqol 5D youth. EQ-5D-3L: Euroqol 5D 3L. HUI: health utilities index. UK: United Kingdom. US: United States of America. CEA: cost-effectiveness analysis. PedsQL: Pediatric quality of life inventory™ version 4.0 generic core scales.

Utility values for childhood obesity interventions: a systematic review and meta-analysis of the evidence for use in economic evaluation

Abstract

Rigorous estimates of preference-based utilities are important inputs into economic evaluations of childhood obesity interventions, yet no published review currently exists examining utility by weight status in paediatric populations. A comprehensive systematic literature review and meta-analysis was therefore undertaken, pooling data on preference-based health state utilities by weight status in children using a random effects model. Tests for heterogeneity were performed and publication bias was assessed. Of 3,434 potentially relevant studies identified, eleven met our eligibility criteria. Estimates of Cohen's d statistic suggested a small effect of weight status on preference-based utilities. Mean utility values were estimated as 0.85 (95% UI 0.84-0.87), 0.83 (95% UI 0.81-0.85), 0.82 (95% UI 0.79-0.84) and 0.83 (95% UI 0.80-0.86) for healthy weight, overweight, obese and overweight/obese states respectively. Meta-analysis of studies reporting utility values for both healthy weight and overweight/obese participants found a statistically significant weighted mean difference (0.015, 95% UI 0.003-0.026). A small but statistically significant difference was also estimated between healthy weight and overweight participants (0.011, 95% UI 0.004-0.018). Study findings suggest that paediatric specific benefits of obesity interventions may not be well reflected by available utility measures, potentially underestimating cost-effectiveness if weight loss in childhood/adolescence improves health or well-being.

1. Introduction

Childhood overweight and obesity is a serious global public health challenge (1). Given society's scarce resources, rigorous economic evaluation of childhood obesity prevention and treatment interventions is a useful tool for priority-setting. Cost-utility analysis (CUA) is a form of economic evaluation that compares the costs of an intervention with the benefits gained, incorporating intervention impact on both the quality and quantity of life. CUA allows for comparison of cost-effectiveness across healthcare programs, and is used as a tool for allocating health resources by governments, including in Australia and the United Kingdom (2, 3).

The quality adjusted life year (QALY) is the most widely used utility-based unit of measurement for CUA (4). The use of QALYs allows for comparison between different health states or health outcomes within diverse populations (for example, both adult and child populations), with an underlying assumption that QALYs are of equal social value irrespective of who accrues them (5). Preference-based health-related quality of life (HRQoL) is incorporated using health utilities that are indexed on a cardinal scale where 0 represents death and 1 represents perfect health (6). Utilities are preference weights, with preference equated with value or desirability (7). Utilities can be measured directly (using techniques such as Standard Gamble or Time Trade Off) or indirectly (using Multi-Attribute Utility Instruments (MAUIs))(8). The choice of utility method and measure can have an effect on estimated utility values, and in turn on cost-effectiveness results (9).

Catalogues of utility values can help to inform the appropriate selection of values for use in economic modelling. Health economists are often not equipped with the time or resources to elicit utility values either directly or indirectly themselves. Reviews of utility values for use in adult populations across health states such as diabetes, cancer and mental disorders exist (9-13). Evidence from such reviews improves the robustness, transparency and rigour of modelled economic evaluations, allowing for a more systematic approach to the selection of appropriate model parameters. For example, the catalogue of weights by Tengs & Wallace (12) was used to inform the economic modelling of long-term health outcomes and economic consequences of intervention for diabetes mellitus (14). The more recent study by Davies et al. (15) incorporated utility values from the review by Beaudet et al. (9) to investigate the cost-effectiveness of different therapies for patients with type 2 diabetes mellitus.

Limited reviews have examined utility values by weight status in adults (13, 16). Measuring utilities for children and adolescents is however only an emerging field of research, given the complexities of valuation in these age groups (17). In particular, whilst MAUIs have gained popularity as a straightforward and easy way to elicit values, most MAUIs have historically been applied in adult populations (18). Limited catalogues or reviews of utility values in child or adolescent populations currently exist (19-22). Additionally, no published reviews or catalogues of utility values for use in CUAs of childhood and adolescent obesity interventions have been identified. To date and to the best of our knowledge, studies undertaking meta-analysis of obesity-related HRQoL in paediatric populations have focused on non-preference based measures (23, 24).

This study aims to systematically review studies reporting utility values by weight status in children and adolescents aged ≤ 18 years. A meta-analysis was undertaken, providing a catalogue of utility values useful for informing CUAs of obesity interventions in paediatric populations. In addition, this study reviews published economic evaluations of obesity interventions in paediatric populations that have measured or referenced utility-based HRQoL. Health state utility values are often reported in economic evaluations, rather than solely in health benefit measurement studies (25). This ensures an extensive scope for the search for utility values by weight status, and also provides an overview of the current state of practice and context for how published utility values are used in the economic modelling of paediatric obesity interventions.

2. Methods

The systematic review was based on the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines (26) and registered with PROSPERO (#CRD42017067226). A search of the literature was conducted using key search strategies agreed in conjunction with a subject specialist librarian (Supporting information).

To be considered for inclusion, studies had to meet the following criteria:

- Studies published in peer-reviewed journals reporting primary data collection of utility values, with results reported by weight status;
- Conducted in paediatric populations (mean or median target age ≤ 18 years);
- Written in the English language;

- Published anytime until May 2017.

Reviews or commentaries were excluded. A search was also conducted for economic evaluations of obesity interventions published in peer-reviewed journals including an analysis of HRQoL that used or referenced a preference-based utility instrument. Economic evaluations needed to include paediatric populations within their study population, but did not have to be limited to paediatric populations. Utility values needed to specifically relate to weight status, rather than the potential health sequelae of obesity that can also appear in economic models that adopt long-term time horizons. The Paediatric Economic Database Evaluation (PEDE) (27) was also searched, using the search terms “obes*”, “body mass index” and “BMI” as keyword/title/abstract search terms. Screening of potential study inclusions was conducted by two authors (VB, ET), with conflicts resolved in consultation with a third author (MM). Reference lists of all included studies were checked for further inclusions (i.e. backward citation searching).

Data were extracted by one reviewer (VB) using Covidence (28) and Microsoft Excel, and checked by a second reviewer (ET). Key characteristics of interest for studies eliciting utility values included study aim, methods, study population, sample size, intervention evaluated (where applicable), utility measure, tariff (where applicable), and QALY weights. Data on weight classification of the study population were extracted as per the BMI cutoff values used in each selected study. Meta-analysis was undertaken with the Microsoft Excel add-in MetaXL version 5.3 (29) using a random effects model. Mean utility values for healthy weight, overweight, obese and overweight/obese (i.e. studies combining overweight and obese into a single category) states were estimated, ignoring potential within-study correlation in estimates because of the small numbers of studies included (30). Utility values from relevant studies were included as separate observations in the meta-analyses. Forest plots were presented, and tests for heterogeneity were performed using I^2 and Cochran’s Q test. Heterogeneity was regarded as substantial when I^2 exceeded 40% or the Q statistic was significant at $p < 0.10$ (31). Formal quality assessment of primary studies eliciting utility values was not undertaken, given a lack of standard systems or checklists for grading the quality of health state utility values (25, 32). We examined potential publication and small study bias visually using funnel and Doi plots, where a symmetrical plot suggests no or little bias (33). The Luis Furuya-Kanamori (LFK) index of asymmetry is also presented from the Doi plot, with an assessment of “no”, “minor” or “major” asymmetry (33).

Studies reporting mean utility values for both healthy weight and overweight/obese participants were also pooled to estimate the weighted mean difference (WMD) in utility values and 95% uncertainty interval (95% UI). Sub-analyses were undertaken for studies reporting mean utility values for both healthy weight and either overweight or obese participants. Cohen's d effect sizes were estimated, where approximately 0.2, 0.5 and 0.8 are respectively considered small, moderate and large (34).

A number of sensitivity analyses were undertaken to test the robustness of meta-analysis results. Firstly, individual studies were sequentially excluded from all meta-analyses to gauge the effect on overall results (35) (Supporting information). Variability in meta-analysis estimates may result from differences in the descriptive systems, valuation protocols and utility ranges of the utility measures. Therefore sensitivity analyses were also undertaken restricting study inclusion in the mean utility and WMD healthy weight and overweight/obese meta-analyses to (i) studies using the EQ-5D-Y (36, 37) and (ii) studies using the CHU-9D (37-39). Sensitivity analyses for WMD between healthy weight and either overweight or obese states were only estimated for studies using the CHU-9D instrument (37-39), due to the small number of included studies using other instruments. Variability in meta-analysis results may also result from the use of different BMI cutoff reference values between studies (Supporting information). Sensitivity analyses were therefore conducted, stratifying by BMI cutoff reference values. Sensitivity analyses were also conducted to stratify study inclusions by age, by pooling studies whose mean study participant age was <10 years or ≥10 years.

Key study variables of interest for economic evaluations of paediatric obesity interventions included those reported in the Consolidated Health Economic Evaluation Reporting Standards (CHEERS) checklist (40). Economic evaluation studies were summarised narratively, focusing on methodology and use of preference-based utility values within the analyses.

3. Results

3.1 Selection and inclusion of studies

After the removal of duplicates, the systematic search resulted in the identification of 3,434 potentially relevant publications (Figure 1). Titles and abstracts were searched by two independent reviewers, following which 3,106 studies were excluded because they did not meet the inclusion criteria. Full texts for the remaining 328 potentially relevant studies were located and searched, with 317 of these studies

excluded because they did not meet the inclusion criteria. No additional papers were identified from the reference lists of included studies or the PEDE database. A total of 11 studies were included in our review, nine health state valuation studies eliciting primary utility values (36-39, 41-45) and two economic evaluations (46, 47).

Figure 1 – PRISMA flowchart of systematic review results

3.2 Study characteristics- primary health state valuation studies eliciting utility values

Health state valuation studies eliciting utility values were undertaken in Australia (38, 42, 43), England (36, 37, 39), the USA (41, 45) and the Netherlands (44). All studies were cross-sectional, with seven studies reporting data collected at single time points within larger intervention studies (37-39, 42-45). Study settings included clinic or specialist care centres (41, 44), primary or middle schools (37, 39, 45), secondary schools (36, 42, 43) or a mix of both primary and secondary schools as part of a community-based intervention (38). The mean age of study participants varied between 6.3 years (39) and 15.7 years (44) (Supporting information). Only one study used parental proxy report for utility (with self-report also undertaken in participants aged ≥ 8 years) (41).

Utility instruments used are reported in Table 1. As might be expected, the most common utility instruments were those developed specifically for use in children and adolescents, with the CHU-9D used in three studies (37-39) and the EQ-5D-Y used in two studies (36, 37). One study (36) collected height and weight outcomes by self-report, with all other studies using objectively measured data. Studies used different weight classification BMI cutoff values (Supporting information), and weight classification categories varied between studies (healthy weight, overweight, obese, severe obesity, overweight/obese). Only three included studies reported that utility data was non-normally distributed (38, 39, 41), with two studies specifically reporting the data as left-skewed (39, 41); only one study reported no underlying differences in the distribution of the utility data in different weight categories (39).

Table 1 – Utility instruments from included studies

Table 2 lists the published mean utility values by weight status from studies included in our review. The study by Boyle et al. (36) invited participants to complete the study survey on two occasions (in winter

and in summer) but analysed the data cross-sectionally. It is unclear how many of the children completed the survey at both time points. Three studies used more than one utility instrument (37, 41, 45); however, only the studies by Canaway & Frew 2014 (37) and Trevino et al. 2013 (45) reported complete results per instrument used.

Table 2 – Published utility weights by weight status from the literature

3.3 Results of meta-analyses

Results from our systematic review demonstrate that there is a relatively wide range of utility values by weight status reported within the literature (range reported for healthy weight 0.73-0.93, overweight 0.66-0.86, obese 0.69-0.84 and overweight/obese 0.67-0.89) (Table 2). Figure 2 reports the forest plots for mean utility values by weight status. The mean utility values were 0.85 (95% UI 0.84-0.87, $I^2=97$, $Q=424$, $p=0.00$), 0.83 (95% UI 0.81-0.85, $I^2=89$, $Q=56.1$, $p=0.00$), 0.82 (95% UI 0.79-0.84, $I^2=86$, $Q=42.1$, $p=0.00$) and 0.83 (95% UI 0.8-0.86, $I^2=87$, $Q=55.5$, $p=0.00$) for healthy weight, overweight, obese and overweight/obese states, respectively. Given that only two studies reported utility values for the severe obesity weight category (40-41), a separate mean utility value for the severe obesity state was not estimated. All meta-analyses displayed a moderate level of heterogeneity.

Figure 2 – Forest plots of mean utility by weight status

After pooling relevant results, the weighted mean difference in utility value between healthy weight and overweight/obese participants was small but statistically significant (WMD 0.015, 95% UI 0.003-0.026, $I^2=9.3$, $Q=7.7$, $p=0.36$) (Table 2). An even smaller but still statistically significant weighted mean difference in utility value was found between healthy weight and overweight participants (WMD 0.011, 95% UI 0.004-0.018, $I^2=0$, $Q=1.6$, $p=0.95$). The WMD between healthy weight and obese participants was slightly larger; however, results suggest heterogeneity exists (0.024, 95% UI 0.009-0.039, $I^2=60.2$, $Q=15.1$, $p=0.02$). Visual inspection of funnel and Doi plots also indicated the possibility of publication bias for all analyses. LFK Indexes for all analyses reported major asymmetry (Table 3).

Table 3 – Pooled estimates of WMD in HRQoL in children and adolescents, by weight category

Sensitivity analyses were undertaken for all analyses and full results are presented in Supporting information files. Omitting individual studies within the analyses of mean utility values for healthy weight, overweight, obese and overweight/obese states did not change results markedly. When considering all analyses, the mean utility value for healthy weight ranged from 0.82 (95% UI 0.78-0.87, when including only studies where the mean participant age was <10 years (37, 39)) to 0.88 (95% UI 0.83-0.93, when including only studies using the EQ-5D-Y (36, 37)). Mean utility values for the overweight/obese state ranged from 0.81 (95% UI 0.76-0.87, when only studies with a mean participant age <10 years were included (37, 39)) to 0.84 (95% UI 0.81-0.88, when only studies with a mean participant age ≥10 years were included (36, 38, 41, 42) or when individual studies were excluded from the analyses (37, 38, 41)).

The sequential exclusion of the study by Bolton et al. (42) in the WMD analysis (healthy weight and overweight/obese), resulted in the WMD estimate no longer reaching statistical significance at the 5% level (WMD 0.014, 95% UI 0-0.029, $I^2=17$, $Q=7.3$). Restriction of inclusions in the analysis of WMD between healthy weight and overweight/obese to studies using the UK BMI cutoff reference values (60) resulted in the WMD estimate no longer reaching statistical significance at the 5% level (WMD 0.019, 95% UI -0.004-0.043, $I^2=43$, $Q=7$). Restrictions of inclusions into the analysis of WMD between healthy weight and overweight/obese to studies using either the EQ-5D-Y (36, 37) or the CHU-9D (37-39) resulted in WMD in health utility no longer reaching statistical significance (Supporting information), although it should be noted that these analyses were limited by the small number of study inclusions by instrument. Similar findings resulted from restricting inclusions into healthy weight and either overweight or obese analyses to those studies using the CHU9D (healthy weight and overweight: WMD 0.011, 95% UI -0.001-0.022, $I^2=0$, $Q=0.07$; healthy weight and obese: WMD 0.019, 95% UI -0.014-0.051, $I^2=61$, $Q=5.1$)(Supporting information).

3.4 *Summary of economic evaluations*

Only two economic evaluations of obesity interventions were identified as including an analysis of weight-related HRQoL using or referencing a utility-based instrument (Supporting information). The study by McAuley et al. (46) evaluated a community-based obesity prevention intervention and utilised the Health Utilities Index (HUI) to collect parental proxy HRQoL data for a sample of New Zealand children aged 5 to 12 years. No statistically significant differences between the intervention and control

groups were reported, and utility by weight status was therefore not reported. The economic evaluation consisted of a cost-effectiveness analysis (CEA), where results were presented as cost per kilogram of weight gain prevented from the intervention.

The study by Robertson et al. (47) evaluated the effectiveness and cost-effectiveness of a community-based programmatic intervention. Data on utility values were collected at baseline, three and twelve months using the EQ-5D-Y based on both child self-report and parental proxy. Although utility values were not specifically reported, a statistically significant change in HRQoL was not found in study participants between intervention timepoints. Among children with complete cost and QALY data over the study period, the intervention was associated with a mean incremental QALY gained of only 0.0009.

4. Discussion

Many of the conditions for which obesity is a risk factor, such as stroke, heart disease and type 2 diabetes, generally occur later in life (61, 62). HRQoL may therefore be the most measurable impact of overweight and obesity in the childhood years, and this is an important consideration when undertaking CUA of obesity interventions in paediatric populations. Given the potential influence on cost-effectiveness results, the selection of appropriate health state utility values is important. Therefore estimates of utility associated with weight status arising from meta-analyses are useful in improving the rigour and reliability of CUA findings, as well as providing an overview of the current state of evidence for the impact of overweight and obesity on preference-based HRQoL in children and adolescents.

Studies in adult populations have estimated lower utility-based HRQoL between healthy weight and overweight or obese states (63). For instance, the study by Sach et al. (64) estimated a statistically significant mean difference of -0.099 between healthy weight and obese class I states using the EQ-5D instrument. A broad systematic review and meta-analysis of childhood health utilities by Kwon et al. has very recently been published (22). However, the breadth of that review did not allow the authors to focus on weight status in children in the granulated way that we do. For example, Kwon et al. (22) estimate weighted averages of mean utility or VAS scores for broad ICD-10 delineated categories that grouped obese status and type II diabetes together. Moreover, Kwon et al. do not estimate utility score differences between children categorised as either healthy weight, overweight, obese and

overweight/obese (22). Our review therefore presents a more in-depth and granulated assessment of utility values by weight status in children and adolescents aged ≤ 18 years.

Our systematic review demonstrated that the current literature on utility values for children and adolescents by weight status is limited. There is also a relatively wide range of values estimated. The results of our meta-analyses displayed high degrees of heterogeneity; however, pooled analyses suggest that in children and adolescents, overweight and obesity states are associated with lower utility values than those for a healthy weight. This is important information for health economists and those interested in priority setting, with the catalogue of preference-based utility values presented here useful as inputs into economic evaluations of obesity interventions in paediatric populations. Whilst we do not recommend these values as definitive given the limited evidence base, our results highlight the potential for impact of interventions that are effective at changing weight status through the childhood years.

Utility measurement in childhood populations is clearly an underexplored area, with more research required to better understand the clinical importance and significance of HRQoL in children and adolescents. We found only 11 relevant studies for our review, although several other studies measured utility values, but were not included in our analyses because they did not report scored utility values (65-68) or ascribe them to weight categories (69). Reviews of non-preference based studies assessing HRQoL in childhood populations suggest poorer HRQoL in obese youth (24, 70-72), with recent studies finding that interventions that reduce BMI are associated with improvements in HRQoL (73, 74). Our meta-analysis results broadly support these findings, however given this is a relatively emerging field within the literature, it is clear that more research is required to better understand potential quantitative associations useful for economic modelling. Our meta-analysis results are also broadly consistent with the findings of other studies reporting an element of ill-health or suboptimal HRQoL in the general population (75). Our estimated utility value for healthy weight children (mean 0.85 (95% UI 0.84-0.87)) is consistent with the findings from Kwon et al. (22), who reported a mean utility using the HUI3 of 0.876 (95% UI 0.788-0.965).

A hypothesis within the literature is that very young children do not experience any loss of HRQoL due to weight status, but that older children and adolescents may (37). We identified only two studies estimating utilities for children under 10 years (37, 39), and so our sensitivity analyses estimating mean utilities in younger populations were limited by the lack of study inclusions. Results demonstrated

overlapping of confidence intervals in utility values between younger children (i.e. <10 years) of healthy weight (0.82 (95% UI 0.78-0.87)) and those that were overweight/obese (0.81 (95% UI 0.76-0.87)). Whilst this may suggest either no or minimal effect of weight status on HRQoL in younger children, it is clear that more research is required for a better understanding of the potential differential effects across the childhood and adolescent years. This is important because if the HRQoL decrements between weight states are small, interventions aimed at reducing obesity in children are only likely to be cost-effective based on QALYs if they are either very low cost or if the intervention effects are able to be sustained through to adulthood, thereby reducing the incidence of chronic diseases in later life. Debate exists within the literature on the appropriateness of using QALY outcome measures in cost-utility analysis for children, rather than cost-effectiveness analysis where benefits are expressed in terms of natural units (for example, cost per BMI unit saved)(76).

Many studies conducted in paediatric populations use non-preference based measures such as the Paediatric Quality of Life Inventory™ Version 4.0 Generic Core Scales (PedsQL™)(77). Non-preference based measures can be mapped or cross-walked onto generic preference-based measures of health; however, there are current limitations in doing so in paediatric populations (20, 78). The current state of the literature allows for mapping from the PedsQL to the CHU9D in adolescents (78) or to the EQ-5D-Y in children aged 11-15 years (79). More research is required to extend these and other mapping algorithms across different populations and age groups.

Measuring child and adolescent HRQoL is challenging, and there are well-recognised limitations to existing methods (17). Accurately capturing information on HRQoL in children requires instruments that are sensitive to detect differences and appropriate to deliver across different age groups. For instance, whilst younger children in particular may require parental proxy reporting, the accuracy of such reporting has recently been questioned with UI-Haq et al. (24) finding that parents may overestimate the impact of obesity on the HRQoL of their children. In addition, the tariffs for scoring utility values for many instruments may not necessarily reflect the values of childhood or adolescent populations if elicited from general populations.

The findings of our systematic review and meta-analysis highlight that HRQoL is a significant source of uncertainty in the economic modelling of obesity interventions in childhood populations. The current paucity of published utility values by weight status is reflected in the lack of published cost-utility

analyses of paediatric obesity interventions incorporating HRQoL in childhood. Many cost-effectiveness analyses of paediatric obesity interventions focus on averting adult obesity, and only include utility values to estimate QALYs in adults (for example, (80, 81)). This may not allow for the HRQoL benefits of an intervention during childhood to be considered (41), and may underestimate intervention cost-effectiveness if overweight or obese states themselves have a significant impact on HRQoL in childhood and adolescence. The recent study by Lal et al. (82) was published after our literature search was completed, but modelled the cost-effectiveness of a sugar-sweetened beverage tax across socioeconomic position using utility values for healthy weight and obesity in children and adolescents sourced from Chen et al. (38). This may reflect the growing interest in trying to capture the full range of costs and benefits of obesity interventions in childhood and adolescence, particularly when disease-specific consequences are largely experienced later in life.

The strengths of our study include the comprehensive and systematic search strategies, as well as the undertaking of a number of sensitivity analyses. This allowed for a range of summary estimates to be produced, for inclusion within economic evaluations at the discretion of study authors given the relative uncertainty within the body of literature.

Our study has some limitations, which should be borne in mind by readers. The small number of studies included in our meta-analyses means that less confidence may be placed in results, although we present all values here as a quantitative summary of the literature as it currently stands and recommend use with caution. The published evidence on which we have based our results has also only been undertaken in Western developed countries, and therefore the generalisability of these results to countries with differing cultural associations around body weight is unknown. Whilst our results build upon the currently limited evidence base on the impact of HRQoL by weight status in paediatric populations, it is clear that further investigation is required.

We also did not conduct quality assessments of studies included in our meta-analyses, and therefore cannot comment on the rigour or reliability of study inclusions. The lack of standard systems or validated checklists for grading the quality of health state utility values for inclusion in meta-analysis is both a limitation and an area for future research (32, 83). Due to the small number of study inclusions and the lack of a validated quality assessment tool, we assumed that the quality across all included studies was of an equal standard, and this may in fact not have been the case. This in turn may

influence results of the meta-analyses, however we have tried to circumvent this through extensive sensitivity analyses omitting individual studies from the analyses in order to gauge overall effects on utility estimates. It is also possible that publication bias may influence these results, with studies finding no statistically significant difference in preference-based utility perhaps less likely to be published than studies that report a difference, and this should be considered.

Results have also been pooled across studies using different utility instruments, BMI cutoff values and tariffs for scoring, in different study populations. The use of meta-analytical methods to pool utility values from different instruments is an ongoing area for investigation, with more research required into the sources of variation in utility values that may be driven by differences in descriptive systems (84). We have again attempted to circumvent this by undertaking several sensitivity analyses, although the low number of studies using each respective instrument or by age group was a limitation. Finally, our systematic review and meta-analysis identified only cross-sectional studies examining HRQoL by weight status. It is possible that reverse causation may be a factor in any association, and given the current evidence base we are unable to establish whether BMI might impact on the HRQoL of children and adolescents or vice versa. More rigorous investigation is therefore required, exploring causation and the complexity of possible associations through multiple pathways.

5. Conclusion

This study represents the first systematic review and meta-analysis of utility values by weight status for childhood populations, demonstrating the relative lack of published preference-based utility values by weight status in paediatric populations. To date, very few economic evaluations of obesity interventions incorporating HRQoL benefits to children and adolescents have been published. This potentially results in under-estimation of the cost-effectiveness of obesity interventions in children and adolescents. Meta-analysis results provide mean utility values by healthy weight, overweight, obese and overweight/obese states, although results should be interpreted with caution given high degrees of heterogeneity. Results demonstrate higher preference-based utility values in healthy weight as compared to overweight or obese children and adolescents, but more evidence is required to improve the rigour and reliability of these estimates.

Figure 1 – PRISMA flowchart of systematic review results

Table 1 – Utility instruments from included studies

Table 2 – Published utility weights by weight status from the literature

Figure 2 – Forest plots of mean utility by weight status

Table 3 – Pooled estimates of WMD in HRQoL in children and adolescents, by weight category

Conflict of Interest Statement

There are no conflicts of interest to declare.

REFERENCES

1. World Health Organisation. Report of the Commission on Ending Childhood Obesity. Geneva: WHO, 2016.
2. Australian Government Department of Health. Guidelines for preparing submissions to the Pharmaceutical Benefits Advisory Committee (PBAC) Canberra, Australia: Australian Government; n.d. [cited 2017 14 August]. Available from: <https://pbac.pbs.gov.au/>.
3. National Institute for Health and Care Excellence. Developing NICE guidelines: the manual United Kingdom 2017 [cited 2017 14 August]. Available from: <https://www.nice.org.uk/process/pmg20/chapter/incorporating-economic-evaluation>.
4. Robinson R. Cost-utility analysis. *BMJ*. 1993;307(6908):859-62.
5. Whitehead SJ, Ali S. Health outcomes in economic evaluation: the QALY and utilities. *Br Med Bull*. 2010;96(1):5-21.
6. Drummond MF, Sculpher MJ, Torrance GW, O'Brien BJ, Stoddart GL. *Methods for the Economic Evaluation of Health Care Programmes*. 3rd edn ed. Oxford: Oxford University Press; 2005.
7. Weinstein MC, Torrance G, McGuire A. QALYs: the basics. *Value Health*. 2009;12(s1):S5-S9.
8. Torrance GW. Preferences for health outcomes and cost-utility analysis. *Am J Manag Care*. 1997;3(Suppl 1):S8-S20.
9. Beaudet A, Clegg J, Thuresson P-O, Lloyd A, McEwan P. Review of utility values for economic modeling in type 2 diabetes. *Value Health*. 2014;17(4):462-70.
10. Sonntag M, König H-H, Konnopka A. The estimation of utility weights in cost-utility analysis for mental disorders: a systematic review. *Pharmacoeconomics*. 2013;31(12):1131-54.
11. Brown J, Cook K, Adamski K, Lau J, Bargo D, Breen S, et al. Utility values associated with advanced or metastatic non-small cell lung cancer: data needs for economic modeling. *Expert Rev Pharmacoecon Outcomes Res*. 2017;17(2):153-64.
12. Tengs TO, Wallace A. One thousand health-related quality-of-life estimates. *Med Care*. 2000;38(6):637-647.
13. Dennett SL, Boye KS, Yurgin NR. The Impact of Body Weight on Patient Utilities with or without Type 2 Diabetes: A Review of the Medical Literature. *Value Health*. 2008;11(3):478-86.
14. Palmer AJ, Roze S, Valentine WJ, Minshall ME, Foos V, Lurati FM, et al. The CORE Diabetes Model: projecting long-term clinical outcomes, costs and costeffectiveness of interventions in diabetes mellitus (types 1 and 2) to support clinical and reimbursement decision-making. *Curr Med Res Opin*. 2004;20(sup1):S5-S26.
15. Davies MJ, Glah D, Chubb B, Konidaris G, McEwan P. Cost effectiveness of IDegLira vs. alternative basal insulin intensification therapies in patients with type 2 diabetes mellitus uncontrolled on basal insulin in a UK setting. *Pharmacoeconomics*. 2016;34(9):953-66.
16. Doyle S, Lloyd A, Moore L, Ray J, Gray A. A systematic review and critical assessment of health state utilities: weight change and type 2 diabetes mellitus. *Pharmacoeconomics*. 2012;30(12):1133-43.
17. Thorrington D, Eames K. Measuring health utilities in children and adolescents: A systematic review of the literature. *PLoS One*. 2015;10(8).
18. Chen G, Ratcliffe J. A Review of the Development and Application of Generic Multi-Attribute Utility Instruments for Paediatric Populations. *Pharmacoeconomics*. 2015;33(10):1013-28.
19. Petrou S. Methodological issues raised by preference-based approaches to measuring the health status of children. *Health Econ*. 2003;12(8):697-702.
20. Petrou S, Kupek E. Estimating Preference-Based Health Utilities Index Mark 3 Utility Scores for Childhood Conditions in England and Scotland. *Med Decis Making*. 2009;29(3):291-303.
21. Herdman M, Cole A, Hoyle CK, Coles V, Carroll S, Devlin N. Sources and Characteristics of Utility Weights for Economic Evaluation of Pediatric Vaccines: A Systematic Review. *Value Health*. 2016;19(2):255-66.

22. Kwon J, Kim SW, Ungar WJ, Tsiplova K, Madan J, Petrou S. A Systematic Review and Meta-Analysis of Childhood Health Utilities. *Med Decis Making*. 2017. In press.
23. Kelly A, Ligthart M, Paulis WD, Djasmo D, Koes BW, van Middelkoop M. Effect of multidisciplinary interventions on quality of life in obese children: A systematic review and meta-analysis. *Qual Life Res*. 2015;24(7):1635.
24. Ul-Haq Z, Mackay DF, Fenwick E, Pell JP. Meta-analysis of the association between body mass index and health-related quality of life among children and adolescents, assessed using the pediatric quality of life inventory index. *J Pediatr*. 2013;162(2):280-6.e1.
25. Papaioannou D, Brazier J, Paisley S. Systematic searching and selection of health state utility values from the literature. *Value Health*. 2013;16(4):686-95.
26. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*. 2009;6(7):e1000097.
27. The Research Institute at The Hospital For Sick Children. Paediatric Economic Database Evaluation (PEDE) Toronto, Canada: The Hospital for Sick Children; 2017 [cited 2017 12 July]. Available from: <http://pede.ccb.sickkids.ca/pede/search.jsp>.
28. Covidence. Covidence 2017 [cited 2017 14 August]. Available from: <https://www.covidence.org/>.
29. EpiGear International. MetaXL Queensland, Australia: EpiGear International; 2017 [cited 2017 3 July]. Available from: http://www.epigear.com/index_files/metaxl.html.
30. Lung TW, Hayes AJ, Hayen A, Farmer A, Clarke PM. A meta-analysis of health state valuations for people with diabetes: explaining the variation across methods and implications for economic evaluation. *Qual Life Res*. 2011;20(10):1669-78.
31. Higgins J, Green S, editors. *Cochrane handbook for systematic reviews of interventions*, version 5.1: The Cochrane Collaboration 2011; 2011.
32. Doyle S, Lloyd A, Moore L, Ray J, Gray A. A systematic review and critical assessment of health state utilities. *Pharmacoeconomics*. 2012;30(12):1133-43.
33. Barendregt JJ, Doi SA. *MetaXL User Guide*. Queensland, Australia: EpiGear International, 2016.
34. Cohen J. The effect size index: d. *Statistical power analysis for the behavioral sciences*. 1988;2:284-8.
35. Le LK-D, Barendregt JJ, Hay P, Mihalopoulos C. Prevention of eating disorders: A systematic review and meta-analysis. *Clin Psychol Rev*. 2017;53:46-58.
36. Boyle SE, Jones GL, Walters SJ. Physical activity, quality of life, weight status and diet in adolescents. *Qual Life Res*. 2010;19(7):943-54.
37. Canaway AG, Frew EJ. Is utility-based quality of life in children aged 6-7 years affected by Body Mass Index (BMI)? *Int J Obesity* (2005). 2014;38(8):1146.
38. Chen G, Ratcliffe J, Olds T, Magarey A, Jones M, Leslie E. BMI, health behaviors, and quality of life in children and adolescents: a school-based study. *Pediatr*. 2014;133(4):e868-e74.
39. Frew EJ, Pallan M, Lancashire E, Hemming K, Adab P. Is utility-based quality of life associated with overweight in children? Evidence from the UK WAVES randomised controlled study. *BMC Pediatr*. 2015;15(211).
40. Husereau D, Drummond M, Petrou S, Carswell C, Moher D, Greenberg D, et al. Consolidated health economic evaluation reporting standards (CHEERS) statement. *Cost Eff Resour Alloc*. 2013;11(1):6.
41. Belfort MB, Zupancic JAF, Riera KM, Turner JHG, Prosser LA. Health state preferences associated with weight status in children and adolescents. *BMC Pediatr*. 2011;11(12).
42. Bolton K, Kremer P, Rossthorn N, Moodie M, Gibbs L, Waters E, et al. The effect of gender and age on the association between weight status and health-related quality of life in Australian adolescents. *BMC Public Health*. 2014;14(898).
43. Keating CL, Moodie ML, Richardson J, Swinburn BA. Utility-based quality of life of overweight and obese adolescents. *Value Health*. 2011;14(5):752-8.

44. Makkes S, Renders CM, Bosmans JE, Baan-Slootweg OHvd, Seidell JC. Cardiometabolic risk factors and quality of life in severely obese children and adolescents in the Netherlands. *BMC Pediatr.* 2013;13(62).
45. Treviño RP, Pham TH, Edelstein SL. Obesity and preference-weighted quality of life of ethnically diverse middle school children: the HEALTHY study. *J Obesity.* 2013;2013.
46. McAuley KA, Taylor RW, Farmer VL, Hansen P, Williams SM, Booker CS, et al. Economic evaluation of a community-based obesity prevention program in children: the APPLE project. *Obesity.* 2010;18(1):131-6.
47. Robertson W, Fleming J, Kamal A, Hamborg T, Khan KA, Griffiths F, et al. Randomised controlled trial and economic evaluation of the 'Families for Health' programme to reduce obesity in children. *Arch Dis Childhood.* 2017;102(5):416-26.
48. Richardson JR, Peacock SJ, Hawthorne G, Iezzi A, Elsworth G, Day NA. Construction of the descriptive system for the assessment of quality of life AQoL-6D utility instrument. *Health Qual Life Outcomes.* 2012;10(1):38.
49. Moodie M, Richardson J, Rankin B, Iezzi A, Sinha K. Predicting time trade-off health state valuations of adolescents in four Pacific countries using the Assessment of Quality-of-Life (AQoL-6D) instrument. *Value Health.* 2010;13(8):1014-27.
50. The University of Sheffield. Measuring and Valuing Health, The development of a paediatric health related quality of life measure for use in economic evaluation: The Child Health Utility 9D (CHU 9D) United Kingdom: The University of Sheffield; 2017 [cited 2017 03 July]. Available from: <https://www.sheffield.ac.uk/scharr/sections/heds/mvh/paediatric>.
51. Stevens K. Valuation of the child health utility 9D index. *Pharmacoeconomics.* 2012;30(8):729.
52. Ratcliffe J, Flynn T, Terlich F, Stevens K, Brazier J, Sawyer M. Developing adolescent-specific health state values for economic evaluation. *Pharmacoeconomics.* 2012;30(8):713.
53. EuroQol. EQ-5D: About The Netherlands: EuroQol; 2017 [cited 2017 03 July]. Available from: <https://euroqol.org/eq-5d-instruments/eq-5d-y-about/>.
54. Dolan P, Gudex C, Kind P, Williams A. A social tariff for EuroQol: results from a UK general population survey. University of York: University of York, 1995.
55. EuroQol Research Foundation. EQ-5D-3L: About The Netherlands: EuroQol Research Foundation; 2017 [cited 2017 11 July]. Available from: <https://euroqol.org/eq-5d-instruments/eq-5d-3l-about/>.
56. Lamers L, Stalmeier P, McDonnell J, Krabbe P, van Busschbach J. Measuring the quality of life in economic evaluations: the Dutch EQ-5D tariff. *Nederlands tijdschrift voor geneeskunde.* 2005;149(28):1574-8.
57. Health Utilities Inc. Health Utility Index Canada: HUI Inc; 2017 [cited 2017 03 July]. Available from: <http://www.healthutilities.com/>.
58. Torrance GW, Feeny DH, Furlong WJ, Barr RD, Zhang Y, Wang Q. Multiattribute utility function for a comprehensive health status classification system: Health Utilities Index Mark 2. *Med Care.* 1996;34(7):702-22.
59. Feeny D, Furlong W, Torrance GW, Goldsmith CH, Zhu Z, DePauw S, et al. Multiattribute and single-attribute utility functions for the health utilities index mark 3 system. *Med Care.* 2002;40(2):113-28.
60. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ.* 2000;320(7244):1240.
61. Finkelstein EA, Brown DS, Trogdon JG, Segel JE, Ben-Joseph RH. Age-specific impact of obesity on prevalence and costs of diabetes and dyslipidemia. *Value Health.* 2007;10:S45-S51.
62. Ezzati M, Lopez AD, Rodgers A, Murray CJ. Comparative quantification of health risks: global and regional burden of disease attributable to selected major risk factors. Geneva: World Health Organisation; 2004.

63. Kolotkin R, Andersen J. A systematic review of reviews: exploring the relationship between obesity, weight loss and health-related quality of life. *Clin Obesity*. 2017;5:273-289.
64. Sach TH, Barton GR, Doherty M, Muir KR, Jenkinson C, Avery AJ. The relationship between body mass index and health-related quality of life: comparing the EQ-5D, EuroQol VAS and SF-6D. *Int J Obesity*. 2006;31:189.
65. Petracci E, Cavrini G. The effect of weight status, lifestyle, and body image perception on health-related quality of life in children: a quantile approach. *Qual Life Res*. 2013;22(9):2607-15.
66. Kesztyűs D, Wirt T, Kobel S, Schreiber A, Kettner S, Dreyhaupt J, et al. Is central obesity associated with poorer health and health-related quality of life in primary school children? Cross-sectional results from the Baden-Württemberg Study. *BMC Public Health*. 2013;13(260).
67. Kesztyűs D, Schreiber A, Kobel S, Wartha O, Kesztyűs T, Kilian R, et al. Illness and determinants of health-related quality of life in a cross-sectional sample of schoolchildren in different weight categories. *GMS*. 2014;12.
68. Wu X, Ohinmaa A, Veugelaers PJ. Diet quality, physical activity, body weight and health-related quality of life among grade 5 students in Canada. *Public Health Nutr*. 2012;15(1):75-81.
69. Kremer P, Waqa G, Vanualailai N, Schultz JT, Roberts G, Moodie M, et al. Reducing unhealthy weight gain in Fijian adolescents: results of the Healthy Youth Healthy Communities study. *Obesity Rev*. 2011;12(s2):29-40.
70. Griffiths LJ, Parsons TJ, Hill AJ. Self-esteem and quality of life in obese children and adolescents: a systematic review. *Int J Pediatr Obesity*. 2010;5(4):282-304.
71. Buttitta M, Iliescu C, Rousseau A, Guerrien A. Quality of life in overweight and obese children and adolescents: a literature review. *Qual Life Res*. 2014;23(4):1117-39.
72. Tsiros MD, Olds T, Buckley JD, Grimshaw P, Brennan L, Walkley J, et al. Health-related quality of life in obese children and adolescents. *Int J Obesity*. 2009;33(4):387-400.
73. Møllerup PM, Nielsen TR, Bøjsøe C, Kloppenborg JT, Baker JL, Holm J-C. Quality of life improves in children and adolescents during a community-based overweight and obesity treatment. *Qual Life Res*. 2017;26(6):1597-608.
74. Steele RG, Gayes LA, Dalton WT, Smith C, Maphis L, Conway-Williams E. Change in health-related quality of life in the context of pediatric obesity interventions: A meta-analytic review. *Health Psychol*. 2016;35(10):1097-109.
75. Richardson J, Khan MA, Iezzi A, Maxwell A. Comparing and explaining differences in the magnitude, content, and sensitivity of utilities predicted by the EQ-5D, SF-6D, HUI 3, 15D, QWB, and AQL-8D multiattribute utility instruments. *Med Decis Making*. 2015;35(3):276-91.
76. Griebisch I, Coast J, Brown J. Quality-adjusted life-years lack quality in pediatric care: a critical review of published cost-utility studies in child health. *Pediatr*. 2005;115(5):e600-e14.
77. Varni JW. The PedsQL Measurement Model for the Pediatric Quality of Life Inventory 2017 [cited 2017 10 July]. Available from: <http://www.pedsq.org/>.
78. Mpundu-Kaambwa C, Chen G, Russo R, Stevens K, Petersen KD, Ratcliffe J. Mapping CHU9D Utility Scores from the PedsQLTM 4.0 SF-15. *Pharmacoeconomics*. 2017;35(4):453-67.
79. Khan KA, Petrou S, Rivero-Arias O, Walters SJ, Boyle SE. Mapping EQ-5D utility scores from the PedsQL™ generic core scales. *Pharmacoeconomics*. 2014;32(7):693-706.
80. Gortmaker SL, Long MW, Resch SC, Ward ZJ, Cradock AL, Barrett JL, et al. Cost Effectiveness of Childhood Obesity Interventions: Evidence and Methods for CHOICES. *Am J Prev Med*. 2015;49(1):102-11.
81. Graziose MM, Koch PA, Wang YC, Lee Gray H, Contento IR. Cost-effectiveness of a Nutrition Education Curriculum Intervention in Elementary Schools. *J Nutr Education Behavior*. 2016;49(8):684-691.
82. Lal A, Mantilla-Herrera AM, Veerman L, Backholer K, Sacks G, Moodie M, et al. Modelled health benefits of a sugar-sweetened beverage tax across different socioeconomic groups in Australia: A cost-effectiveness and equity analysis. *PLOS Med* 2017;14(6):e1002326.

83. Nerich V, Saing S, Gamper E-M, Holzner B, Pivot X, Viney R, et al. Critical appraisal of health-state utility values used in breast cancer-related cost–utility analyses. *Breast Cancer Res Treatment*. 2017;1-10.
84. Peasgood T, Brazier J. Is meta-analysis for utility values appropriate given the potential impact different elicitation methods have on values? *Pharmacoeconomics*. 2015;33(11):1101-5.
85. Centers for Disease Control and Prevention. Defining Childhood Obesity USA: CDC; 2016 [cited 2017 05 July]. Available from: <https://www.cdc.gov/obesity/childhood/defining.html>.
86. World Health Organisation. Growth reference 5-19 years Geneva, Switzerland: WHO; 2017 [cited 2017 5 July]. Available from: <http://www.who.int/growthref/en/>.
87. Cole TJ, Freeman JV, Preece MA. Body mass index reference curves for the UK, 1990. *Arch Disease Childhood*. 1995;73(1):25-9.
88. Fredriks AM, Van Buuren S, Burgmeijer RJ, Meulmeester JF, Beuker RJ, Brugman E, et al. Continuing positive secular growth change in The Netherlands 1955–1997. *Pediatr Res*. 2000;47(3):316-23.