

1 Functional amyloids promote retention of public goods in bacteria

2 John B. Bruce¹, Stuart A. West¹ & Ashleigh S. Griffin¹

3 ¹Department of Zoology, University of Oxford, Oxford, UK

4 Keywords: Cooperation, Social evolution, *Pseudomonas*, Functional amyloid,
5 Pyoverdine,

6 **Abstract**

7 The growth and virulence of bacteria depends upon a number of factors that are
8 secreted into the environment. These factors can diffuse away from the producing
9 cells, to be either lost or utilized by cells that do not produce them (cheats).

10 Mechanisms that act to reduce the loss of secreted factors through diffusion are
11 expected to be favoured. One such mechanism may be the production of Fap fibrils,
12 needle-like fibres on the cell surface observed in *P.aeruginosa*, which can transiently
13 bind several secreted metabolites produced by cells. We test whether Fap fibrils help
14 retain a secreted factor, the iron-scavenging molecule pyoverdine, and hence reduce
15 the potential for exploitation by non-producing, cheating cells. We found that: (1)
16 wildtype cells retain more iron-chelating metabolites than fibril non-producers; (2)
17 purified Fap fibrils can prevent the loss of the iron-chelators PQS (*Pseudomonas*
18 quinolone signal) and pyoverdine; and (3) pyoverdine non-producers have higher
19 fitness in competition with fibril non-producers than with wildtype cells. Our results
20 suggest that by limiting the loss of a costly public good, Fap fibrils may play an
21 important role in stabilizing cooperative production of secreted factors.

Introduction

Bacteria often rely upon the secretion of extracellular public goods to survive and thrive in an environment. These goods are metabolically costly to produce but may be essential for effective communication between cells, scavenging of iron and nutrients, protection from biotic and abiotic stresses and engaging in antagonistic competition (West & Buckling. 2003, Ghequire & De Mot 2014, Popat *et al.* 2015, Flemming *et al.* 2016,). Production of these goods can also benefit neighbouring cells. Thus, public goods producers can potentially be exploited by non-producing ‘cheats’ that benefit from the available goods while paying none of the metabolic costs of production (Griffin *et al.* 2004, Diggle *et al.* 2007a, Ghoul *et al.* 2014). Exploitation by cheats can be detrimental to the population; significantly decreasing its density and subsequently increasing its chances of collapse (Griffin *et al.* 2004, Sanchez & Gore 2013). Cheats have been observed to arise *de novo* under laboratory conditions, and putative social cheats have been identified in a number of natural bacterial populations (Sandoz *et al.* 2007, Cordero *et al.* 2012, West *et al.* 2012, Andersen *et al.* 2015, Bruce *et al.* 2017).

As well as exploitation by social cheats, cells may lose return on investment in public good production simply through diffusion or advection of costly goods away from the cell in turbulent environments (Kummerli *et al.* 2009, Emge *et al.* 2016, Mund *et al.* 2016). These losses can be minimized through a number of different mechanisms. Cells can avoid exploitation by cheats by ensuring the benefits of public good production accrue to relatives. This can be achieved at its simplest through clonal growth but bacteria can also engage in conflict with, and kill, local unrelated cells (Joulu *et al.* 2013, Nadell *et al.* 2013, McNally *et al.* 2016). Forming a physical barrier can also prevent loss of public goods by slowing diffusion and/or reducing fluid flow rate around the colony. For example, thick layers of biofilm produced by *Vibrio cholera* can slow the diffusion of digested chitin, preventing its use by protease-deficient mutants (Drescher *et al.* 2014).

A recently described mechanism that may slow or prevent the loss of secreted goods is the production of needle-like fibrils known collectively as functional amyloids (Dueholm *et al.* 2010). These fibrils are assembled on the cell surface and are thought to be involved in surface attachment, providing structural rigidity to the extracellular matrix and even aiding in the evasion of host immune responses (Barnhart & Chapman 2006, Zeng *et al.* 2015, Taglialegna *et al.* 2016). Recently, it

was demonstrated that functional amyloids produced by *Pseudomonas aeruginosa*, Fap fibrils, are capable of binding several hydrophobic metabolites secreted by the cell such as pyocyanin and the signaling molecule PQS (*Pseudomonas* quinolone signal) (Dueholm *et al.* 2013, Seviour *et al.* 2015). Binding of the metabolites is transient: raising the possibility that serial association and dissociation with Fap fibrils may allow secreted metabolites to be locally ‘bioavailable’ while ensuring they are retained close to producing cells. While the social consequences of metabolite binding by Fap fibrils have yet to be described, if public goods producers are able to maintain preferential access to the costly goods they secrete into the environment, this could have important implications for cooperator-cheat dynamics in bacterial populations (Ross Gillespie *et al.* 2007).

We test whether Fap fibrils help cells retain secreted metabolites involved in iron acquisition and if this can influence cooperator-cheat dynamics. Using strains of *Pseudomonas aeruginosa* that differ in their ability to produce Fap fibrils, we assess whether production of fibrils influences growth and investment in the iron-scavenging molecule pyoverdine. We also isolate Fap fibrils from cells to determine whether fibrils can bind, and prevent the loss by dilution, of the dedicated siderophore pyoverdine and the signaling molecule PQS, which is also a strong iron-chelator. We subsequently test whether any iron-chelating metabolites retained by Fap fibrils are functionally available to cells. Finally, we determine whether producing Fap fibrils influences the ability of pyoverdine producers to resist invasion by pyoverdine non-producing cheats in an iron-limited environment.

Material and Methods

Media and Bacterial Strains

To culture from frozen, we used Kings Broth (KB) medium (20g Protease Peptone N°3 (BD Biosciences), 10g glycerol, 1.5g K₂HPO₄·3H₂O and 1.5g MgSO₄·7H₂O per litre of dH₂O) and for growth in an iron-limited environment we used Casamino Acid (CAA) medium (5g Casamino acids, 1.18g K₂HPO₄·3H₂O and 0.25g MgSO₄·7H₂O per litre of dH₂O) supplemented with 20mM sodium bicarbonate and 100µg ml⁻¹ of the iron chelator human apo-transferrin. We washed and diluted bacterial cultures using M9 minimal salt media (6.8g Na₂HPO₄, 3g KH₂PO₄, 0.5g NaCl and 10 g NH₄Cl per litre of dH₂O). For purification of Fap fibrils, we cultured strains in colony factor

antigen (CFA) medium (10g hydrolyzed casein, 50mg MgSO₄ and 1.5g yeast extract per litre of dH₂O). All reagents were purchased from Sigma unless otherwise stated.

We used strains that differ in their ability to produce Fap fibrils and the siderophore pyoverdine in our experiments (Table 1). The PAO1 strain produces wildtype levels of Fap fibrils and pyoverdine. PAO1ΔFap is a mutant with the genes involved in Fap fibril production, FapA-FapF, inactivated by allelic replacement with a gentamicin resistance cassette (Dueholm *et al.* 2013). We also use a mutant that cannot produce the iron-scavenging siderophore pyoverdine. PAO1ΔpvdD is a clean deletion mutant defective for the pyoverdine synthetase gene pvdD (Ghysels *et al.* 2004). For isolating Fap fibrils, we use a PAO1 strain harboring a plasmid containing the *Pseudomonas* Fap operons under the control of an isopropyl β-D-1-thiogalactopyranoside (IPTG) inducible lacUV5 promoter (PAO1 pFap). Prior to our experiments, we cultured strains overnight for 15 hours at 37°C in 6ml of KB media, at 200rpm on an orbital shaker.

Table 1. Summarising phenotypic differences between strains used in the study.

Strain	Fap Fibrils	Pyoverdine
PAO1	+	+
PAO1ΔFap	-	+
PAO1 pFap	+	+
PAO1ΔPvdD	+	-

Measuring Growth and Pyoverdine Production

To determine whether producing Fap fibrils influenced the growth of strains, or altered their investment in pyoverdine, we cultured fibril producers and non-producers under iron-limited conditions. We did not quantify the production of pyochelin, the other siderophore produced by *P.aeruginosa*, as its production is almost completely repressed in the stringently iron-limited conditions used here (Dumas *et al.* 2013). We washed overnight cell cultures in M9 media, standardized to an optical density (A₆₀₀) of 0.2 and diluted 100-fold before inoculating 200μl of iron-limited CAA media with 2μl of diluted culture in a 96-well plate. Cell density (absorbance at 600nm) and pyoverdine production (Relative Fluorescent Units, using an excitation of 400nm and emission of 460nm respectively) were measured every 30 minutes for 24 hours using

a SpectraMax i3x multi-mode platform (Molecular Devices). We also calculated pyoverdine available per cell at each time point ($\text{RFU}_{400-460} / A_{600}$) and the rate of pyoverdine production per cell per minute ($(\text{RFU}_{400-460(2)} - \text{RFU}_{400-460(1)}) / 30 \text{ min} / A_{600(2)}$). Each treatment was replicated 6 times per strain.

Cells retention of iron-chelating metabolites

To determine whether Fap fibrils influence cells ability to retain iron-chelating metabolites, we measured the iron-chelating activity of fibril producers and non-producers after removing unbound metabolites, using a CAS assay (Schwyn & Neilands. 1987). Overnight cell cultures in M9 media were washed, standardized to an optical density (A_{600}) of 0.2 and diluted 100-fold before inoculating 6ml of iron-limited CAA media with 60 μl of diluted culture. We incubated vials statically for 48 hours at 37°C before vortexing and passing 2ml of culture through a 0.22 μm syringe filter. Cells held on the membrane were washed to remove unbound iron-chelating metabolites by passing 20ml of M9 solution through the syringe filter and re-suspended by passing 2ml of M9 solution through the opposite end of the filter. 100 μl of CAS solution (Schwyn & Neilands 1987) was added to 100 μl of re-suspended cells in a 96-well plate before we incubated the plate in darkness for 30 minutes. Iron-chelating molecules remove iron from the CAS solution, changing the colour of the solution. We quantified this change by measuring absorbance at A_{630} . We also measured the optical density of the re-suspended cells (A_{600}) and calculated the iron-chelating activity per cell ($1 - A_{630} / A_{600}$). Each treatment was replicated 22 times.

Binding of iron-chelators to Fap fibrils.

We wished to determine if Fap fibrils can bind iron-chelating metabolites, and whether bound metabolites are functionally available to cells. Isolated fibrils were exposed to the iron-chelators PQS and pyoverdine and the iron-chelating activity of the fibrils after repeated washing to remove unbound molecules was quantified. We also assessed the ability of fibrils exposed to iron-chelators to promote the growth of a pyoverdine deficient mutant in iron-limited media.

Isolation of Fap fibrils

We inoculated PAO1 pFAP, an inducible Fap fibril overproducing strain, into KB media and incubated overnight (37°C, 200rpm) before inoculating 400ml of CFA

media with 4ml of the overnight culture in a 2L flask with 40µg/ml of tetracycline. Cultures were incubated at 37°C, 200rpm to an optical density (A_{600}) of ~0.5 before inducing expression of Fap fibrils with 1mM of IPTG and incubating for a further 8 hours. We harvested cells by centrifugation (28,000g, 30 mins, 20°C), resuspended cells in 30ml of buffer (10mM Tris-HCL, pH 8.0) and homogenized manually before adding 10ml of enzyme mix (0.4mg/ml RNaseA, 0.4mg/ml DNase I, 4mg/ml lysozyme, 4mM MgCL₂ and 0.4% Triton X-100). This mix was homogenized by three cycles of freeze-thawing using a -80°C freezer and a 37°C water bath, followed by a final incubation of 2hrs at 37°C. We then added 5ml of 20% sodium dodecyl sulfate (SDS), boiled the sample for 20 minutes and collected insoluble material by centrifugation (28,000g, 30 mins, 20°C). The pellet was then resuspended in 27ml buffer with 3ml of 20% SDS and boiled for 20 minutes before insoluble material was collected by centrifugation. We repeated this step twice. To wash away residual SDS, we resuspended the pellet in 25ml of buffer and centrifuged (28,000g, 30 mins, 20°C), repeating this step a further two times before re-suspending the pellet in 3ml of buffer after the final centrifugation.

To confirm the presence of amyloid structures in our purified protein solution, we used a Thioflavin T fluorescence assay. We suspended 0.5mg/ml of lysozyme and 0.5mg/ml of purified protein solution in 10mM Tris-HCl (pH 8.0) and 40µM Thioflavin T. 10mM Tris-HCl (pH 8.0) and lysozyme, a protein lacking any amyloid structures, served as negative controls. Using a Spectramax i3x, we recorded the emission spectra of the samples from 470-600nm after excitation at 450nm.

Fap fibril retention of iron-chelating metabolites

We suspended purified Fap fibrils in 400µl buffer (10mM HEPES, pH 7.4) to concentrations of 5µg/ml and 50µg/ml, and included a control with no fibrils (0µg/ml). PQS or pyoverdine were added to final concentrations of 200µM and 100µM respectively and the mixtures were incubated overnight (22°C, 400rpm). Each treatment was replicated 6 times. The mixtures were centrifuged (5 minutes, 1000g), the supernatant discarded and the retentate resuspended in 1ml of buffer and vortexed. This step was repeated 3 times and following the final centrifugation, retentate was resuspended in 400µl of buffer.

To determine whether fibrils prevented the loss of PQS and pyoverdine by dilution, we quantified the iron chelating activity of the mixes after repeated

washing. We added 100µl of each mixture to 100µl of CAS solution in a 96-well plate, incubated the plate in darkness for 30 minutes and measured absorbance at 630nm.

We also determined whether iron-chelators retained by fibrils were available for use by cells and could stimulate the growth of a pyoverdine non-producer strain, PAO1ΔpvdD, in iron-limited media. Overnight cultures of the pyoverdine non-producer were washed in M9 media, standardized to an OD of 0.2 and diluted 100-fold into 180µl of iron-limited CAA media in a 96-well plate. We then made up the volume in each well to 200µl with the different concentrations of fibrils (0, 5, or 50µg/ml) exposed to buffer, PQS or pyoverdine. Plates were incubated statically at 37°C and we measured the optical density of cultures (A_{600}) at 0hrs and 24hrs respectively.

Relative fitness of pyoverdine non-producers

To determine whether producing Fap fibrils influenced the ability of public goods producers to resist invasion by social cheats, we competed fibril producers and fibril non-producers (both pyoverdine producers) against the pyoverdine non-producer PAO1ΔPvdD under iron-limited conditions. After washing overnight cultures in M9 media, standardizing optical density (A_{600}) to 0.2 and diluting 100-fold, we mixed each combination of producer and non-producer at a ratio of 10:1 (producers: non-producers). 6ml of iron-limited CAA media was inoculated with 60µl of each mixture and incubated cultures statically at 37°C for 24 hours. Each treatment was replicated 20 times. We measured the frequency of pyoverdine producers and non-producers before and after 24hrs of incubation by plating onto KB agar (12g agar per litre of KB medium) and counting the number of colonies of producers and non-producers. We calculated the relative fitness of non-producers using the formula $w = (p_2(1-p_1))/(p_1(1-p_2))$ where p_1 and p_2 are the proportion of pyoverdine non-producers in the population before and after competition occurs respectively. A value of $w > 1$ indicates that the non-producer has a higher fitness than the pyoverdine producer and $w < 1$ indicates that the producer has a higher fitness than the non-producer.

Statistical Analyses

We carried out all statistical analyses in the R statistical environment (v3.3.3, <http://www.R-project.org>). Except where stated, we carried out standard analyses (T-

211 Test, Linear models, etc.) assuming normal errors. All analyses using Linear Mixed-
212 Effect Models (LMEM) included day as a random effect, to account for the fact that
213 replicates of the experiments were carried out on different days.

214 **Results**

215 **Population Growth and Pyoverdine Production.**

216 To determine whether the deletion of the Fap operon has consequences for growth or
217 alters investment in pyoverdine, we measured growth and pyoverdine production of
218 fibril producers and non-producers. After 24hrs of growth in iron-limited media, we
219 found no significant difference in cell density ($t=0.519$, $df=10$, $p=0.615$), pyoverdine
220 production ($t=-0.135$, $df=10$, $p=0.895$) and pyoverdine availability per cell ($t=-0.899$,
221 $df=10$, $p=0.389$) (Fig 1). These results suggest that not producing fibrils does not
222 significantly alter fibril non-producers investment in pyoverdine production nor
223 detrimentally affect its growth in an iron-limited environment.





Figure 1. Growth and pyoverdine production of Fap fibril producers (black lines, grey envelope) and non-producers (grey lines, pink envelope) over 24hrs in iron-limited media. Mean and standard error envelope are plotted for each. (a) Growth of strains, measured by cell density at A_{600} (b) Cumulative pyoverdine available, measured by $RFU_{400-460}$ (c) Pyoverdine available per cell, measured by $RFU_{400-460}/A_{600}$ (d) Rate of pyoverdine production, measured per cell per 30 minutes $((RFU_2 - RFU_1)/30 \text{ min})/A_{600(2)}$. Cell density, available pyoverdine and pyoverdine available per cell were not significantly different between fibril producers and non-producers.

Retention of iron-chelating metabolites.

After cells had been washed to remove unattached metabolites, we found that the iron-chelating activity, standardized by cell density, of fibril producers was significantly greater than that of non-producer cells (LMEM, $t=2.256$, $df=41$,

236 p=0.0295) (Fig 2). This suggests that producing fibrils can promote retention of iron-
 237 chelating secreted metabolites under conditions where otherwise they would be
 238 removed.



239 **Figure 2.** Retention of iron-chelating metabolites by fibril producers and non-producers. Iron
 240 chelating activity was quantified using a CAS assay and standardized by cell density (Iron-
 241 chelating activity per cell ($(1-A_{630})/A_{600}$)). Boxplots give the median value, boxes show the
 242 interquartile range and whiskers extend to the largest or smallest value respectively. Iron-
 243 chelating activity of fibril producers was significantly greater than that of non-producers.

244 We also isolated fibrils and determined whether they could prevent the loss of the
 245 known iron-chelators, pyoverdine and PQS, by dilution. A strong fluorescence

emission, with a maximum at ~490 nm when excited at 450 nm from our Thioflavin T assays confirmed the presence of amyloid structures in our purified protein solution (Fig S1). We found that isolated fibrils exposed to pyoverdine or PQS showed increased iron-chelating activity relative to fibrils exposed to buffer, as revealed by a significant interaction effect between fibril concentration and pyoverdine ($F_{3,32}=12.984$, $p=0.00105$) or PQS ($F_{3,32}=82.95$, $p=2.12 \times 10^{-10}$) (Fig 3a). Increasing concentrations of fibrils, without the addition of pyoverdine or PQS, did not increase levels of iron-chelation ($F_{1,16}=0.008$, $p=0.931$). This suggests that the increase in iron-chelating activity is due to the retention of pyoverdine and PQS by fibrils and not the fibrils themselves.

We also determined whether pyoverdine or PQS molecules retained by fibrils could be utilized by pyoverdine non-producers in an iron-limited environment. We found that fibrils exposed to pyoverdine increased the growth of a pyoverdine non-producer strain, PAO1 Δ pvdD, relative to fibrils exposed to buffer, demonstrated by a significant interaction effect between fibril concentration and pyoverdine ($F_{3,32}=50.43$, $p=4.68 \times 10^{-8}$) (Fig 3b). Fibrils alone did not increase the growth of cells ($F_{1,16}=1.281$, $p=0.274$), suggesting that the growth of the pyoverdine non-producer was promoted by the increasing availability of the siderophore pyoverdine in the iron-limited environment. We found that fibrils exposed to PQS actually repressed the growth of the pyoverdine non-producer relative to fibrils exposed to buffer ($F_{3,32}=4.983$, $p=0.032$) (Fig 3b). This effect is likely due to the fact that PQS is an iron-chelator but not a dedicated siderophore. Thus, it will bind iron but if cells lack a means of recovering the iron from PQS (by using, for example, a dedicated siderophore such as pyoverdine) it only serves to further iron-limit the surrounding environment.





Figure 3. Retention of iron-chelating metabolites by Fap fibrils. (a). Iron-chelating activity, measured as $(1-A_{630})$, of 0 $\mu\text{g/ml}$, 5 $\mu\text{g/ml}$ and 50 $\mu\text{g/ml}$ of untreated purified Fap fibrils (white) or purified Fap fibrils treated with PQS (cyan) or pyoverdine (green). Iron-chelating activity increases with increasing fibril concentration for both pyoverdine and PQS treatments. (b) Growth, measured as $A_{600(24\text{hrs})}-A_{600(0\text{hrs})}$, of a pyoverdine non-producer (PAO1 ΔpvdD) in iron-limited media supplemented with 0 $\mu\text{g/ml}$, 5 $\mu\text{g/ml}$ and 50 $\mu\text{g/ml}$ of untreated purified Fap fibrils or purified Fap fibrils treated with PQS (cyan) or pyoverdine (green). Growth of a pyoverdine non-producer increases with increasing concentrations of fibrils treated with pyoverdine and decreases with increasing concentrations of fibrils treated with PQS.

Competition Assays

282 To determine whether producing fibrils can influence the outcome of cooperator-
 283 cheat competitive dynamics, we competed fibril producers and non-producers with a
 284 pyoverdine non-producing strain (PAO1ΔPvdD). We found that in mixed cultures,
 285 pyoverdine non-producers increase in frequency when in competition with both fibril
 286 producers ($w=5.67$, $t=7.64$, $df=5$, $p=3.041 \times 10^{-4}$) and non-producers ($w=8.76$, $t=8.39$,
 287 $df=5$, $p=1.959 \times 10^{-4}$) (Fig 4). However, pyoverdine non-producer relative fitness is
 288 significantly greater in competition with Fap fibril non-producers than with fibril
 289 producers ($t=2.789$, $df=10$, $p=0.0192$). These results suggest that producing fibrils can
 290 limit the invasion of a pyoverdine non-producer into a population of pyoverdine
 291 producers.



292 **Figure 4.** Relative fitness (RF) of a pyoverdine non-producer (PAO1ΔPvdD) in competition
 293 with fibril producers and non-producers in iron-limited media. Boxplots give the median
 294 value, boxes show the interquartile range and whiskers extend to either 1.5 times the

interquartile range or to the largest or smallest value respectively. Outliers are shown as small circles. Pyoverdine non-producer significantly increase in frequency over 24hrs in competition with both fibril producers and non-producers, but pyoverdine non-producer relative fitness is significantly higher when competed against fibril non-producers than against fibril producers.

Discussion

Bacterial cells rely upon the secretion of extracellular public goods to acquire resources from the environment. However, cells fail to benefit from these goods if they are lost to the environment through diffusion or are utilised by social cheats. We present evidence that Fap fibril production in *P. aeruginosa* limits the loss of secreted public goods involved in iron acquisition, and that this can impact upon competition between public goods producers and cheats over iron. Fibril-producing cells show improved retention of iron-chelating metabolites (Fig. 2) and isolated Fap fibrils bind both PQS and pyoverdine (Fig. 3), supporting a role for Fap fibrils in retaining these costly public goods. Secreted public goods such as pyoverdine are often exploited by social cheats: subsequently, cells employ mechanisms that constrain the ability of cheats to benefit from products they produce. We show that fibril production limits the extent to which public goods cheats invade a population of cooperators (Fig 4), suggesting that retention of secreted goods by Fap fibrils also has consequences for cooperator-cheat competitive dynamics.

After washing to strip away unbound metabolites from the environment surrounding cells, fibril producers showed improved retention of iron-chelating metabolites (Fig 2), and purified fibrils appear to bind both pyoverdine and PQS, preventing their loss by dilution (Fig 3a). This supports a role for fibrils in retaining costly iron chelating metabolites. Transient binding of PQS by fibrils occurs due to the hydrophobic nature of both: potentially explaining the strong iron chelating activity of fibrils exposed to PQS despite repeated washing (Fig 3a)(Seviour *et al.* 2015). Pyoverdine, as a larger amphiphilic molecule, is not expected to bind strongly through hydrophobic interactions (Kummerli *et al.* 2014). However, it may possess hydrophobic domains that allow a more limited binding to fibrils (Visca *et al.* 2007). Why is binding, and subsequently limited diffusion, of secreted metabolites likely of

importance to cells? The majority of cells will experience conditions in their natural environments that remove costly secreted metabolites by diffusion or convection, limiting their utility to producing cells (Liu & Tay. 2002, Drescher *et al.* 2014,). Limiting diffusion of secreted metabolites may allow cells to invest in less of a costly, recyclable public good if it is not constantly being lost to the environment (Kummerli & Brown. 2010). It may also increase the range of conditions under which producing extracellular metabolites is a viable strategy: for example, limiting diffusion of quorum sensing molecules may expand the range of conditions under which QS controlled behaviours are inducible (Emge *et al.* 2016, Kim *et al.* 2016).

How does the presence or absence of Fap fibrils influence competition over iron? Our results show that pyoverdine non-producers do significantly better in competition with fibril non-producers than with fibril producers (Fig 4). One possible reason for this is that failure to produce fibrils alters investment in pyoverdine production. This is important because non-producers gain a competitive advantage only when a cooperator is paying the cost of producing a public good (Ghoul *et al.* 2016). Thus, if cells that do not produce fibrils increase per cell pyoverdine production, or increase the duration of pyoverdine production, this may increase the competitive advantage to non-producers. However, we see no significant difference in per cell production of pyoverdine between fibril producers and non-producers (Fig 1), suggesting that altered investment in pyoverdine does not explain the success of pyoverdine non-producers in the population.

As fibrils appear to prevent the loss of pyoverdine by dilution (Fig 3a), it may be that during competition with social cheats, fibril producers can limit or slow the diffusion of pyoverdine away from producing cells. This would have the effect of partially privatizing pyoverdine: ensuring the benefits of pyoverdine production accrue to producing cells and proximate cooperators, and limiting the competitive advantage of non-producers (Dobay *et al.* 2014, Scholz & Greenberg 2016). Another, non-mutually exclusive explanation, is that fibrils can help trap iron close to cells by binding PQS. Our results show that PQS is bound by fibrils but fails to stimulate the growth of a pyoverdine non-producing mutant in an iron-limited environment (Fig 3a, 3b), PQS, while a strong iron chelator, is not a dedicated siderophore as it cannot transfer iron across cell membranes. Thus it has been suggested that PQS may act as an iron trap, binding iron close to cells and making it easier for dedicated siderophores such as pyoverdine to shuttle iron into cells (Diggle *et al.* 2007b).

Previous studies have demonstrated that diffusion of secreted public goods can be limited, through maintaining cell-cell contacts or simply by environmental viscosity (Kummerli *et al.* 2009, Julou *et al.* 2013). Our results suggest that binding of secreted metabolites by Fap fibrils may also limit the diffusion of secreted metabolites and consequently play a role in shaping cooperator-cheat dynamics.

Is the production of functional amyloids an adaptation to reduce or prevent exploitation by social cheats? We have demonstrated the ability of Fap fibrils to partially-privatize secreted metabolites but functional amyloids also play important roles in surface adhesion and biofilm formation and structure in *Pseudomonads* and many other bacterial species (Barnhart & Chapman 2006, Zeng *et al.* 2015, Taglialegna *et al.* 2016). This suggests that functional amyloids may primarily play a role in surface attachment and biofilm formation and maintenance but have been co-opted by *Pseudomonas aeruginosa* as a means of retaining costly secreted metabolites. Exaptation is not uncommon in bacteria, with many traits co-opted to perform additional, or even novel, functions. For example, PQS is a signaling molecule that also plays an important role in iron acquisition, while bacterial conjugation systems have been co-opted for roles in effector translocation in Type IV secretion systems (Diggle *et al.* 2007b, Guglielmini *et al.* 2013).

Cooperative public goods production is common in bacterial populations despite the potential loss of costly public goods through diffusion or advection, or to social cheats (Hibbing *et al.* 2010). We have demonstrated that Fap fibrils can prevent loss of iron-chelating metabolites by dilution and limit the invasion of social cheats during competition over iron. Interestingly, other secreted metabolites that serve as public goods, such as pyochelin, pyocyanin and the QS molecules C4-HSL and C12-HSL are also small, hydrophobic molecules and likely predisposed to binding to Fap fibrils. We have discussed only the benefits of retaining secreted metabolites. However, there may also be costs associated with retention. For example, retaining pyocyanin could increase local oxidative stress. Future work should seek to reveal if Fap fibrils can limit the loss of these secreted metabolites, and any associated costs and benefits of doing so. There is also evidence for intraspecific and interspecific variation in Fap fibril production in *Pseudomonads* (Dueholm *et al.* 2013). It would also be interesting to see if levels of fibril production correlate with an isolate's ability to retain secreted metabolites in turbulent environments. “By

393 helping limit the diffusion of public goods, Fap fibrils may play an important role in
394 explaining the stability of secreted public goods in bacterial populations.

395 **References**

396 Andersen, S. B., Marvig, R. L., Molin, S., Krogh Johansen, H., & Griffin, A. S. 2015.
397 Long-term social dynamics drive loss of function in pathogenic bacteria. *Proceedings*
398 *of the National Academy of Sciences*. 112(34): 10756–10761.

399 Barnhart, M.M. and Chapman, M.R., 2006. Curli biogenesis and function. *Annual*.
400 *Reviews Microbiology*, 60, pp.131-147.

401 Bruce, J.B., Cooper, G.A., Chabas, H., West, S.A. and Griffin, A.S., 2017. Cheating
402 and resistance to cheating in natural populations of the bacterium *Pseudomonas*
403 *fluorescens*. *Evolution*, 71(10), pp.2484-2495.

404 Butaitė, E., Baumgartner, M., Wyder, S. and Kümmerli, R., 2017. Siderophore
405 cheating and cheating resistance shape competition for iron in soil and freshwater
406 *Pseudomonas* communities. *Nature Communications*, 8(1), p.414.

407 Cordero, O. X., & Ventouras, L. A. 2012. Public good dynamics drive evolution of
408 iron acquisition strategies in natural bacterioplankton populations. *Proceedings of the*
409 *National Academy of Sciences*. 109(49): 20059-64

410 Diggle, S.P., Griffin, A.S., Campbell, G.S. & West, S.A. 2007a. Cooperation and
411 conflict in quorum-sensing bacterial populations. *Nature*. 450(7168): 411-4.

412 Diggle, S.P., Matthijs, S., Wright, V.J., Fletcher, M.P., Chhabra, S.R., Lamont, I.L.,
 413 Kong, X., Hider, R.C., Cornelis, P., Cámara, M. and Williams, P., 2007b. The
 414 *Pseudomonas aeruginosa* 4-quinolone signal molecules HHQ and PQS play
 415 multifunctional roles in quorum sensing and iron entrapment. *Chemistry &*
 416 *biology*, 14(1), pp.87-96.

417 Dobay, A., Bagheri, H.C., Messina, A., Kümmerli, R. and Rankin, D.J., 2014.
 418 Interaction effects of cell diffusion, cell density and public goods properties on the
 419 evolution of cooperation in digital microbes. *Journal of Evolutionary Biology*, 27(9),
 420 pp.1869-1877.

421 Drescher, K., Nadell, C.D., Stone, H.A., Wingreen, N.S. and Bassler, B.L., 2014.
 422 Solutions to the public goods dilemma in bacterial biofilms. *Current Biology*, 24(1),
 423 pp.50-55.

424 Dueholm, M.S., Søndergaard, M.T., Nilsson, M., Christiansen, G., Stensballe, A.,
 425 Overgaard, M.T., Givskov, M., Tolker-Nielsen, T., Otzen, D.E. and Nielsen, P.H.,
 426 2013. Expression of Fap amyloids in *Pseudomonas aeruginosa*, *P. fluorescens*, and *P.*
 427 *putida* results in aggregation and increased biofilm formation. *Microbiology*
 428 *Open*, 2(3), pp.365-382.

429 Dumas, Z., Ross-Gillespie, A. and Kümmerli, R., 2013. Switching between
 430 apparently redundant iron-uptake mechanisms benefits bacteria in changeable

431 environments. *Proceedings of the Royal Society B: Biological Sciences*, 280(1764),
 432 p.20131055.

433 Emge, P., Moeller, J., Jang, H., Rusconi, R., Yawata, Y., Stocker, R. and Vogel, V.,
 434 2016. Resilience of bacterial quorum sensing against fluid flow. *Scientific reports*, 6,
 435 p.33115.

436 Flemming, H.C., Wingender, J., Szewzyk, U., Steinberg, P., Rice, S.A. and
 437 Kjelleberg, S., 2016. Biofilms: an emergent form of bacterial life. *Nature Reviews*
 438 *Microbiology*, 14(9), pp.563-575.

439 Ghequire, M. G. K., & De Mot, R. (2014). Ribosomally encoded antibacterial
 440 proteins and peptides from *Pseudomonas*. *FEMS Microbiology Reviews*. 38(4): 523–
 441 568.

442 Ghoul, M., Griffin, A.S. & West, S.A. 2014(a). Toward an evolutionary definition of
 443 cheating. *Evolution*. 68(2): 318-31

444 Ghoul, M., West, S.A., McCorkell, F.A., Lee, Z.B., Bruce, J.B. and Griffin, A.S.,
 445 2016. Pyoverdine cheats fail to invade bacterial populations in stationary
 446 phase. *Journal of Evolutionary Biology*, 29(9), pp.1728-1736.

447 Ghysels, B., Thi Min Dieu, B., Beatson, S., Pirnay, J.-P., Ochsner, U., Vasil, M. &
 448 Cornelis, P. 2004. FpvB, an alternative type I ferripyoverdine receptor of
 449 *Pseudomonas aeruginosa*. *Microbiology* 150: 1671–1680.

450 Griffin, A.S., West, S.A. & Buckling, A. 2004. Cooperation and competition in
 451 pathogenic bacteria. *Nature*. 430(7003): 1024-7

452 Guglielmini, J., De La Cruz, F. and Rocha, E.P., 2012. Evolution of conjugation and
 453 type IV secretion systems. *Mol. Biol. Evo*, 30(2), pp.315-331.

454 Hibbing, M.E., Fuqua, C., Parsek, M.R. and Peterson, S.B., 2010. Bacterial
 455 competition: surviving and thriving in the microbial jungle. *Nature Reviews*
 456 *Microbiology*, 8(1), pp.15-25.

457 Julou, T., Mora, T., Guillon, L., Croquette, V., Schalk, I.J., Bensimon, D. and
 458 Desprat, N., 2013. Cell–cell contacts confine public goods diffusion inside
 459 *Pseudomonas aeruginosa* clonal microcolonies. *Proceedings of the National Academy*
 460 *of Sciences*, 110(31), pp.12577-12582.

461 Kim, M.K., Ingremeau, F., Zhao, A., Bassler, B.L. and Stone, H.A., 2016. Local and
 462 global consequences of flow on bacterial quorum sensing. *Nature microbiology*, 1,
 463 p.15005.

464 Kümmerli, R., Griffin, A.S., West, S.A., Buckling, A. and Harrison, F., 2009. Viscous
 465 medium promotes cooperation in the pathogenic bacterium *Pseudomonas*
 466 *aeruginosa*. *Proceedings of the Royal Society of London B: Biological*
 467 *Sciences*, 276(1672), pp.3531-3538.

468 Kümmerli, R. and Brown, S.P., 2010. Molecular and regulatory properties of a public
 469 good shape the evolution of cooperation. *Proceedings of the National Academy of*
 470 *Sciences*, 107(44), pp.18921-18926.
 471
 472 Kümmerli, R., Schiessl, K.T., Waldvogel, T., McNeill, K. and Ackermann, M., 2014.
 473 Habitat structure and the evolution of diffusible siderophores in bacteria. *Ecology*
 474 *Letters*, 17(12), pp.1536-1544.

 475 Liu, Y. and Tay, J.H., 2002. The essential role of hydrodynamic shear force in the
 476 formation of biofilm and granular sludge. *Water research*, 36(7), pp.1653-1665.

 477 McNally, L., Bernardy, E., Thomas, J., Kalzigi, A., Pentz, J., Brown, S.P., Hammer,
 478 B.K., Yunker, P.J. and Ratcliff, W.C., 2017. Killing by Type VI secretion drives
 479 genetic phase separation and correlates with increased cooperation. *Nature*
 480 *Communications*, 8, p.14371.

 481 Mund, A., Diggle, S.P. and Harrison, F., 2017. The fitness of *Pseudomonas*
 482 *aeruginosa* quorum sensing signal cheats is influenced by the diffusivity of the
 483 environment. *mBio*, 8(3), pp.e00353-17.

 484 Nadell, C.D., Bucci, V., Drescher, K., Levin, S.A., Bassler, B.L. and Xavier, J.B.,
 485 2013, March. Cutting through the complexity of cell collectives. In *Proc. R. Soc.*
 486 *B* (Vol. 280, No. 1755, p. 20122770).

487 Popat, R., Cornforth, D.M., McNally, L. and Brown, S.P., 2015. Collective sensing
 488 and collective responses in quorum-sensing bacteria. *Journal of the Royal Society*
 489 *Interface*, 12(103), p.20140882.

490 Ross-Gillespie, A., Gardner, A., West, S.A. and Griffin, A.S., 2007. Frequency
 491 dependence and cooperation: theory and a test with bacteria. *The American*
 492 *Naturalist*, 170(3), pp.331-342.

493 Sandoz, K.M., Mitzimberg, S.M. & Schuster, M. 2007. Social cheating in
 494 *Pseudomonas aeruginosa* quorum sensing. *Proceedings of the National Academy of*
 495 *Sciences*. 104(40): 15876-81

496 Seviour, T., Hansen, S.H., Yang, L., Yau, Y.H., Wang, V.B., Stenvang, M.R.,
 497 Christiansen, G., Marsili, E., Givskov, M., Chen, Y. and Otzen, D.E., 2015.
 498 Functional amyloids keep quorum-sensing molecules in check. *Journal of Biological*
 499 *Chemistry*, 290(10), pp.6457-6469.

500 Scholz, R.L. and Greenberg, E.P., 2015. Sociality in *Escherichia coli*: enterochelin is
 501 a private good at low cell density and can be shared at high cell density. *Journal of*
 502 *Bacteriology*, 197(13), pp.2122-2128.

503 Schwyn, B. and Neilands, J.B., 1987. Universal chemical assay for the detection and
 504 determination of siderophores. *Analytical Biochemistry*, 160(1), pp.47-56.

- 505 Taglialegna, A., Lasa, I. and Valle, J., 2016. Amyloid structures as biofilm matrix
506 scaffolds. *Journal of Bacteriology*, 198(19), pp.2579-2588.
- 507 Visca, P., Imperi, F. and Lamont, I.L., 2007. Pyoverdine siderophores: from
508 biogenesis to biosignificance. *Trends microbiol*, 15(1), pp.22-30.
- 509 West, S.A. and Buckling, A., 2003. Cooperation, virulence and siderophore
510 production in bacterial parasites. *Proceedings of the Royal Society of London B:*
511 *Biological Sciences*, 270(1510), pp.37-44.
- 512 West, S.A., Winzer, K., Gardner, A. and Diggle, S.P., 2012. Quorum sensing and the
513 confusion about diffusion. *Trends in microbiology*, 20(12), pp.586-594.
- 514 Zeng, G., Vad, B.S., Dueholm, M.S., Christiansen, G., Nilsson, M., Tolker-Nielsen,
515 T., Nielsen, P.H., Meyer, R.L. and Otzen, D.E., 2015. Functional bacterial amyloid
516 increases *Pseudomonas* biofilm hydrophobicity and stiffness. *Frontiers in*
517 *Microbiology*, 6.