

Synthesis of the Ca^{2+} -mobilizing messengers NAADP and cADPR by intracellular CD38 enzyme in the mouse heart: Role in β -adrenoceptor signaling

Received for publication, April 3, 2017, and in revised form, May 13, 2017. Published, Papers in Press, May 24, 2017. DOI 10.1074/jbc.M117.789347

Wee K. Lin[‡], Emma L. Bolton[‡], Wilian A. Cortopassi^{§¶}, Yanwen Wang^{||}, Fiona O'Brien[‡], Matylda Maciejewska[‡], Matthew P. Jacobson[¶], Clive Garnham[‡], Margarida Ruas[‡], John Parrington[‡], Ming Lei^{‡1}, Rebecca Sitsapesan^{‡2}, Antony Galione^{‡3}, and Derek A. Terrar^{‡4}

From the [‡]Department of Pharmacology, University of Oxford, Mansfield Road, Oxford OX1 3QT, United Kingdom, the

[§]Department of Chemistry, Chemistry Research Laboratory, University of Oxford, Mansfield Road, Oxford OX1 3TA, United

Kingdom, the [¶]Department of Pharmaceutical Chemistry, University of California, San Francisco, California 94158, and the ^{||}Faculty of Biology, Medicine, and Health, University of Manchester, Manchester M13 9NT, United Kingdom

Edited by Roger J. Colbran

Nicotinic acid adenine dinucleotide phosphate (NAADP) and cyclic ADP-ribose (cADPR) are Ca^{2+} -mobilizing messengers important for modulating cardiac excitation–contraction coupling and pathophysiology. CD38, which belongs to the ADP-ribosyl cyclase family, catalyzes synthesis of both NAADP and cADPR *in vitro*. However, it remains unclear whether this is the main enzyme for their production under physiological conditions. Here we show that membrane fractions from WT but not *CD38*^{−/−} mouse hearts supported NAADP and cADPR synthesis. Membrane permeabilization of cardiac myocytes with saponin and/or Triton X-100 increased NAADP synthesis, indicating that intracellular CD38 contributes to NAADP production. The permeabilization also permitted immunostaining of CD38, with a striated pattern in WT myocytes, whereas *CD38*^{−/−} myocytes and nonpermeabilized WT myocytes showed little or no staining, without striation. A component of β -adrenoreceptor signaling in the heart involves NAADP and lysosomes. Accordingly, in the presence of isoproterenol, Ca^{2+} transients and contraction amplitudes were smaller in *CD38*^{−/−} myocytes than in the WT. In addition, suppressing lysosomal function with bafilomycin A1 reduced the isoproterenol-induced increase in Ca^{2+} transients in cardiac myocytes from WT but not *CD38*^{−/−} mice. Whole hearts isolated from *CD38*^{−/−} mice and exposed to isoproterenol showed reduced arrhythmias. SAN4825, an ADP-ribosyl cyclase inhibitor that reduces cADPR and NAADP synthesis in mouse membrane fractions, was shown to bind to CD38 in docking simulations and reduced the isoproterenol-induced arrhythmias in WT hearts. These observations support generation of NAADP and cADPR by intracellular CD38, which contributes to effects of β -adrenoreceptor stimulation to

increase both Ca^{2+} transients and the tendency to disturb heart rhythm.

ADP-ribosyl cyclases (ARCs)⁵ are enzymes capable of converting NAD into cyclic adenosine diphosphate ribose (cADPR). In addition, some of these cyclases (for example, CD38 and *Aplysia* cyclase), can (at least under *in vitro* conditions) also produce nicotinic acid adenine dinucleotide phosphate (NAADP) using NADP as substrate in the presence of nicotinic acid (NA) (see reviews in Ref. 1). cADPR and NAADP are both potent Ca^{2+} -releasing messengers (2, 3) that play significant roles in various signaling pathways in plants and animals, including humans (see reviews in Refs. 4, 5). Cellular functions mediated by these two messengers include hormonal secretion, smooth muscle contraction, immune responses, fertilization, and neuromodulation (4, 5). In the heart, both cADPR and NAADP modulate excitation–contraction coupling (6–9). These effects seem particularly important after stimulation of β -adrenoreceptors, presumably associated with the elevated levels of NAADP and cADPR (7, 10, 11). It has been postulated that the increased concentrations of these messengers result from enhanced production by endogenous enzymes and that excessive production of these molecules contributes to the development of cardiac hypertrophy and arrhythmias associated with stimulation of β -adrenoreceptors (9, 12, 13). The enzyme(s) responsible for the production of cADPR and NAADP is/are therefore a potential therapeutic target, and drugs, including SAN4825, to target these enzyme(s) are under development (14). However, the nature and location of ARCs and/or NAADP-synthesizing enzymes in the heart remain unclear.

It was mentioned above that CD38 can function as an ARC catalyzing the synthesis of both cADPR and NAADP. However,

The authors declare that they have no conflicts of interest with the contents of this article.

Author's Choice—Final version free via Creative Commons CC-BY license.

This article contains supplemental Figs. S1–S7.

¹ Funded by the British Heart Foundation (BHF), Medical Research Council, and BHF Centre of Research Excellence (CRE).

² Funded by the BHF and BHF CRE.

³ Funded by a Wellcome Trust Senior Investigatorship and BHF CRE.

⁴ To whom correspondence should be addressed. Tel.: 44-1865271943; Fax: 44-1865271853; E-mail: derek.terrar@pharm.ox.ac.uk.

⁵ The abbreviations used are: ARC, ADP-ribosyl cyclase; cADPR, cyclic ADP-ribose; NAADP, nicotinic acid adenine dinucleotide phosphate; SR, sarcoplasmic reticulum; SUEH, sea urchin egg homogenate; NGD, nicotinamide guanine dinucleotide; cGDP, cyclic GDP-ribose; NA, nicotinic acid; HSR, heavy sarcoplasmic reticulum fraction; LHR, light sarcoplasmic reticulum fraction; CaT, Ca^{2+} transient; RyR, ryanodine receptor.

a recent study on cardiac tissue suggests that CD38 mediates the synthesis of cADPR but not NAADP (15). This paper proposes that another cardiac enzyme on lysosomes synthesizes NAADP in response to intracellular Ca^{2+} fluxes in the myocyte following stimulation of β -adrenoceptors. The enzyme was referred to as NAADP-synthesizing enzyme. A further proposal was that NAADP produced within the lysosome causes Ca^{2+} release from this organelle and that this Ca^{2+} promotes cADPR synthesis by CD38 in neighboring endosomes. cADPR transported out of the endosome was then proposed to regulate ryanodine receptor opening in the sarcoplasmic reticulum (SR). Suppression of CD38 expression in $\text{CD38}^{-/-}$ mice was shown to reduce the cardiac hypertrophy that is associated with chronic exposure to the β -adrenoceptor agonist isoproterenol (15). The proposal that CD38 promotes cADPR synthesis within the organelle is consistent with the commonly accepted view that CD38 can show activity as an ectoenzyme at the plasma membrane with an active site facing away from the cytosol (16). Work on lymphocytes first established the external orientation of CD38 (17). Because targets for NAADP and cADPR in cardiac muscle appear to be intracellular, the physiological relevance in the heart of enzyme activity of an ecto-CD38 remains unclear. Some have proposed expression of intraorganelle ARC (with an active site facing the lumen) in other tissues for cADPR and NAADP synthesis (see reviews in Refs. 18, 19). Evidence from sea urchin egg supports the synthesis of cADPR within organelles, followed by transport of the active substance into the cytosol to enable its participation in Ca^{2+} signaling (20). However, it has recently been suggested that CD38 can also exist in the opposite orientation, with the active site of the enzyme facing the cytosol, and that the active site can still function to promote synthesis of cADPR in this topological orientation (21–23).

The aim of this study was to further characterize the nature and identity of cADPR- and NAADP-synthesizing enzymes in cardiac muscle, exploring their possible location on intracellular membranes and investigating the hypothesis that CD38 might function as the main enzyme responsible for the synthesis for both Ca^{2+} -mobilizing messengers. The contributions of this pathway to physiological excitation–contraction coupling and pathological development of disturbances of cardiac rhythm were also investigated because this pathway provides a novel target for the development of antiarrhythmic drugs.

Results

CD38 is the main enzyme responsible for cardiac synthesis of NAADP and cADPR

A membrane-enriched preparation containing sarcolemma and SR was derived from mouse heart muscle both from WT and $\text{CD38}^{-/-}$ mice. A sensitive bioassay for measuring NAADP was provided by sea urchin egg homogenates (SUEHs) (24), which show a homologous desensitization of the Ca^{2+} -releasing effect of NAADP at remarkably low concentrations (from $\sim 10^{-10}$ to 10^{-8} M). Example traces and a calibration curve are shown in Fig. 1, A and B. The membrane-enriched preparation

from WT hearts produced NAADP when supplied with NADP and NA (Fig. 1, C and D). Interestingly, there was no comparable synthesis of NAADP with mixed membranes from $\text{CD38}^{-/-}$ hearts ($n = 3$; $p \leq 0.001$; Fig. 1, C and D). A widely accepted conventional nicotinamide guanine dinucleotide (NGD) assay was used to determine the rate of cADPR synthesis (25). Using this assay, we detected the cyclization of NGD (an analogue of NAD) into cyclic GDP-ribose (cGDPR, a fluorescent analogue of cADPR, by membranes from WT hearts but not from $\text{CD38}^{-/-}$ hearts ($n = 6$; $p \leq 0.001$; Fig. 1, E and F). These observations are consistent with the hypothesis that CD38 is the main enzyme responsible for the cardiac synthesis of both NAADP and cADPR.

Intracellular location of CD38 and possible association with the SR

Fig. 2A shows that, when the enzymatic activity of intact cardiac ventricular myocytes from WT mice was investigated using the same assay procedures as for the membrane-enriched preparation from heart muscle above, there was very little synthesis of NAADP when cells were supplied with NADP and NA in the extracellular solution. However, permeabilization of the ventricular cell membranes with saponin (0.01% w/v) caused a substantial increase in the rate of production of NAADP, consistent with access of substrates to the cell interior of healthy cells, permitting production of NAADP by enzymes with intracellular active sites (Fig. 2A). As saponin preferentially targets cholesterol-rich membranes (26) such as the sarcolemma, rather than the SR, we used Triton X-100 (0.1% v/v) to further expose potential intracellular enzymes (perhaps including those with lumen-facing active sites in various organelles). However, Triton X-100 permeabilization in the presence of saponin failed to cause a substantial increase in the rate of NAADP production (Fig. 2A). There appeared to be little or no difference between the effects of the two membrane-permeabilizing agents applied alone or in combination (Fig. 2B). The above experiments were repeated on intact myocytes from $\text{CD38}^{-/-}$ hearts, and there was little or no synthesis of NAADP both in intact myocytes and after membrane permeabilization with Triton X-100 (Fig. 2B). Control experiments showed that neither Triton X-100 nor saponin at the concentration used in these experiments affected the ability of the sea urchin egg homogenate bioassay to measure NAADP (supplemental Fig. S1).

A rabbit polyclonal antibody to human CD38 was used to further investigate the presence of this enzyme in mouse ventricular myocytes because the amino acid sequence (residues 1–170) used to generate the antibody showed extensive similarities between the two species (supplemental Fig. S2). When the plasma membrane was intact in the absence of Triton X-100, there was little immunolabeling in WT mouse cardiac myocytes, and this labeling appeared to be concentrated in membrane patches (Fig. 2C, top panels). When Triton X-100 was used to permeabilize the cell membrane, the labeling by the CD38 antibody was much more extensive and showed striations with a spacing corresponding to the sarcomere length of slightly less than $2 \mu\text{m}$ (Fig. 2C, center panels). Convincing evidence that the human CD38 antibody was indeed labeling the





Figure 1. *In vitro* ADP-ribosyl cyclase activities and NAADP production were not detected in mixed-membrane preparations from $CD38^{-/-}$ mouse heart. We evaluated the ability of a membrane preparation containing plasmalemma and SR from WT but not $CD38^{-/-}$ mouse heart to catalyze the synthesis of NAADP and cADPR. *A*, example traces of the NAADP measurement assay system based on SUEH, in which increasing concentrations of NAADP show a desensitization response to the activating substance NAADP. *B*, observations reflecting enhanced desensitization at increasing concentrations of NAADP were used to construct the calibration curve. *C*, using this assay, we show that a membrane preparation from WT mice catalyzed the synthesis of NAADP, whereas a similar preparation from $CD38^{-/-}$ mouse heart lacked this ability. The time course of NAADP production is shown. *D*, the average rate of NAADP synthesis from these data ($n = 4$ for both groups). *E* and *F*, the membrane preparation from WT mice catalyzed synthesis of cGDP (an analogue of cADPR) with NGD as a substrate (the time course is shown in *E*, and a bar graph showing the average rate of cGDP synthesis is shown in *F*), whereas membranes from $CD38^{-/-}$ mice lacked this ability ($n = 6$ for both groups). *r.f.u.* indicates relative fluorescence unit. Note that WT membranes showed clear synthesis of both messengers, whereas there was no detectable synthesis with $CD38^{-/-}$ membranes. Data are expressed as mean \pm S.E. ***, $p \leq 0.001$; n = number of preparations.

mouse CD38 was provided by the observation that little or no labeling was detected in permeabilized cardiac ventricular myocytes from $CD38^{-/-}$ mice (Fig. 2C, bottom panels). This striated pattern with a spacing close to $2 \mu\text{m}$ in myocytes from WT but not $CD38^{-/-}$ mice is also supported by Fig. 2D, which shows plots of intensity for a line positioned along the long axis of the cell. Labeling by the antibody to human CD38 was also investigated in ventricular myocytes isolated from rabbit heart. In the presence of Triton X-100 to permeabilize cell membranes, immunolabeling with a striated pattern was again observed (Fig. 2E) and appeared to be strikingly similar to that observed in WT mouse myocytes. Again, the separation

between striations appeared to correspond to sarcomere length, and labeling of ryanodine receptors (RyR2) showed a similar spacing. This immunolabeling with a striated pattern was not observed when Triton X-100 or the primary antibody was omitted (Fig. 2F). It therefore appears that membrane permeabilization was necessary for maximal NAADP synthesis and for labeling of intracellular CD38 in ventricular myocytes from WT mice and that neither NAADP synthesis nor intracellular labeling of CD38 were observed in ventricular myocytes from $CD38^{-/-}$ mice.

In addition, CD38 labeling on permeabilized rabbit atrial myocytes was also found to have a striated pattern, similar to



The observation that CD38 appears to be associated with the SR prompted us to investigate enzyme activity in a sheep SR vesicle preparation that is commonly used to investigate heart SR proteins, including RyR2 (27–34). cGDPH synthesis with NGD as substrate was observed in this sheep cardiac SR preparation, consistent with the presence of ARC activity (supplemental Fig. S4A). As expected, the effects were reduced in the

The effects of isoproterenol to increase the amplitudes of Ca^{2+} transients (CaTs) and contractions accompanying electrical stimulation were smaller in cardiac myocytes from $\text{CD38}^{-/-}$ mice than in those from WT myocytes

The functional effects of genetic knockout of CD38 were investigated in cardiac ventricular myocytes. Previous work has shown that levels of NAADP and cADPR are elevated following β -adrenoceptor stimulation with isoproterenol (7, 11). It was also found that both NAADP and cADPR contribute, by different mechanisms, to the isoproterenol-induced increase in the amplitude of CaTs and contractions evoked by electrical stimulation in cardiac myocytes (7–9, 12). Consistent with these observations and the hypothesis that CD38 provides the major synthetic pathway for NAADP and cADPR, it was found that the amplitudes of CaTs and contractions in the presence of isoproterenol were smaller in cardiac myocytes from $\text{CD38}^{-/-}$ mice than in myocytes from WT mice (Fig 3, A–D).

Bafilomycin A1, to inhibit lysosomal NAADP function, reduced the increase in amplitude of CaTs caused by isoproterenol in electrically stimulated myocytes from WT but not $\text{CD38}^{-/-}$ mice

In the case of NAADP actions, previous work has demonstrated that a component of the isoproterenol-induced increase in the amplitude of CaTs induced by electrical stimulation appears to depend on NAADP acting via TPC2 channels in lysosomal membranes (9). Lysosomes have also been shown to form associations with the SR, consistent with the formation of Ca^{2+} microdomains between the organellar membranes (36). In addition, it has been shown that, when lysosomal NAADP function is disrupted with bafilomycin A1, the increase in CaT amplitude caused by isoproterenol is reduced (7, 9). The reduction in the isoproterenol-induced increase in the amplitude of CaT following bafilomycin A1 treatment (10 μM) is shown again here for WT myocytes but, in contrast bafilomycin A1, failed to reduce the isoproterenol-induced increase in CaT amplitude in myocytes from $\text{CD38}^{-/-}$ mice (Fig. 3E). This was also evident from the ratio of effects of isoproterenol with and without bafilomycin A1 (Fig. 3F, showing a lack of difference in $\text{CD38}^{-/-}$ myocytes compared with a reduced effect in the WT). Sample records are shown in Fig. 3G. These observations are therefore consistent with the hypothesis that cardiac myocytes from $\text{CD38}^{-/-}$ mice are unable to synthesize significant amounts of NAADP and that, therefore, this pathway can no longer be functionally suppressed by bafilomycin A1. The observations provide further support for a role for NAADP synthesized by CD38 in this physiological process in WT cardiac muscle. The remaining isoproterenol-induced increases in myocytes from $\text{CD38}^{-/-}$ mice are likely to have resulted, at least in part, from the expected PKA-mediated effects on L-type Ca^{2+} channels and phospholamban that are thought not to involve CD38 (37). The CD38-dependent pathway underlying these observations is considered at greater length under “Discussion.”

In vitro and in silico evidence showed that a novel inhibitor of cardiac ARC, SAN4825, also inhibited NAADP synthesis at physiological pH, possibly through targeting CD38

SAN4825 is a novel drug developed as an inhibitor of cardiac-specific ARCs and has been shown (14) to suppress the ability of a rat SR membrane preparation to synthesize a fluorescent analogue of cADPR ($\text{IC}_{50} \approx 1.3 \mu\text{M}$). This drug was also shown to partially inhibit the ARC activity of human purified CD38 at higher concentration (14). We tested the ability of SAN4825 to inhibit both NAADP and cADPR synthesis by the mouse membrane preparation for which the evidence reported above shows CD38 to be the major enzyme catalyzing synthesis of these Ca^{2+} -mobilizing messengers. SAN4825 was observed to cause a concentration-dependent inhibition of both NAADP synthesis (Fig. 4A) and cADPR synthesis (Fig. 4B). A similar inhibition of NAADP synthesis by SAN4825 was also observed in saponin-permeabilized myocytes from mouse hearts (supplemental Fig. S5). SAN4825 inhibition of NAADP synthesis was observed to be greater at pH 7.2, and this will be considered at greater length under “Discussion.”

The active site residues Glu-150, Asp-159, Glu-230, Trp-129, and Trp-193 in mouse CD38 correspond to Glu-146, Glu-226, Asp-155, Trp-125, and Trp-189 in human CD38. These residues in human CD38 are believed to play an important role in NAADP and cADPR synthesis following mutagenesis studies (38). In a docking simulation using Glide, we demonstrate that the most favorable docking conformation of SAN4825 at pH 7.2 in mouse CD38 (PDB code 2EG9 (39)) is in close proximity to these active-site residues (Fig. 4C). Further superimposition of the binding conformation of SAN4825 in mouse CD38 with the binding conformations of an NAADP analogue and ADPRP in human CD38 (PDB code 4F46 (40)) suggests that SAN4825 occupies the same binding site as these substrates, in close proximity to Glu-230 of mouse CD38 (Fig. 4C). The pyridine ring of SAN4825 also showed π – π stacking with Trp-193 of mouse CD38. These π – π interactions were also observed in the crystal structure of wild-type human CD38 in complex with the NAADP analogue and ADPRP (40).

Further computational studies also demonstrated that the single active site is the only plausible binding site of SAN4825. Specifically, blind docking results using SwissDock showed that approximately 90% of the predicted docking sites (250 predictions in total) of SAN4825 in mouse CD38 are within 4.5 Å from the center of the active site (Fig. 4, D and E). SiteMap also presented a similar result, showing that the active site is the only site to present a SiteScore value closer to 1.0 (0.97), indicative of a difficult but druggable site with an estimated volume of 286 Å³. The other binding sites presented SiteScore values close to 0.8, suggestive of undruggable cavities with volumes smaller than 120 Å³. Taken together, these observations suggest that SAN4825 acts as a competitive inhibitor of NAADP and cADPR synthesis.

$\text{CD38}^{-/-}$ hearts were more resistant to arrhythmias during overstimulation of the β -adrenoceptor pathway

It is known that excessive β -adrenoceptor stimulation can lead to cardiac arrhythmias (41). The tendency to arrhythmias




Figure 3. Contribution of NAADP signaling pathway to the β -adrenoceptor-mediated positive inotropy was lost in the $CD38^{-/-}$ ventricular myocytes. Ca^{2+} transients were triggered by cardiac action potentials in ventricular myocytes from WT and $CD38^{-/-}$ mice. **A**, the amplitudes of Ca^{2+} transients in WT ($n = 9$) and $CD38^{-/-}$ myocytes ($n = 15$) are shown in the presence of isoproterenol. **B**, example images. Under these conditions, amplitudes of CaTs accompanying action potentials were smaller in cardiac myocytes from $CD38^{-/-}$ mice than from WT mice. ISO, isoproterenol. **C**, sarcomere shortening in the presence of isoproterenol accompanying action potentials was also smaller in myocytes from $CD38^{-/-}$ mice than from WT mice (WT, $n = 12$; $CD38^{-/-}$, $n = 11$). **D**, example traces. The observations support a functional role for the CD38 enzyme during β -adrenergic receptor stimulation. **E**, the effects of isoproterenol on the amplitude of CaTs with and without bafilomycin A1 ($10 \mu M$) in ventricular myocytes from WT and $CD38^{-/-}$ mice ($n = 16$ – 25 in each group). **F**, these observations are replotted to show the magnitude of the isoproterenol-induced changes with and without bafilomycin A1 in myocytes from WT and $CD38^{-/-}$ mice. **G**, representative traces. Bafilomycin A1 reduced the effect of isoproterenol on the amplitude of CaTs in myocytes from WT but not $CD38^{-/-}$ mice. The isoproterenol-induced increase in amplitudes of CaTs was therefore less in the presence of bafilomycin A1 in WT myocytes, whereas, in myocytes from $CD38^{-/-}$ mice, the isoproterenol-induced increases in CaT amplitude were similar with and without bafilomycin A1. Data are expressed as mean \pm S.E. *, $p \leq 0.05$; **, $p \leq 0.01$; n = number of cells.

in intact mouse hearts was assessed by four different protocols (burst pacing with increasing frequency, 50-Hz burst pacing with increasing current, S1S2 pacing, and dynamic pacing; see “Experimental Procedures” for details and scoring methods). WT hearts showed a significant increase in arrhythmogenicity after application of 300 nM isoproterenol

(Fig. 5A). In contrast, no significant increase in arrhythmogenicity was observed following isoproterenol application in hearts from $CD38^{-/-}$ mice (Fig. 5A). Two other methods of data presentation were used to highlight changes in arrhythmogenicity. Fig. 5B shows the increase in arrhythmic events after isoproterenol and demonstrates the significant





Figure 4. *In vitro* and *in silico* evidence for the action of SAN4825 to inhibit NAADP and cADPR production through targeting CD38. **A** and **B**, SAN4825 inhibited NAADP (**A**) and cADPR (**B**) synthesis in mouse mixed-membrane preparations at both pH 7.2 and pH 4.5. Note that the synthesis of NAADP in the absence of SAN4825 was higher at pH 4.5 than at pH 7.2. **C**, superimposition of the binding conformation of the NAADP analogue in human CD38 (*left, red*, PDB code 4F46, chain B) on the most favorable binding conformation of SAN4825 (*left, blue*) in mouse CD38 (PDB code 2EG9) at pH 7.2 following Glide docking. A superimposition of the binding conformation of ADPRP in human CD38 (*right, red*, PDB code 4F46, chain A) on the most favorable binding conformation of SAN4825 (*right, blue*) in mouse CD38 (PDB code 2EG9) is also shown (*right*). **D**, the predicted binding sites (*orange*) for SAN4825 in mouse CD38 following blind docking in the SwissDock server. Inner cavities are shown in *yellow*. **E**, the distances between the predicted docking sites and the center of the active site were plotted. Data are expressed as mean \pm S.E. *, $p \leq 0.05$; **, $p \leq 0.01$; ***, $p \leq 0.001$; n.s., not significant; $n \geq 3$ preparations in each group.

reduction in arrhythmogenicity in hearts from $CD38^{-/-}$ mice compared with WT mice. When the observations were plotted as pie charts representing the proportion of hearts showing arrhythmias, it was clear that the proportion of hearts from $CD38^{-/-}$ mice that were susceptible to isoproterenol-induced arrhythmogenicity (one of six hearts) was less than the corresponding proportion of WT hearts (seven of eight hearts, Fig 5C).

The cumulative percentage of hearts that became arrhythmic in the presence of isoproterenol (300 nM) during the burst pacing protocol with increasing pacing frequency and increasing pacing current is shown in Fig. 5D, again showing that hearts

from $CD38^{-/-}$ mice had a reduced tendency to arrhythmias under these conditions. Hearts from WT mice also had a significantly lower current threshold for initiation of arrhythmias, as shown in Fig. 5E. The percentage of $CD38^{-/-}$ hearts that exhibited signs of arrhythmias (in the absence and presence of 300 nM isoprenaline) during dynamic pacing and S1S2 pacing protocols was also lower than those of WT hearts (Fig. 5F). Representative traces are shown in Fig. 5G. Taken together, these observations show that the tendency for arrhythmias during exposure to high concentrations of β -adrenoceptor agonist was greatly reduced in hearts from $CD38^{-/-}$ mice.



β -adrenoreceptor stimulation were broadly similar to the consequences of genetic knockdown of CD38 described above (Fig. 5, A–G), although the drug effects appeared to be less than CD38 knockout.

Discussion

The main findings reported here are that CD38 is the principal enzyme in heart muscle responsible for synthesis of two Ca^{2+} -mobilizing messengers, NAADP and cADPR, and that a major component of CD38 appears to be intracellular, perhaps associated with the SR (although there may be some additional CD38 enzyme on the cell surface). These findings are consistent with the conclusions of Gul *et al.* (15) that CD38 mediates synthesis of cADPR in cardiac myocytes but contrast with Gul *et al.* (15) in their proposal that a different enzyme (NAADP-synthesizing enzyme), is responsible for NAADP production in these cells.

When comparing our observations with those of Gul *et al.* (15), it is important to note differences in the experimental conditions, particularly with respect to the actions of isoproterenol. In our experiments, the effects of β -adrenoceptor stimulation were studied under a physiological condition where cardiac myocytes were paced and the amplitude of CaTs triggered by electrical stimulation were investigated at low concentrations of isoproterenol (2–5 nM). In contrast, the increases in cytosolic Ca^{2+} reported by Gul *et al.* (15) were in cardiac myocytes that were not electrically paced and for which the isoproterenol concentration was 1000 times greater (2 μM). These very high concentrations of isoproterenol are likely to have caused spontaneous Ca^{2+} waves and/or oscillations, and the recorded maintained increases in cytosolic Ca^{2+} may have been the result of summation of signals (with an image collection rate of one per 3 s) from many cardiac myocytes in the field of view.

Gul *et al.* (15) have suggested that NAADP production precedes synthesis of cADPR and propose that Ca^{2+} released from lysosomes by NAADP reaches nearby CD38 in endosomes, causing an increase in the enzyme activity of CD38 to produce cADPR. This interpretation relies on observations in their Fig. 2, although it could be argued that the quite small observed differences in apparent time to peak synthesis of NAADP or cADPR (that appeared to be 15 s, or one measurement time point) may result from the ability to detect these messengers rather than reflect differences in the underlying physiology. Previous work in whole hearts showed that production of both cADPR and NAADP occurs rapidly following isoproterenol addition (11). It should also be mentioned that protein structural work shows that binding of Ca^{2+} causes conformational changes in CD38 that are expected to be inhibitory rather than stimulatory for enzyme action, whereas functional studies show little or no change in enzyme activity at the Ca^{2+} concentrations likely to be experienced by intracellular CD38 (42).

The possibility that the major active enzyme CD38 is located at or close to SR membranes is supported by our immunohistochemical observations showing that CD38 labeling has a striated pattern with a spacing that closely resembles that of the SR protein RyR2 both in ventricular and atrial myocytes. The lack of similar immunohistochemical staining in myocytes from CD38^{-/-} mice demonstrates that these observations are unlikely to result from lack of specificity of the antibody used to identify CD38. Because atrial myocytes lack the extensive t-tubule network seen in ventricular myocytes (and atrial t-tubules do not show the regular sarcomere spacing observed in the

ventricle), the striated pattern also seen in atrial myocytes must result from staining of CD38 that appears to be associated with the SR (which does show a striated pattern in both atrial and ventricular muscle). It has recently been shown that lysosomes also show some association with the SR (as well as mitochondria), but the striated pattern for lysosomal markers is less distinct than the observations reported here for CD38 (36).

Although evidence for NAADP and cADPR function in cardiac myocytes has so far been demonstrated only in mice, rats, and guinea pigs (7–10, 43, 44) it is our hypothesis that this pathway operates in all mammalian hearts and that CD38 associated with the SR is the main synthetic enzyme. In the past, SR membrane preparations from the hearts of various animals have been used as a model to study the synthesis of cADPR or cGDPR by ARC (10, 14, 45, 46). In particular, sheep SR preparations have been widely used for the study of RyR2 and other SR proteins (27–34). We are the first to show that, in addition to catalysis of the cyclization reactions, sheep SR vesicles can also catalyze the base exchange reaction that produces NAADP (supplemental Fig. S4). Although the functional effects of this pathway have yet to be investigated in sheep, this species has the advantage of allowing easy preparation of amounts of biological material that are orders of magnitude greater than what can be prepared from much smaller mouse hearts. The relative purity of similar preparations is supported by electron microscopy and biochemical studies (45, 47, 48). Most importantly, to date, CD38 is the only mammalian enzyme that has been identified to have the ability to synthesize both cADPR and NAADP as a consequence of reactions at a multifunctional active site (see reviews in Ref. 49). Although we have yet to fully characterize the ARC on sheep SR, our initial investigations show that the enzymatic properties observed in sheep SR preparations have a high resemblance to those of human and mouse CD38. Therefore, our observations on sheep SR vesicles corroborate our hypothesis that CD38 is expressed intracellularly and appears to show a preferential association with the SR in cardiac tissue.

The functional observations with the Sanofi drug SAN4825, which reduced synthesis of both NAADP and cADPR, are consistent with its inhibitory actions at a single enzyme. The most favorable Glide docking conformations of SAN4825 with mouse CD38 at pH 7.2 suggest that this compound occupies the same binding site as the NAADP analogue and ADPRP, in close proximity with Glu-226. The pyridine ring of SAN4825 also shows π – π interaction with Trp-189 of mouse CD38. The blind docking results from the SwissDock server also predicted a single inhibitory binding site for SAN4825, and this site corresponds to the enzymatic active site of CD38. These SAN4825–CD38 binding conformations showed some resemblance to the native binding conformation between CD38 and its substrates, such as NAD, nicotinamide mononucleotide (NMN), cADPR, and NAADP (40, 50, 51).

Interestingly, the inhibition of NAADP but not cADPR synthesis by SAN4825 was observed to be greater at the cytosolic pH of 7.2 than at the acidic pH of 4.5 (Fig. 4, A and B). Therefore, we further investigated the binding conformations of SAN4825 at the active site of mouse CD38 with considerations of different protonation states of SAN4825 and mouse CD38 protein amino acids predicted by Epik and Propka 3.1, respec-

tively (supplemental Fig. S6). The SAN4825 molecule can be protonated or unprotonated, and CD38 will adopt different protein conformations at different pH values. Nevertheless, with all four combinations (two SAN4825 protonation states and two CD38 conformations), all predicted binding sites of SAN4825 were still in proximity of the catalytic residues of mouse CD38 (supplemental Fig. S6). When considering the weakening of inhibition of NAADP synthesis by SAN4825 at pH 4.5, the following factors need to be taken into account: NAADP synthesis by human CD38 (3) and mouse CD38 (Fig. 4A) was observed to be enhanced at pH 4.5; inhibition of cADPR synthesis by SAN4825 remained unchanged at pH 4.5 (Fig. 4B); SAN4825 was predicted to bind to the CD38 active site at both pH 7.2 and pH 4.5 (supplemental Fig. S6); no NAADP synthesis was observed in membrane fractions from $CD38^{-/-}$ mice at pH 4.5 (supplemental Fig. S7). In view of these observations and predictions, it appears that the observed weakened inhibition of SAN4825 at pH 4.5 does not result from a degradation of the drug or from a major change in the binding conformation of the drug and cannot be accounted for by a non-CD38 enzyme synthesizing NAADP. Although a weakened inhibition of NAADP synthesis by SAN4825 was observed at pH 4.5, evidence from our docking simulations and *in vitro* experiments still supports the ability of SAN4825 to inhibit the synthesis of NAADP and cADPR as a competitive inhibitor.

What are the implications of our findings for the normal physiology of the heart? Our observations show that the amplitudes of CaTs initiated by electrical stimulation at a constant rate were increased by β -adrenoceptor stimulation with isoproterenol at low concentrations (2–5 nM) and that part of this response was prevented in WT mice by the use of bafilomycin A1 to suppress the contribution of the lysosomal pathway involving NAADP and TPC2 (9). In contrast, in myocytes from $CD38^{-/-}$ mice, bafilomycin A1 no longer suppressed this component of the isoproterenol action, consistent with the importance of CD38 in the mechanism underlying NAADP production and the subsequent response. This is consistent with the need for coupling of CD38 to receptors, leading to NAADP synthesis and action via TPC2 proteins in the cardiac lysosomal signaling pathway (7, 44).

Previous experiments support elevation of NAADP and cADPR levels in response to β -adrenoceptor stimulation by isoproterenol (7, 10, 11, 52). Our observations show that genetic knockout and pharmacological inhibition of CD38 provide resistance for arrhythmias associated with acute exposure to excessive stimulation with β -adrenoceptors compared with WT hearts. Taken together, these observations are consistent with an effect of β -adrenoceptor stimulation by as yet unidentified mechanisms to enhance the catalytic activity of CD38, leading to increased levels of these Ca^{2+} -mobilizing messengers (7, 11), which, in turn, enhance CaTs evoked by cardiac action potentials or trigger arrhythmic events resulting from excessive Ca^{2+} -mobilizing messengers.

We have shown recently that the cardiac hypertrophy and associated tendency to arrhythmias that is provoked by chronic exposure to β -adrenoceptor stimulation is markedly reduced when the influence of the NAADP pathway is suppressed in mice lacking NAADP-regulated TPC2 proteins (9). Because

CD38 is upstream in this signaling pathway, it seems reasonable to propose that CD38 contributes to these processes and that hearts from $CD38^{-/-}$ mice might also be resistant to cardiac hypertrophy and associated arrhythmias following chronic exposure to β -adrenoceptor stimulation. The recent observations from Gul *et al.* (15) show that this is indeed the case and provide further support for this hypothesis.

Graeff *et al.* (53) have suggested that a protonation of NA at acidic pH is essential for the binding of NA to acidic residues (Glu-146 and Asp-155) on CD38 because, at neutral pH, the electrostatic repulsion between them would reduce substrate binding and, consequently, the rate of NAADP synthesis. However, our observations showed that cardiac muscle CD38 can catalyze the formation of NAADP at physiological pH, although at a lower rate than at acidic pH. It should also be noted that, as a result of the remarkably high potency of NAADP as a Ca^{2+} -releasing messenger, the threshold concentration for NAADP to cause lysosomal Ca^{2+} release is very low, probably in the tens of nanomolar range (7), and, therefore, even a small amount of synthesized NAADP at pH 7.2 would be enough to activate the NAADP signaling pathway. This hypothesis is further supported by the work of Lee and co-workers (23), demonstrating that the active site of CD38 can remain functional under the reducing conditions of the cytosol and at cytosolic pH.

Although CD38 was thought to exist as a type II transmembrane protein with its C-terminal catalytic domain on the outside of the cell, Zhao *et al.* (22) showed, in cell lines and primary human cells, the presence of a proportion of CD38 in the type III form, with the catalytic domain facing the cytosol. In our study, substrates to permit synthesis of NAADP and/or cADPR were not included in the perfusion solution for intact heart or single-cell studies, and even if there was extracellular synthesis of these messengers, it seems likely that any relevant transporter of molecules will be made redundant or inefficient as a consequence of the constant flow of perfusion solution under the conditions of our experiments. These factors are expected to prevent the cytosolic accumulation of NAADP and cADPR from an extracellular source, and we have therefore favored arguments that, in cardiac cells, CD38 in the type III form might mediate intracellular synthesis of these Ca^{2+} -mobilizing agents in WT type animals, leading to the observed effects on CaTs, contraction and arrhythmogenicity of the heart. This is consistent with our observations that permeabilized single cells (using saponin and/or Triton X-100) have higher enzymatic activity. Permeabilization was also required for immunolabeling of CD38. The finding that a functionally important component of CD38 appears to be intracellular and associated with the SR in the heart should not be considered to be in conflict with observations showing other locations for similar enzymes in different cell types (16, 19, 20, 54). It is possible that the Ca^{2+} signaling molecules cADPR and NAADP are so important in many diverse roles in the plant and animal kingdoms that different locations of synthetic enzymes may have evolved to suit different functional needs.

Because the known major modulatory influences of NAADP and cADPR on CaTs in cardiac myocytes have been shown to be mediated through SR-associated proteins (sarco-endoplasmic reticulum Ca^{2+} -ATPase and RyRs) (9, 43, 55), it seems reason-

able to propose that the most efficient operation of the signaling pathways involving NAADP and cADPR might be best served by a CD38 synthetic enzyme that is predominantly intracellular (perhaps located at or close to the SR membrane). An interesting hypothesis would be that CD38, NAADP, and cADPR as well as the relevant substrates could act within microdomains close to sites of Ca^{2+} uptake and/or release on the SR membrane, thereby facilitating the signaling pathways (see scheme in Ref. 44). Our recent observations concerning the location of lysosomes in close proximity to the SR are relevant in this context (36).

In the discussion above, the main focus is to highlight the pivotal role of intracellular CD38 in catalyzing the cyclization and base exchange reactions in hearts, particularly following activation of the β -adrenoceptor pathway. There are several limitations to this study. Although we have proposed that the CD38 enzyme appears to be associated with SR membranes, we have not addressed the topology of the enzyme, particularly concerning the orientation of the active site, and this will be the subject of future experiments. In addition, although this study adds to previous work showing a link between β -adrenoceptor stimulation and synthesis of NAADP and cADPR, further work is needed to establish the extent of activation of the CD38 pathway in the absence of this stimulation and to investigate the cellular mechanisms by which β -adrenoceptor stimulation up-regulates CD38 activity.

In conclusion, this study provides novel evidence for cardiac synthesis of two potent and important Ca^{2+} mobilizing molecules, NAADP and cADPR, by intracellular CD38, probably associated with or close to the sarcoplasmic reticulum. The cellular pathway starting with CD38 and involving NAADP and cADPR acts to enhance excitation–contraction coupling, particularly during activation of β -adrenoceptors. Our observations demonstrating that $\text{CD38}^{-/-}$ hearts are resistant to β -adrenoceptor-associated arrhythmias also reveal the potential of targeting CD38-signaling pathways to treat cardiac disease.

Experimental procedures

Single-cell studies

Cell isolation—The $\text{CD38}^{-/-}$ mouse line was developed by Cockayne *et al.* (56) and obtained from The Jackson Laboratory ($\text{Cd38}^{\text{tm1Lnd/J}}$). The protocol for generating and maintaining the mice was approved by the Oxford University Ethical Review Committee (Pharmacology Subcommittee) and Home Office. We confirm that all methods were performed in accordance with the relevant United Kingdom Home Office and institutional guidelines and regulations. Ventricular myocytes were isolated from male mice (C57BL6 or $\text{CD38}^{-/-}$, 16–24 weeks old) and male New Zealand White rabbits (~1 kg) using 0.3–0.5 mg ml⁻¹ collagenase (type II, Worthington Biochemical Corp.) The detailed methods were described in Refs. 9, 36. Rabbit atrial myocytes were isolated using a similar protocol to that used to isolate rabbit ventricular myocytes.

Ca^{2+} imaging in single cardiac myocytes—Mouse myocytes were incubated with Fluo-5F/AM (5 μM) for 20 min. CaTs were stimulated at 1 Hz by carbon fiber electrodes placed at the side

of the superfusion bath. Mouse ventricular myocytes were visualized using a Nikon Axiovert 200 inverted microscope with an attached Nipkow spinning disk confocal unit (CSU-10, Andor Technology). Excitation light was provided by a 488-nm diode laser (Cairn Research Ltd., Kent, UK) passed through the Nipkow unit and delivered to the sample through the objective. Emitted light passed back through the CSU-10 unit and was detected using an Andor iXON897 electron multiplying charge-coupled device (EMCCD) camera (Andor Technology) at 50 frames/s. Images were recorded and analyzed using Andor iQ software (Andor Technology). For analysis, background fluorescence was subtracted, and multiple transients were averaged to obtain the basal CaTs before isoproterenol application and the peak CaTs amplitude during 5 min of isoproterenol application. Data are presented as F/F_0 so that fluorescence data are presented relative to diastolic fluorescence.

Contraction studies—Cells were stimulated at 1 Hz by carbon fiber electrodes placed at the side of the superfusion bath, and the contractile properties were studied using the IonOptix system (IonOptix Corp.) to measure sarcomere length. Cells were visualized via a $\times 40$ oil objective using an IonOptix MyoCam (IonOptix Corp.) that sampled images at a frequency of 240 Hz. Sinusoidal optical density traces arising from the alternating light and dark bands of the contractile machinery were then transformed into a signal of sarcomere length by application of a fast Fourier transform by the IonWizard sarcomere length acquisition software (IonOptix Corp.). Length measurements were calibrated using a stage micrometer with 2- μm graduations so that the number of pixels per micrometer recorded by the MyoCam could be entered into the software as a fixed value. Amplitude of sarcomere shortening was calculated by deduction of systolic from diastolic sarcomere length. Analysis was performed using IonWizard 5 software (IonOptix Corp.). All values represent an average of 10 contractions. Cells for both Ca^{2+} imaging experiments and contraction studies were superfused with medium containing 130 mM NaCl, 5.4 mM KCl, 3.5 mM MgCl_2 , 1.8 mM CaCl_2 , 10 mM glucose, 5 mM HEPES, and 0.4 mM NaH_2PO_4 (pH 7.4) at 36 °C.

In vitro biochemical assay

Mixed-membrane and sarcoplasmic reticulum-enriched membrane preparations—As described previously, mixed-membrane vesicles were prepared from sheep (obtained from an abattoir) or male mouse (C57BL6 or $\text{CD38}^{-/-}$, 12–20 weeks old) cardiac tissue (57). A discontinuous sucrose gradient was used to separate the sarcolemmal membrane from the LSR and HSR membrane fractions. For each tissue, the protein concentration was calculated using a Bradford assay (58), and the various membrane fractions were aliquoted and stored at -80°C .

Assessment of the enzymatic cyclization rate of ADP-ribosyl cyclase—A final concentration of 500 μM NGD was added to the protein sample to initiate the enzymatic reaction. cGDPR formation was monitored through its fluorescence intensity at 410 ± 5 nm during excitation, with light of a wavelength of 300 ± 3 nm (25). The cyclase activity of each reaction mixture was then defined by the change in fluorescence with time, *i.e.* the slope of the line in the fluorescence *versus* the time plot obtained by linear regression.

Detection and quantification of NAADP production—The level of NAADP production was quantified by means of an NAADP inactivation bioassay using SUEHs (24). The SUEHs were prepared using a method reported previously (59). Free Ca^{2+} concentration was measured with Fluo-3 by monitoring fluorescence intensity at excitation and emission wavelengths of 490 and 535 nm, respectively. To synthesize NAADP, 500 μM NADP and 7 mM NA were preincubated with the SR-enriched cardiac preparation at 37 °C. Then the protein–substrate mixtures were diluted before addition to the sea urchin egg homogenate. After 5 min, a standard 500 nM purified NAADP was further added to the sea urchin egg homogenate to elicit submaximal NAADP-induced Ca^{2+} release. The Ca^{2+} release response to the standard 500 nM NAADP is dependent on the subthreshold concentration of the NAADP in the preceding protein–substrate mixture addition. The concentration of NAADP in the protein–substrate mixture was quantified through interpolation of a standard dose-response curve from known concentrations of purified NAADP.

For both the NGD and SUEH assay, 10–20 μl of the SR-enriched or mixed-membrane cardiac preparation was incubated in a solution containing 150 mM KCl, 0.5 mM MgCl_2 , and 10 mM HEPES (pH 7.2). An acetate-based buffer was used in the condition where an acidic pH of 4.5 is required. For experiments investigating the enzymatic activities of permeabilized myocytes, myocytes were incubated with a solution comprising 70 mM KCl, 5 mM MgCl_2 , 5 mM potassium glutamine, 20 mM taurine, 0.04 mM EGTA, 5 mM succinic acid, 20 mM KH_2PO_4 , 10 mM glucose, and 5 mM HEPES (pH 7.2) with KOH. This solution contained either 0.01% saponin (w/v) or 0.1% Triton X-100 (v/v). Fluorimetry was performed in a Novostar plate reader (BMG Labtech).

Immunocytochemistry studies

Cells were fixed in 4% paraformaldehyde/PBS for 15 min, washed in PBS (three changes, 5 min each), permeabilized using 0.1% Triton X-100 (Sigma-Aldrich) for 10 min, washed in PBS, and blocked with 1% bovine serum albumin for 60 min before being incubated with the primary antibody at 4 °C overnight. The next day, cells were first washed with PBS before being incubated with secondary antibody at room temperature for 2 h (either Alexa Fluor 488–conjugated donkey anti-rabbit or DyLight 550 donkey anti-mouse, 1:1000 dilution) and then washed. Finally, coverslips were mounted using Vectashield® and permanently sealed. Cells were stored in the dark at 4 °C and visualized within 2 days. For control experiments, we performed the same procedure with omission of the primary antibodies or with cells from $\text{CD38}^{-/-}$ mice. Primary antibodies against CD38 (sc-15362, Santa Cruz Biotechnology) and RyR2 (ab2827, Abcam) were used at 1:100 and 1:200 dilution, respectively. Observations were carried out using a Nikon A1 confocal laser-scanning microscope equipped with a $\times 60$ objective, and images were processed using ImageJ software.

Ex vivo programmed electrical stimulations

Hearts were isolated from male mice (16–24 weeks old) and mounted in a Langendorff perfusion system. Hearts were perfused with Krebs solution comprising 119 mM NaCl, 25 mM NaHCO_3 , 4 mM KCl, 1.8 mM sodium pyruvate, 1.2 mM KH_2PO_4 ,

1 mM MgCl_2 , 1.8 mM CaCl_2 , and 10 mM glucose (pH 7.4) at 37 °C. Hearts were left to stabilize for 30 min. During this 30-min time window, hearts were also preincubated with drugs or vehicle present inside the Krebs solutions. To assess propensity to ventricular arrhythmias, hearts were subjected to four different programmed electrical stimulations in the absence and presence of 300 nM isoproterenol.

Frequency-varying burst pacing protocol—For the frequency-varying burst pacing protocol, a set of three trains of stimuli was delivered to the heart. Each train of stimuli consisted of 50 stimuli with intervals (cycle lengths) of 100 ms and was separated by a pacing-free interval of 2 s. The cycle length was progressively reduced from 100 to 20 ms in 10-ms decrements after each set.

Amplitude-varying burst pacing protocol—For the amplitude-varying burst pacing protocol, a set of three trains of stimuli was delivered to the heart. Each train of stimuli consisted of 50 stimuli with intervals of 20 ms and was separated by a pacing-free interval of 2 s. The current amplitude was progressively increased by factors of the pacing threshold current, *i.e.* the minimum current to evoke an action potential in a 1:1 manner. The protocol was terminated when a ventricular arrhythmia was observed or a current of 35 mA was reached.

Dynamic S1 pacing—Cycles of a decremental paced electrogram fractionation sequence comprising a 100-stimulus (S1) drive train were applied at a pacing interval of 135 ms initially. The pacing interval was reduced by 5 ms between successive drive trains until a pacing interval of 10 ms was reached.

S1S2 pacing—Cycles of a decremental paced electrogram fractionation sequence comprising an 8-Hz, eight-beat stimulus (S1) drive train, followed by an extra stimulus (S2), were applied. Initially, the interval between S1 and S2 was set to be 125 ms and was progressively reduced by 1 ms between successive drive trains until the preparation became refractory.

Ventricular arrhythmia was defined as six or more consecutive regular premature waveforms or irregular fibrillating waveforms. The arrhythmogenicity of the heart was determined by the number of protocols that successfully triggered arrhythmias. A scoring system was devised on this basis. For example, the score for arrhythmogenicity was 3 when a heart exhibited signs of arrhythmias under three of four protocols. The maximum score was 4 when all four protocols successfully triggered arrhythmias.

In silico study

A conformational search in implicit solvent (water) with the OPLS_2005 force field (60) was performed for SAN4825 using MacroModel Schrödinger release 2017-1. An adapted blind docking of all possible conformations in the mouse CD38 (PDB code 2EG9, chain A) was performed with Glide (61, 62), considering all inside cavities, as determined by SiteMap (63). Protein was prepared using the protein preparation tool available in Maestro, filling in missing side chains and loops with Prime (64, 65) and minimizing the resulting structure with the OPLS_2005 force field. The protonation states of SAN4825 were predicted using the Epik software (66). Protein residue protonation states were determined by PropKa 3.1 (67, 68).

Following docking, binding energies were calculated for ranking the best docking conformations using the molecular mechanics/generalized born surface area scoring function. For the scoring, residues close to 5 Å were considered flexible, allowing optimization of the docking conformations. To confirm that SAN4825 binds only to the active site containing the most important residues for NAADP and cADPR synthesis, SAN4825 was further submitted to a blind docking using the SwissDock server (69, 70). The distance between the predicted binding sites and the active site of CD38 was measured. The center of the active site of CD38 was determined to be the center of geometry of amino acid residues Trp-129, Glu-150, Asp-159, Trp-193, and Glu-230 of mouse CD38. These amino acid residues correspond to Trp-125, Glu-146, Asp-155, Trp-189, and Glu-226 in human CD38, which are important in CD38-mediated enzymatic syntheses, as identified in a site-directed mutagenesis study (38).

Drugs and reagents

The SAN4825 compound was a kind gift from Sanofi (Monsieur, France). β -NGD, NADP, NA, nicotinamide, Triton X-100, saponin, and bafilomycin A1 were from Sigma (St. Louis, MO).

Statistical analysis

All statistical comparisons were made by two-tailed Student's *t* tests unless stated otherwise. Mann-Whitney test and Fisher's exact test were used to compare arrhythmogenicity scores and categorical data, respectively. As only one in six hearts from CD38^{-/-} mice showed arrhythmias, one-sample *t* test ($H_0, \mu = 7$) was used for statistical analysis in Fig. 5E. All data are expressed as mean \pm S.E.

Author contributions—W. K. L. carried out all experiments, either solely or in collaboration with W. A. C. and M. P. J. (docking experiment), E. L. B. (contraction studies), Y. W. (MAP recording), F. O. B. (isolating SR fractions), and M. M. (sheep cardiac SR experiments). C. G. provided technical support for the plate reader experiments. M. R. was responsible for the breeding the CD38^{-/-} mouse line, and J. P. was the Home Office project license holder. W. K. L., A. G., M. L., and D. A. T. were responsible for the experimental design. All experiments were carried out in the laboratories of D. A. T., A. G., R. S., and M. L. All authors contributed to the writing and editing of the manuscript.

Acknowledgments—We thank Sanofi for providing the SAN4825 compounds. We also thank Dr. Daniel Aston for providing rabbit ventricular and atrial myocytes and Dr. Grant Churchill for constructive comments.

References

1. Ferrero, E., Lo Buono, N., Horenstein, A. L., Funaro, A., and Malavasi, F. (2014) The ADP-ribosyl cyclases: the current evolutionary state of the ARCs. *Front. Biosci.* **19**, 986–1002
2. Howard, M., Grimaldi, J. C., Bazan, J. F., Lund, F. E., Santos-Argumedo, L., Parkhouse, R. M., Walseth, T. F., and Lee, H. C. (1993) Formation and hydrolysis of cyclic ADP-ribose catalyzed by lymphocyte antigen CD38. *Science* **262**, 1056–1059

3. Aarhus, R., Graeff, R. M., Dickey, D. M., Walseth, T. F., and Lee, H. C. (1995) ADP-ribosyl cyclase and CD38 catalyze the synthesis of a calcium-mobilizing metabolite from NADP. *J. Biol. Chem.* **270**, 30327–30333
4. Lee, H. C. (2001) Physiological functions of cyclic ADP-ribose and NAADP as calcium messengers. *Annu. Rev. Pharmacol. Toxicol.* **41**, 317–345
5. Galione, A. (2015) A primer of NAADP-mediated Ca^{2+} signalling: from sea urchin eggs to mammalian cells. *Cell Calcium* **58**, 27–47
6. Rakovic, S., Galione, A., Ashamu, G. A., Potter, B. V., and Terrar, D. A. (1996) A specific cyclic ADP-ribose antagonist inhibits cardiac excitation-contraction coupling. *Curr. Biol.* **6**, 989–996
7. Macgregor, A., Yamasaki, M., Rakovic, S., Sanders, L., Parkesh, R., Churchill, G. C., Galione, A., and Terrar, D. A. (2007) NAADP controls cross-talk between distinct Ca^{2+} stores in the heart. *J. Biol. Chem.* **282**, 15302–15311
8. Collins, T. P., Bayliss, R., Churchill, G. C., Galione, A., and Terrar, D. A. (2011) NAADP influences excitation-contraction coupling by releasing calcium from lysosomes in atrial myocytes. *Cell Calcium* **50**, 449–458
9. Capel, R. A., Bolton, E. L., Lin, W. K., Aston, D., Wang, Y., Liu, W., Wang, X., Burton, R. A., Bloor-Young, D., Shade, K. T., Ruas, M., Parrington, J., Churchill, G. C., Lei, M., Galione, A., and Terrar, D. A. (2015) Two pore channels (TPC2s) and nicotinic acid adenine dinucleotide phosphate (NAADP) at lysosomal-sarcoplasmic reticular junctions contribute to acute and chronic β -adrenoceptor signaling in the heart. *J. Biol. Chem.* **290**, 30087–30098
10. Rakovic, S., and Terrar, D. A. (2002) In *Cyclic ADP-Ribose and NAADP* (Lee, H. C., ed), pp. 319–341. Springer, Boston
11. Lewis, A. M., Aley, P. K., Roomi, A., Thomas, J. M., Masgrau, R., Garnham, C., Shipman, K., Paramore, C., Bloor-Young, D., Sanders, L. E., Terrar, D. A., Galione, A., and Churchill, G. C. (2012) β -Adrenergic receptor signaling increases NAADP and +cADPR levels in the heart. *Biochem. Biophys. Res. Commun.* **427**, 326–329
12. Rakovic, S., Cui, Y., Iino, S., Galione, A., Ashamu, G. A., Potter, B. V., and Terrar, D. A. (1999) An antagonist of cADP-ribose inhibits arrhythmogenic oscillations of intracellular Ca^{2+} in heart cells. *J. Biol. Chem.* **274**, 17820–17827
13. Nebel, M., Schworer, A. P., Warszta, D., Siebrands, C. C., Limbrock, A. C., Swarbrick, J. M., Fliegert, R., Weber, K., Bruhn, S., Hohenegger, M., Geisler, A., Herich, L., Schlegel, S., Carrier, L., Eschenhagen, T., et al. (2013) Nicotinic acid adenine dinucleotide phosphate (NAADP)-mediated calcium signaling and arrhythmias in the heart evoked by β -adrenergic stimulation. *J. Biol. Chem.* **288**, 16017–16030
14. Kannt, A., Sicka, K., Kroll, K., Kadereit, D., and Gögelein, H. (2012) Selective inhibitors of cardiac ADPR cyclase as novel anti-arrhythmic compounds. *Naunyn-Schmiedeberg's Arch. Pharmacol.* **385**, 717–727
15. Gul, R., Park, D. R., Shawl, A. I., Im, S. Y., Nam, T. S., Lee, S. H., Ko, J. K., Jang, K. Y., Kim, D., and Kim, U. H. (2016) Nicotinic acid adenine dinucleotide phosphate (NAADP) and cyclic ADP-ribose (cADPR) mediate Ca^{2+} signaling in cardiac hypertrophy induced by β -adrenergic stimulation. *PLoS ONE* **11**, e0149125
16. De Flora, A., Franco, L., Guida, L., Bruzzone, S., and Zocchi, E. (1998) Ectocellular CD38-catalyzed synthesis and intracellular Ca^{2+} -mobilizing activity of cyclic ADP-ribose. *Cell Biochem. Biophys.* **28**, 45–62
17. Jackson, D. G., and Bell, J. I. (1990) Isolation of a cDNA encoding the human CD38 (T10) molecule, a cell surface glycoprotein with an unusual discontinuous pattern of expression during lymphocyte differentiation. *J. Immunol.* **144**, 2811–2815
18. De Flora, A., Zocchi, E., Guida, L., Franco, L., and Bruzzone, S. (2004) Autocrine and paracrine calcium signaling by the CD38/NAD⁺/cyclic ADP-ribose system. *Ann. N.Y. Acad. Sci.* **1028**, 176–191
19. Bruzzone, S., Guida, L., Sturla, L., Usai, C., Zocchi, E., and De Flora, A. (2012) Subcellular and intercellular traffic of NAD⁺, NAD⁺ precursors and NAD⁺-derived signal metabolites and second messengers: old and new topological paradoxes. *Messenger* **1**, 34–52
20. Davis, L. C., Morgan, A. J., Ruas, M., Wong, J. L., Graeff, R. M., Poustka, A. J., Lee, H. C., Wessel, G. M., Parrington, J., and Galione, A. (2008) Ca^{2+} signaling occurs via second messenger release from intraorganelle synthesis sites. *Curr. Biol.* **18**, 1612–1618

21. Zhao, Y. J., Zhang, H. M., Lam, C. M., Hao, Q., and Lee, H. C. (2011) Cytosolic CD38 protein forms intact disulfides and is active in elevating intracellular cyclic ADP-ribose. *J. Biol. Chem.* **286**, 22170–22177
22. Zhao, Y. J., Lam, C. M., and Lee, H. C. (2012) The membrane-bound enzyme CD38 exists in two opposing orientations. *Sci. Signal.* **5**, ra67
23. Zhao, Y. J., Zhu, W. J., Wang, X. W., Zhang, L. H., and Lee, H. C. (2015) Determinants of the membrane orientation of a calcium signaling enzyme CD38. *Biochim. Biophys. Acta* **1853**, 2095–2103
24. Genazzani, A. A., Empson, R. M., and Galione, A. (1996) Unique inactivation properties of NAADP-sensitive Ca^{2+} release. *J. Biol. Chem.* **271**, 11599–11602
25. Graeff, R. M., Walseth, T. F., Fryxell, K., Branton, W. D., and Lee, H. C. (1994) Enzymatic synthesis and characterizations of cyclic GDP-ribose: a procedure for distinguishing enzymes with ADP-ribosyl cyclase activity. *J. Biol. Chem.* **269**, 30260–30267
26. Jamur, M. C., and Oliver, C. (2010) Permeabilization of cell membranes. *Methods Mol. Biol.* **588**, 63–66
27. Mahony, L. (1988) Maturation of calcium transport in cardiac sarcoplasmic reticulum. *Pediatr. Res.* **24**, 639–643
28. Michalak, M. (1988) Identification of the Ca^{2+} -release activity and ryanodine receptor in sarcoplasmic-reticulum membranes during cardiac myogenesis. *Biochem. J.* **253**, 631–636
29. Sitsapasan, R., and Williams, A. J. (1990) Mechanisms of caffeine activation of single calcium-release channels of sheep cardiac sarcoplasmic reticulum. *J. Physiol.* **423**, 425–439
30. Sitsapasan, R., McGarry, S. J., and Williams, A. J. (1994) Cyclic ADP-ribose competes with ATP for the adenine nucleotide binding site on the cardiac ryanodine receptor Ca^{2+} release channel. *Circ. Res.* **75**, 596–600
31. Zissimopoulos, S., and Lai, F. A. (2005) Interaction of FKBP12.6 with the cardiac ryanodine receptor C-terminal domain. *J. Biol. Chem.* **280**, 5475–5485
32. Laver, D. R. (2007) Ca^{2+} stores regulate ryanodine receptor Ca^{2+} release channels via luminal and cytosolic Ca^{2+} sites. *Clin. Exp. Pharmacol. Physiol.* **34**, 889–896
33. Sikkel, M. B., Collins, T. P., Rowlands, C., Shah, M., O'Gara, P., Williams, A. J., Harding, S. E., Lyon, A. R., and MacLeod, K. T. (2013) Flecainide reduces Ca^{2+} spark and wave frequency via inhibition of the sarcolemmal sodium current. *Circ. Res.* **98**, 286–296
34. Li, L., Mirza, S., Richardson, S. J., Gallant, E. M., Thekkedam, C., Pace, S. M., Zorzato, F., Liu, D., Beard, N. A., and Dulhunty, A. F. (2015) A new cytoplasmic interaction between junctin and ryanodine receptor Ca^{2+} release channels. *J. Cell Sci.* **128**, 951–963
35. Chini, E. N. (2009) CD38 as a regulator of cellular NAD: a novel potential pharmacological target for metabolic conditions. *Curr. Pharm. Des.* **15**, 57–63
36. Aston, D., Capel, R. A., Ford, K. L., Christian, H. C., Mirams, G. R., Rog-Zielinska, E. A., Kohl, P., Galione, A., Burton, R. A., and Terrar, D. A. (2017) High resolution structural evidence suggests the sarcoplasmic reticulum forms microdomains with acidic stores (lysosomes) in the heart. *Sci. Rep.* **7**, 40620
37. Wächter, S. B., and Gilbert, E. M. (2012) β -Adrenergic receptors, from their discovery and characterization through their manipulation to beneficial clinical application. *Cardiology* **122**, 104–112
38. Munshi, C., Aarhus, R., Graeff, R., Walseth, T. F., Levitt, D., and Lee, H. C. (2000) Identification of the enzymatic active site of CD38 by site-directed mutagenesis. *J. Biol. Chem.* **275**, 21566–21571
39. Hara-Yokoyama, M., Kukimoto-Niino, M., Terasawa, K., Harumiya, S., Podyma-Inoue, K. A., Hino, N., Sakamoto, K., Itoh, S., Hashii, N., Hiruta, Y., Kawasaki, N., Mishima-Tsumagari, C., Kaitsu, Y., Matsumoto, T., Wakiyama, M., et al. (2012) Tetrameric interaction of the ectoenzyme CD38 on the cell surface enables its catalytic and raft-association activities. *Structure* **20**, 1585–1595
40. Zhang, H., Graeff, R., Lee, H. C., and Hao, Q. (2013) Crystal structures of human CD38 in complex with NAADP and ADPRP. *Messenger* **2**, 44–53
41. Dorian, P. (2005) Antiarrhythmic action of β -blockers: potential mechanisms. *J. Cardiovasc. Pharmacol. Ther.* **10**, S15–S22
42. Liu, Q., Graeff, R., Kriksunov, I. A., Lam, C. M. C., Lee, H. C., and Hao, Q. (2008) Conformational closure of the catalytic site of human CD38 induced by calcium. *Biochemistry* **47**, 13966–13973
43. Macgregor, A. T., Rakovic, S., Galione, A., and Terrar, D. A. (2007) Dual effects of cyclic ADP-ribose on sarcoplasmic reticulum Ca^{2+} release and storage in cardiac myocytes isolated from guinea-pig and rat ventricle. *Cell Calcium* **41**, 537–546
44. Terrar, D. A. (2015) The roles of NAADP, two pore channels and lysosomes in Ca^{2+} signaling in cardiac muscle. *Messenger* **4**, 23–33
45. Mészáros, L. G., Wrenn, R. W., and Váradi, G. (1997) Sarcoplasmic reticulum-associated and protein kinase C-regulated ADP-ribosyl cyclase in cardiac muscle. *Biochem. Biophys. Res. Commun.* **234**, 252–256
46. Xie, G. H., Rah, S. Y., Yi, K. S., Han, M. K., Chae, S. W., Im, M. J., and Kim, U. H. (2003) Increase of intracellular Ca^{2+} during ischemia/reperfusion injury of heart is mediated by cyclic ADP-ribose. *Biochem. Biophys. Res. Commun.* **307**, 713–718
47. Saito, A., Seiler, S., Chu, A., and Fleischer, S. (1984) Preparation and morphology of sarcoplasmic reticulum terminal cisternae from rabbit skeletal muscle. *J. Cell Biol.* **99**, 875–885
48. Holmberg, S. R., and Williams, A. J. (1990) The cardiac sarcoplasmic reticulum calcium-release channel: modulation of ryanodine binding and single-channel activity. *Biochim. Biophys. Acta* **1022**, 187–193
49. Lee, H. C. (2012) Cyclic ADP-ribose and nicotinic acid adenine dinucleotide phosphate (NAADP) as messengers for calcium mobilization. *J. Biol. Chem.* **287**, 31633–31640
50. Liu, Q., Kriksunov, I. A., Graeff, R., Munshi, C., Lee, H. C., and Hao, Q. (2005) Crystal structure of human CD38 extracellular domain. *Structure* **13**, 1331–1339
51. Liu, Q., Kriksunov, I. A., Graeff, R., Munshi, C., Lee, H. C., and Hao, Q. (2006) Structural basis for the mechanistic understanding of human CD38-controlled multiple catalysis. *J. Biol. Chem.* **281**, 32861–32869
52. Higashida, H., Egorova, A., Higashida, C., Zhong, Z. G., Yokoyama, S., Noda, M., and Zhang, J. S. (1999) Sympathetic potentiation of cyclic ADP-ribose formation in rat cardiac myocytes. *J. Biol. Chem.* **274**, 33348–33354
53. Graeff, R., Liu, Q., Kriksunov, I. A., Hao, Q., and Lee, H. C. (2006) Acidic residues at the active sites of CD38 and ADP-ribosyl cyclase determine nicotinic acid adenine dinucleotide phosphate (NAADP) synthesis and hydrolysis activities. *J. Biol. Chem.* **281**, 28951–28957
54. Cosker, F., Cheviron, N., Yamasaki, M., Menteigne, A., Lund, F. E., Moutin, M. J., Galione, A., and Cancela, J. M. (2010) The ecto-enzyme CD38 is a nicotinic acid adenine dinucleotide phosphate (NAADP) synthase that couples receptor activation to Ca^{2+} mobilization from lysosomes in pancreatic acinar cells. *J. Biol. Chem.* **285**, 38251–38259
55. Venturi, E., Pitt, S., Galfré, E., and Sitsapasan, R. (2012) From eggs to hearts: what is the link between cyclic ADP-ribose and ryanodine receptors? *Cardiovasc. Ther.* **30**, 109–116
56. Cockayne, D. A., Muchamuel, T., Grimaldi, J. C., Muller-Steffner, H., Randall, T. D., Lund, F. E., Murray, R., Schuber, F., and Howard, M. C. (1998) Mice deficient for the ecto-nicotinamide adenine dinucleotide glycohydrolase CD38 exhibit altered humoral immune responses. *Blood* **92**, 1324–1333
57. Galfré, E., Pitt, S. J., Venturi, E., Sitsapasan, R., Zaccari, N. R., Tsaneva-Atanasova, K., O'Neill, S., and Sitsapasan, R. (2012) Fkbp12 activates the cardiac ryanodine receptor Ca^{2+} -release channel and is antagonised by fkb12.6. *PLoS ONE* **7**, e31956
58. Noble, J. E., and Bailey, M. J. (2009) Quantitation of Protein. *Methods Enzymol.* **463**, 73–95
59. Clapper, D. L., Walseth, T. F., Dargie, P. J., and Lee, H. C. (1987) Pyridine nucleotide metabolites stimulate calcium release from sea urchin egg microsomes desensitized to inositol trisphosphate. *J. Biol. Chem.* **262**, 9561–9568
60. Jorgensen, W. L., Maxwell, D. S., and Tirado-Rives, J. (1996) Development and testing of the OPLS all-atom force field on conformational energetics and properties of organic liquids. *J. Am. Chem. Soc.* **118**, 11225–11236
61. Friesner, R. A., Banks, J. L., Murphy, R. B., Halgren, T. A., Klicic, J. J., Mainz, D. T., Repasky, M. P., Knoll, E. H., Shelley, M., Perry, J. K., Shaw,

- D. E., Francis, P., and Shenkin, P. S. (2004) Glide: a new approach for rapid, accurate docking and scoring: 1: method and assessment of docking accuracy. *J. Med. Chem.* **47**, 1739–1749
62. Halgren, T. A., Murphy, R. B., Friesner, R. A., Beard, H. S., Frye, L. L., Pollard, W. T., and Banks, J. L. (2004) Glide: a new approach for rapid, accurate docking and scoring: 2: enrichment factors in database screening. *J. Med. Chem.* **47**, 1750–1759
63. Halgren, T. A. (2009) Identifying and characterizing binding sites and assessing druggability. *J. Chem. Inf. Model.* **49**, 377–389
64. Jacobson, M. P., Friesner, R. A., Xiang, Z., and Honig, B. (2002) On the role of the crystal environment in determining protein side-chain conformations. *J. Mol. Biol.* **320**, 597–608
65. Jacobson, M. P., Pincus, D. L., Rapp, C. S., Day, T. J., Honig, B., Shaw, D. E., and Friesner, R. A. (2004) A hierarchical approach to all-atom protein loop prediction. *Proteins* **55**, 351–367
66. Shelley, J. C., Cholleti, A., Frye, L. L., Greenwood, J. R., Timlin, M. R., and Uchimaya, M. (2007) Epik: a software program for pKa prediction and protonation state generation for drug-like molecules. *J. Comput. Aided Mol. Des.* **21**, 681–691
67. Søndergaard, C. R., Olsson, M. H., Rostkowski, M., and Jensen, J. H. (2011) Improved treatment of ligands and coupling effects in empirical calculation and rationalization of pKa values. *J. Chem. Theory Comput.* **7**, 2284–2295
68. Li, H., Robertson, A. D., and Jensen, J. H. (2005) Very fast empirical and rationalization of protein pKa values. *Proteins* **61**, 704–721
69. Grosdidier, A., Zoete, V., and Michielin, O. (2011) SwissDock, a protein-small molecule docking web service based on EADock DSS. *Nucleic Acids Res.* **39**, W270–W277
70. Grosdidier, A., Zoete, V., and Michielin, O. (2011) Fast docking using the CHARMM force field with EADock DSS. *J. Comput. Chem.* **32**, 2149–2159

Synthesis of the Ca²⁺-mobilizing messengers NAADP and cADPR by intracellular CD38 enzyme in the mouse heart: Role in β -adrenoceptor signaling

Wee K. Lin, Emma L. Bolton, Wilian A. Cortopassi, Yanwen Wang, Fiona O'Brien, Matylda Maciejewska, Matthew P. Jacobson, Clive Garnham, Margarida Ruas, John Parrington, Ming Lei, Rebecca Sitsapesan, Antony Galione and Derek A. Terrar

J. Biol. Chem. 2017, 292:13243-13257.

doi: 10.1074/jbc.M117.789347 originally published online May 24, 2017

Access the most updated version of this article at doi: [10.1074/jbc.M117.789347](https://doi.org/10.1074/jbc.M117.789347)

Alerts:

- [When this article is cited](#)
- [When a correction for this article is posted](#)

[Click here](#) to choose from all of JBC's e-mail alerts

Supplemental material:

<http://www.jbc.org/content/suppl/2017/05/24/M117.789347.DC1>

This article cites 69 references, 24 of which can be accessed free at <http://www.jbc.org/content/292/32/13243.full.html#ref-list-1>

Synthesis of the Ca^{2+} -mobilizing messengers, NAADP and cADPR, by intracellular CD38 enzyme in mouse heart: role in β -adrenoceptor signaling

Wee K. Lin¹, Emma L. Bolton¹, Wilian A. Cortopassi^{3,4}, Yanwen Wang², Fiona O'Brien¹, Matylda Maciejewska¹, Matthew P. Jacobson⁴, Clive Garnham¹, Margarida Ruas¹, John Parrington¹, Ming Lei¹, Rebecca Sitsapesan¹, Antony Galione¹, Derek A. Terrar¹.

From the ¹Department of Pharmacology, University of Oxford, Mansfield Road, Oxford, OX1 3QT, UK

²Faculty of Biology, Medicine and Health, University of Manchester, Manchester, M13 9NT, UK

³Department of Chemistry, Chemistry Research Laboratory, University of Oxford, Mansfield Road, Oxford, OX1 3TA, UK

⁴Department of Pharmaceutical Chemistry, University of California, San Francisco, California 94158, United States

Running title: *Intracellular CD38 in heart and β -adrenoceptor signaling*

To whom correspondence should be addressed: Prof Derek A Terrar, Department of Pharmacology, University of Oxford, Mansfield Road, Oxford OX1 3QT, UK. Telephone: (+44) 1865271943; FAX: (+44) 1865271853; E-mail: derek.terrar@pharm.ox.ac.uk

Keywords: Ca^{2+} , beta-adrenoceptor, NAADP, cADPR, CD38, lysosomes, sarcoplasmic reticulum, heart, cardiac arrhythmia, cardiac hypertrophy

List of figures:

Figure S1: Effects of H_2O , Triton X-100, saponin and DMSO on the sea urchin egg homogenates bioassay

Figure S2: Alignment of sequence of human CD38 and mouse CD38

Figure S3: Immunolabeling of CD38 on rabbit atrial myocytes

Figure S4: Sheep cardiac SR showed ADP-ribosyl cyclase activities and NAADP production with properties similar to those of CD38

Figure S5: SAN4825 inhibited the production of NAADP in saponin-permeabilized single ventricular myocytes from mouse hearts

Figure S6: Binding conformations of SAN4825 at different pH

Figure S7: NAADP synthesis at pH 4.5 in cardiac membrane preparation from $\text{CD38}^{-/-}$ mice

REFERENCES

1. Altschul, S. F., Gish, W., Miller, W., Myers, E. W., and Lipman, D. J. (1990) Basic local alignment search tool., *J. molecular biology* **215**(3), 403–10. 10.1016/S0022-2836(05)80360-2
2. Altschul, S. F., Madden, T. L., Schäffer, A. A., Zhang, J., Zhang, Z., Miller, W., and Lipman, D. J. (1997) Gapped BLAST and PSI-BLAST: A new generation of protein database search programs
3. Munshi, C., Aarhus, R., Graeff, R., Walseth, T. F., Levitt, D., and Lee, H. C. (2000) Identification of the enzymatic active site of CD38 by site-directed mutagenesis, *J. Biol. Chem.* **275**(28), 21566–21571. 10.1074/jbc.M909365199

FIGURES




Figure S1**A****B****C**

Figure S1: Effects of H₂O, Triton X-100, saponin and DMSO on the sea urchin egg homogenates bioassay

A shows the homologous inactivation properties of NAADP in the SUEH, while **B** shows the ability of NAADP to mediate Ca²⁺ release in SUEH in the presence of H₂O, Triton X-100, saponin or DMSO. **C** shows the change in Ca²⁺-dependent fluorescence after the addition of H₂O, Triton X-100, saponin or DMSO in the SUEH bioassay. Note that all addition artifacts are relatively small (< 10 %) compared to the change in Ca²⁺-dependent fluorescence caused by 500 nM NAADP. These control experiments showed that neither the homologous inactivation properties nor the NAADP-mediated Ca²⁺ release was affected by permeabilizing agents or solvents used in our studies, supporting the reliability of NAADP measurement using this assay system under the conditions of our experiments. The concentrations of Triton X-100 (5×10^{-5} % v/v), saponin (5×10^{-6} % w/v) and DMSO (5×10^{-4} % v/v) that were used in these experiments correspond to the highest final concentrations present in SUEH in our studies. Data are expressed as the mean \pm SEM; n = 3 independent experiments.

Figure S2

```

MANYEFSQVSGDRPGCRLSRKAQIGLGVGLLVLIALVVGIVVILLRPRSL51 mouse
MANCEFSQVSGDKPCCRLSRRAQLCLGVSIILVLILVVVLAVVV---PRWRQ48 human
VWTGEPSTTKHFSIDIFLGRCLITYQILRPEMRDQNCQEILSTFKGAFVSKNP102 mouse
QWSGPGSTTKRFPETVLRARCVKYTEI-HPEMRHVDCQSVWDAFKGAFISKHP98 human
CNITREDYAPLVKLVTQTIPCNKTLFWSKSKHLAQYTWIQGKMFTLEDTL153 mouse
CNITEEDYQPLMKLGTQTVPCNKILLWSRIKDLAQFTQVQVQDMFTLEDTL149 human
LGYIADDLRWCGDPSTSDMNYVSCPHWSENCNNPITVFWKVISQKFAEDA204 mouse
LGYLADDLTWCGEFNTSKINYQSCPDWRKDCNNPVSVFWKTVSRRFAEAA200 human
CGVVQVMLNGSLREPFYKNSTFGSVEVFSLDPNKVHKLQAWVMHDIEGASS255 mouse
CDVVHVMLNGSRSKIIFDKNSTFGSVEVHNLQPEKVQTLQAWVIHGGREDSR251 human

NACSSSSSLNELKMIVQKRNMIIFACVDNYRPARFLQCVKNPEHPSC300 mouse
DLCQDPTIKELLESIIISKRNIIQFSCKNIYRPDKFLQCVKNPEDSSC296 human

```

Figure S2: Alignment of sequence of human CD38 and mouse CD38

The region used to raise the antibody was highlighted in yellow. The sequences were aligned using protein-protein Blast (1, 2) and the identical amino acid residues from mouse and human CD38 were labeled with red fonts. The amino acid residues surrounded by green boxes are those shown to be critical for the human CD38 enzymatic activities, determined by a site-mutagenesis study (3). The mouse catalytic residues corresponding to the Trp125, Glu146, Asp155, Trp189 and Glu226 of human CD38 are Trp129, Glu150, Asp159, Trp193 and Glu230, respectively. All catalytic residues are conserved between mouse and human CD38.

Figure S3




Figure S3: Immunolabeling of CD38 on rabbit atrial myocytes

Figure S3 shows the immunolabeling of CD38 with a striated pattern corresponding to the location of SR (and not that of surface membrane). Images shown are the representative staining of the majority (> 75%) observations (number of cells observed >20 isolated from 2 rabbit hearts). Antibodies used: rabbit polyclonal anti-CD38 antibody (sc-15362); scale bar: 20 μ m.

Figure S4




Figure S4: Sheep cardiac SR showed ADP-ribosyl cyclase activities and NAADP production with properties similar to those of CD38

A shows an example trace of synthesis of fluorescent cGDPR by sheep cardiac SR with NGD as substrate. This synthesis was suppressed in the presence of a CD38 inhibitor, nicotinamide. **B** shows the time course of NAADP synthesis in preparations enriched in heavy or light SR (similar in both preparations; $p = 0.9226$). **C** shows an example trace of NAADP synthesis with NADP as a substrate in the presence of NA, while synthesis was suppressed with either NADP or NA alone. Data in **B** are expressed as the mean \pm SEM; $n \geq 3$ independent experiments.

Figure S5





Figure S5: SAN4825 inhibited the production of NAADP in saponin-permeabilized single ventricular myocytes from mouse hearts

Figure S5 shows that the NAADP synthesis rate in saponin-permeabilized single ventricular myocytes from mouse hearts was significantly reduced in the presence of 3 μ M SAN4825. Both control and SAN4825-treated myocytes were first permeabilized with saponin (0.01 % w/v), preincubated with vehicle or drug for 30 min before being supplemented with 500 μ M NADP and 7 mM NA to initiate the NAADP production. Data are expressed as the mean \pm SEM; ** $p < 0.01$ relative to control, determined by Student's t-test. $n = 4$ independent experiments.

Figure S6**Figure S6: Binding conformations of SAN4825 at different pH**

The four images, each showing a SAN4825-CD38 complex, represent the predicted binding conformation of SAN4825 at the active site of mouse CD38 (PDB ID: 2EG9) under different conditions. The predictions were performed with consideration of the influence of pH on the protonation states of SAN4825 and on the amino acids of the CD38 protein, which were predicted by Epik and Propka 3.1, respectively. The predicted pKa values for SAN4825 are shown in top left. The protonation of the nitrogen (highlighted in blue) of the pyridine with a pKa value of 5.1 is likely to change in the pH range of 4.5 to 7.2. The term “protein at pH 4.5” refers to a mouse CD38 protein with protonated states predicted at pH 4.5 while the term “protein at pH 7.2” refers to a mouse CD38 protein with protonated states predicted at pH 7.2. The chemical structure above each image depicts the protonation state of SAN4825 considered.


Figure S7

Figure S7: Lack of *In vitro* NAADP synthesis at pH 4.5 in cardiac membrane preparation from $CD38^{-/-}$ mice

A and **B** show the time course and average rate of NAADP synthesis by cardiac membrane preparation from WT and $CD38^{-/-}$ mice at different pH. The NAADP synthesis of membrane preparation from $CD38^{-/-}$ mice remained reduced and comparable with those of $CD38^{-/-}$ mice at pH 7.2. Data are expressed as the mean \pm SEM; ** $p < 0.01$ relative to control, determined by Student's t-test. $n = 3$ independent experiments.