

# Cryptic infection of a broad taxonomic and geographic diversity of tadpoles by *Perkinsea* protists

Aur lie Chambouvet<sup>1,2</sup>, David J. Gower<sup>2</sup>, Miloslav Jirk <sup>3</sup>, Michael J. Yabsley<sup>4</sup>, Andrew K. Davis<sup>5</sup>, Guy Leonard<sup>1</sup>, Finlay Maguire<sup>2,7</sup>, Thomas M. Doherty-Bone<sup>2,8</sup>, Gabriela Bueno Bittencourt-Silva<sup>6</sup>, Mark Wilkinson<sup>2</sup>, Thomas A. Richards<sup>1,9</sup>

<sup>1</sup>Biosciences, University of Exeter, Geoffrey Pope Building, Exeter, EX4 4QD, UK. <sup>2</sup>Department of Life Sciences, The Natural History Museum, Cromwell Road, London SW7 5BD, UK. <sup>3</sup>Institute of Parasitology, Biology Centre, Academy of Sciences of the Czech Republic, Branisovska 31, 370 05 Ceske Budejovice, Czech Republic. <sup>4</sup>Warenell School of Forestry and Natural Resources and the Southeastern Cooperative Wildlife Disease Study, Department of Population Health, College of Veterinary Medicine, The University of Georgia, Athens, Georgia 30602, USA. <sup>5</sup>Odum School of Ecology, The University of Georgia, Athens, Georgia 30602, USA. <sup>6</sup>Department of Environmental Sciences, University of Basel, Klingelbergstr. 27, CH-4056, Switzerland. <sup>7</sup>Genomics, Ecology and Evolution, University College London, Darwin Building Gower Street, London WC1E 6BT, UK. <sup>8</sup>School of Geography & School of Biology, University of Leeds, Leeds, LS2 9JT, UK. <sup>9</sup>Canadian Institute for Advanced Research, CIFAR Program in Integrated Microbial Biodiversity, Toronto, ON, M5G 1Z8

Submitted to Proceedings of the National Academy of Sciences of the United States of America

The decline of amphibian populations, particularly frogs, is often cited as an example in support of the claim that Earth is undergoing its sixth mass extinction event. Amphibians appear to be particularly sensitive to emerging diseases (e.g. fungal and viral pathogens), yet the diversity and geographic distribution of infectious agents are only starting to be investigated. Recent work has linked a previously undescribed protist with mass-mortality events in the USA, in which infected frog tadpoles have an abnormally enlarged yellowish liver filled with protist cells of a presumed parasite. Phylogenetic analyses revealed that this infectious agent was affiliated with the *Perkinsea*; a parasitic group within the alveolates exemplified by *Perkinsus* sp. a 'marine' protist responsible for mass-mortality events in commercial shellfish populations. Using small subunit rDNA sequencing, we developed a targeted PCR protocol for preferentially sampling a clade of the *Perkinsea*. We test this protocol on freshwater environmental DNA revealing a wide diversity of *Perkinsea* lineages in these environments. Then we used the same protocol to test for *Perkinsea*-like lineages in livers of 182 tadpoles from multiple families of frogs. We identified a distinct *Perkinsea* clade, encompassing a low level of SSU rDNA variation different from the lineage previously associated with tadpole mass-mortality events. Members of this clade were present in 38 tadpoles sampled from 14 distinct genera/phylogroups, from five countries across three continents. This provides the first evidence that *Perkinsea*-like protists infect tadpoles across a wide taxonomic range of frogs in tropical and temperate environments, including oceanic islands.

Frog decline | emerging disease | parasite | alveolates | molecular diversity

## Introduction:

It is widely recognized that amphibians are among the most threatened animal groups, for example in 2008 32% of species were listed as 'threatened or extinct' and 42% were listed as in decline (<http://www.iucnredlist.org/initiatives/amphibians/analysis> - accessed 29-Oct-2014) (1, 2). The main causes of amphibian decline have been identified as habitat loss, environmental change, and the introduction of non-native species (e.g. (3-6)). Emerging infectious diseases have also been shown to play a key role in many amphibian declines, for example the chytrid fungal pathogen *Batrachochytrium dendrobatidis* has caused mass mortality events (MMEs) in Australia, Europe and across the Americas (e.g. (7-9)). MMEs have also been associated with infection by *Ranavirus* in, for example, the United Kingdom (UK), United States of America (USA), and Canada (10, 11). Recent work has linked local MMEs in the USA with the infection of larval frogs (tadpoles) of the genera *Lithobates* and *Acris* by a protist (see SI Appendix, Fig. S1 - (12, 13)). In 2006, histological examinations of tadpole tissues revealed the presence of thousands of


small spherical cells preferentially infecting livers of tadpoles of the Southern Leopard Frog (*Lithobates sphenoccephalus*, formerly *Rana sphenoccephala*) sampled from a MME in Georgia (USA) (12). SSU rDNA PCR and direct-amplicon sequencing combined with phylogenetic tree reconstruction showed that a lineage of protists closely related to *Perkinsus*, a parasite of marine bivalves (14), was the likely infectious agent (12).

*Perkinsea* alveolates were first described as being affiliated with the Apicomplexa (14, 15), which includes important human pathogens such as *Toxoplasma gondii* and *Plasmodium* spp. (the causative agents of malaria). Phylogenetic analysis has shown that *Perkinsea* are a deeply divergent sister-group of dinoflagellate alveolates (16). Only three representative groups of *Perkinsea* were previously described morphologically and taxonomically: *Perkinsus* spp., parasites of marine bivalves (e.g. oysters and clams), *Parvilucifera* spp., parasites of dinoflagellates, and *Rastrimonas subtilis* (previously *Cryptophagus subtilis*), parasites of cryptophyte algae (17-20). However, environmental sequence analysis has considerably expanded the known diversity of *Perkinsea*-like organisms (21-25). The putative causal agent of the *L. sphenoccephalus* MME lies within a clade (monophyletic group) of this environmental sequence diversity (Fig. 1A - (12)) named here, for convenience, Novel Alveolate Group 01 (NAG01).

## Significance

Amphibians are among the most threatened animal groups. Population declines and extinctions have been linked, in part, to emerging infectious diseases. One such emerging disease has been attributed to *Perkinsea*-like protists causing mass mortality events in the USA. Using molecular methods we evaluated the diversity of *Perkinsea* parasites in livers sampled from a wide taxonomic range of tadpoles from six countries across three continents. We discovered a previously unidentified phylogenetically distinct infectious agent of tadpole livers present in a broad range of frogs from both tropical and temperate sites and across all sampled continents. These data demonstrate the high prevalence and global distribution of this newly identified infectious protist.


## Reserved for Publication Footnotes



**Fig. 1. A.** Phylogenetic tree of Perkinsia SSU rDNA sequences focusing on the NAG01 group that includes two separate phylogenetic groups recovered from tadpole liver tissue samples. The phylogeny is estimated from a masked alignment consisting of 292 taxa and 776 characters. Bayesian posterior probability, LogDet distance bootstrap (1,000 pseudoreplicates) and maximum likelihood bootstrap (1,000 pseudoreplicates) values are added to each node using the following convention: support values are summarised by black circles when all are equal to or greater than 0.9/80%/80% and white circle when this is not the case but all values are equal to or greater than 0.6/50%/50%. Five sequences of *Amoebophrya* sp. were used as outgroup. Each square represents one environmental operational taxonomic unit (OTU) and the provenance of the OTUs are indicated by coloured boxes (see key for the detail of sample provenance, SI Appendix Table S2 provides more details on the environments sampled). A red diamond indicates the individual clone sequences from the *L. sphenoccephalus* 2006 mass mortality event in Georgia, USA (12). A subset of the published environmental sequences have been reduced to representative triangles (see SI Appendix, Table S10 for detail of each environmental clade). **B.** Representation of the V4 hyper-variable region of the template SSU rDNA and the relative position of the different primers used in this study (not to scale). **C.** Histogram representing the percentage of clones per clade A, B and C within each host superfamily from infected tadpoles.  $n_1$  represents the number of total clones sequenced per host superfamily/number of infected tadpoles per host superfamily. **D.** Geographical distribution of clade A, B and C. Pie charts represent the proportion of clones per clade A, B and C in each of the five geographical locations where NAG01 was detected (UK, French Guiana, São Tomé and Príncipe, Cameroon and Tanzania).  $n_2$  represents the number of clones sequenced per geographical location/number of infected tadpoles per geographical location.


In this study, we develop SSU rDNA primers that preferentially target the NAG01 group. We use these PCR primers

for targeted screening of NAG01 diversity in both tropical and temperate freshwater environments demonstrating the efficacy



**Fig. 2. A maximum likelihood phylogenetic circle tree (inverted) showing clades A, B and C and demonstrating the provenance of NAG01 phylotypes detected in frog liver tissue.** The RAxML phylogeny is estimated from a masked alignment consisting of 177 and 806 characters from NAG01 sequences from infected tadpoles. Branches proportionally shortened by  $\frac{1}{2}$  are labelled with a double-slashed line and by  $\frac{3}{4}$  with two double-slashed lines. Each clone sequenced detected from the same liver sample are connected. The colours of the connected lines were defined by the geographical location of tadpoles sampled (see the key). Black lines on the branches mark sequence variation confirmed across multiple samples and therefore cannot be the product of PCR error during clone library construction.

of this PCR protocol and expanding the diversity of Perkinsa- like sequences sampled from freshwater environments. Using the



**Fig. 3. Bayesian 16S rDNA phylogenetic tree of tadpole diversity sampled in this study with histograms showing prevalence of NAG01 detection.** The phylogeny is inferred from a masked alignment consisting of 247 taxa and 440 characters. Bayesian posterior probability (6,000 MCMC generation samples), LogDet distance bootstrap (1,000 replicates) and maximum likelihood bootstrap (1,000 replicates). Support values are summarised by black circles when all are equal to or greater than 0.9/80%/80% and white circle when this is not the case but all values are equal to or greater than 0.6/50%/50%. Sequences of *Ambystoma* sp. and *Pleurodeles* sp. (salamanders) were used as outgroup. Some frog species with multiple non-identical 16S rDNA sequences recorded in GenBank are retained. The colour-coded histogram represents the number of NAG01 negative tadpole samples (uncoloured bars) and the number of NAG01 positive samples (coloured bars). Each colour corresponds to the tadpole's country of origin as detailed in the key. The super-family and sub-order of the tadpoles tested is indicated on the histogram. The circled star indicates the host species described by Davis *et al.* (2007) during the 2006 USA mortality event.

same protocol we also detected NAG01 from livers of a wide taxonomic diversity of tadpoles from five countries across three

continents. Within our study sample we find evidence of infection by lineages different to the Perkins-like *L. spheenocephalus*



pathogen identified by Davis *et al.* (2007), suggesting multiple *Perkinsea* lineages infect tadpoles.

## Results and Discussion

### *Development of a targeted PCR assay for Perkinsea-like infection of frog tissue*

Culture-independent environmental DNA methods can be useful for detecting microbial lineages from the environment including from the tissues of plants and animals (e.g. (26-28)). To investigate the prevalence and diversity of *Perkinsea*-like parasites infecting tadpoles we designed two independent sets of PCR primers targeting ~800 base pairs (bp) of the small subunit (SSU rRNA) encoding gene including the V4 variable region (29) (Fig. 1B & SI Appendix, Table S1 for primers used in this study). The primers were designed to amplify the rDNA sequences of the wider NAG01 group (Fig. 1A), including the previously sampled SSU rDNA sequence of the *Perkinsea*-like infectious agent of the Southern Leopard Frog (12). To investigate the specificity of these two pairs of primers we performed PCR on ten environmental DNA samples from three tropical (French Guiana) and seven temperate (UK) planktonic freshwater samples (SI Appendix, Table S2). A total of 248 clones were sampled from the UK temperate water masses and 60 clones from the French Guiana water masses (SI Appendix, Table S2). Among these 308 clones, sequences from 240 clustered together into 46 non-identical sample specific NAG01 sequences (SI Appendix, Table S2 & S3) that were included in the phylogenetic analysis (Fig. 1A). The remaining 68 clones were non-NAG01 sequences encompassing a mixed assemblage of Fungi, Cryptophyta and nematode sequences. Based on these results, our PCR protocol was judged as adequate for preferentially targeted environmental clone library analyses of NAG01, including the previously identified *Perkinsea*-like *L. spheocephalus* infectious agent and seven environmental DNA sequences present in the GenBank database that were recovered from freshwater planktonic samples ((30-32) - see SI Appendix Table S4).

### *Investigating the global prevalence of NAG01 infections in tadpoles.*

Using the same PCR protocol used to detect NAG01 from freshwater environmental samples, we screened for NAG01 sequences in DNA extractions from liver samples dissected from 182 ethanol preserved tadpoles. We sampled tadpoles from French Guiana (80 individuals from eight sampling localities), Cameroon (37 from 14), Tanzania (15 from one), the island of São Tomé (four from one), the UK (40 from five) and the Czech Republic (six from three), of which 38 (21%) were PCR positive for NAG01. See SI Appendix Table S5 for more details, including: GPS location, sampling date and description of environment where the tadpoles were found. For each positive sample, the PCR reaction was repeated three times and the amplicons pooled. Each PCR product was checked on a 1% agarose gel for the presence of a single band of ~800bp. The PCR product was then purified, cloned and sequenced. We sequenced six clones using M13F primers from each PCR positive liver sample resulting in 228 sequences. Based on preliminary sequence analysis of the forward reads, all NAG01 SSU sequences that were unique were double-strand sequenced. This encompassed a minimum of four clones per tadpole liver DNA clone library even though most libraries included fewer non-identical clones (see SI Appendix Table S6).

Phylogenetic analysis demonstrates that all 177 sample-specific, unique NAG01 sequences recovered from tadpole livers form three discrete strongly supported and closely related clades (labelled A, B, and C on Fig. 1A). Clades A, B and C do not correspond directly with host taxon or geographic origin (Fig. 1C/D, 2A/B), a similar result to that described for *Perkinsus* sp. (33) parasites of marine bivalves. Interestingly, 21 of the 38 (55%)

NAG01-positive tadpole liver samples yielded sequences from both clade A and B (Fig. 2), suggesting that these livers harboured representatives from different non-clonal strains/species or that there is intranuclear SSU rDNA variation within NAG01 genomes sampled. Indeed, there is less than 3% nucleotide variation among clade A, B and C sequences (SI Appendix, Fig. S2) consistent with intranuclear variation such as that observed in SSU-5.8S-LSU paralogues and pseudogenes. For example, variant copies of the rRNA gene are known to occur in alveolates (34, 35) with SSU rRNA gene paralogues with 11% difference (36) transcribed at different stages of *Plasmodium* spp. life cycle (37).

It is possible that the NAG01 DNA detected could have arisen from contamination from the environment and/or gastrointestinal tissue in the liver samples. However, this was judged unlikely because all of the NAG01 SSU sequences detected from tadpole liver sequences grouped into one discrete phylogenetic subgroup, while the primers used were capable of detecting a wide diversity of *Perkinsea*-like sequences (Fig. 1A). To further test for cases of environmental contamination, DNA was extracted from tail samples (muscle and fin) taken from the same 182 tadpoles as a control to identify possible sources of non-tissue specific PCR detection or environmental contamination of NAG01 sequences. Although infections of an unknown alveolate-like parasite have been identified in the muscles of adult frogs (38), the tadpoles sampled here showed no evidence of disease progression so this was judged as an adequate control to identify cases of environmental and wider animal tissue contamination. All controls were negative for the two primer-paired NAG01 specific PCR protocols, suggesting that detection of NAG01 was not an artefact of environmental contamination but instead a tissue-specific signal consistent with infection of a *Perkinsea*-like protist associated with the liver of these tadpoles. Interestingly, these liver-derived NAG01 sequences are closely related to some of the sequences recovered from filtered plankton environmental DNA sample sequences, suggesting that this protist group is found both associated with tadpoles and either as free living stage or associated with additional microbial hosts. Indeed, experimental manipulations have shown that infection of putative members of the NAG01 clade, likely occur through ingestion of spores and/or zoospores from the watercolumn (39).

### *Diversity of Perkinsea parasites recovered from liver tissues from USA mass mortality event*

Sequences generated from the tadpole livers sampled here are from a different NAG01 *Perkinsea* lineage than that detected in liver tissues of *L. spheocephalus* tadpoles sampled from a MME in Georgia (USA) in 2006 (see Fig. 1A). Only formalin-fixed, paraffin-embedded liver samples were available from this event (12). We successfully performed two DNA extractions from a historical sample. The standard NAG01 primers failed to generate an amplicon from both DNA samples, most likely because the template DNA was highly fragmented. Consequently we targeted a shorter template using the 300F-B forward primer in combination with a eukaryotic general reverse primer 600R (SI Appendix, Table S1), we amplified 414 bp of the SSU rDNA. PCR reactions were repeated three times for both DNA samples, amplicons pooled and cloned separately for the two DNA samples. In total, across the two DNA samples, we sequenced 45 clones in forward and reverse direction. Phylogenetic analysis revealed that 44 clones represented four unique sequences that are closely related to the published sequence of the *L. spheocephalus* infectious agent (EF675616) and one unique sequence closely related to NAG01 clade B (Fig. 1A).

### *Barcode sequencing reveals a wide host-taxon diversity and biogeography for NAG01*

Precise taxonomic identification of tadpoles using morphological characters can be difficult, especially in geographic re-

gions with high species diversity. We amplified a mitochondrial 16S rDNA barcode, shown to be effective for higher taxonomic assignment of amphibians (40, 41), for all 182 sampled tadpoles. The majority ( $n = 175$ ) of the tadpoles sampled are members of the Neobatrachia (which comprises  $> 95\%$  of extant frogs), whereas the other seven are tadpoles of the Pipidae i.e. *Xenopus* (Fig. 3B). We note that, although the pipid tadpoles sampled all tested negative, increased sampling is required to more confidently determine the presence/prevalence of this protist in this group of frogs.

NAG01 DNA was detected in tadpoles of two of the largest groups of neobatrachians, with 6% of sampled Hyloidea ( $n = 102$ ) infected and 42% of Ranoidea ( $n = 73$ ) infected (Fig. 3). In separate studies a Perkinsea-like parasite linked to local mortality events was detected using histology in seven species of Ranoidea (*Lithobates sphenoccephalus*, *L. capito*, *L. catesbeiana*, *L. catesbeianus*, *L. heckscheri*, and *L. sylvaticus*) and one species of Hyloidea (*Acris gryllus*) ([http://www.nwhc.usgs.gov/publications/quarterly\\_reports](http://www.nwhc.usgs.gov/publications/quarterly_reports) - accessed 29-Oct-2014) (SI Appendix, Fig. S1). Similar dissection based approaches have also shown related infectious agents present in a wide diversity of ranids as well as *Acris gryllus*, *Hyla femoralis*, *Hyla gratiosa*, *Pseudacris ornata* and *Gastrophryne carolinensis* (39). In the absence of molecular data it is not clear if these infections were of protists from NAG01 clades A, B, or C reported here, the *L. sphenoccephalus* parasite detected by Davis *et al.* (2007), or a lineage not yet sampled for DNA analysis. However, the NAG01 sequences from tadpole livers detected in this study were recovered in five of the six countries sampled including both tropical and temperate environments and an oceanic island. Mean prevalence estimate per country and 95% confidence intervals using the Jeffreys method was: 3% (four of 80, 95% CI of 0.4-34%) in French Guiana (eight event/site sampled), 55.2% (13 of 37, 11-78% CI) in Cameroon (13 event/site sampled), 93.3% (14 of 15, 72-99% CI) in Tanzania (one event/site sampled), 100% on the Island of São Tomé (four of four, 73-100% CI - one sampling event/site), and 9% (three of 40, 1-40% CI) in the UK (three event/site sampled). Taken together these data suggest a high prevalence and broad spatial distribution of infection by a specific sub-clade of NAG01 Perkinsea-like protists in neobatrachian tadpole populations specifically of the superfamily Ranoidea. Information connecting the putative phylogeny inferred taxonomy of the host tadpole to the presence of the NAG01 sequence-type is given in SI Appendix Table S5 to supplement the results shown in Fig. 3B.

#### NAG01 Perkinsea-like protist and disease

All our tadpole field samples were preserved in ethanol and as such we could not attempt to purify NAG01 clade A, B or C into culture. Infection by Perkinsea-like parasites identified using histological and/or dissection microscopy techniques have been reported as visible from Gosner life cycle 24-42 (39, 42). In some cases the disease phenotype of tadpoles infected by Perkinsea-like parasites has previously been described as bloated, lethargic and showing cutaneous haemorrhages (12, 13, 42, 43), although we note that this is not diagnostic for Perkinsea disease of tadpoles, because other diseases can cause similar pathologies. The majority of the 38 infected tadpoles analysed in this study were sampled from early Gosner stages, with the majority being of, or close to, Gosner stage 25 with one at Gosner stage 42 (see SI Appendix Table S5). The infected tadpoles showed no gross morphological symptoms of disease. A more precise definition of the disease in the cases of Perkinsea infection is infiltration of the liver and other visceral organs by large numbers of Perkinsea-like organisms (12, 38). Indications of tissue level disease in samples showing molecular evidence for the presence of NAG01 were sought through histology of liver with H&E staining of five  $\mu\text{m}$  sections from representative samples (two from the French

Guiana, three from Cameroon and three from the Tanzania - for examples see SI Appendix Fig S3). These data showed no identifiable tissue damage consistent with disease and no cells attributable to the NAG01 microbes detected using molecular methods, demonstrating that these infectious agents are either a small cellular form and/or badly preserved in histological sections, or alternatively the population of NAG01 is very low suggesting a low infection intensity in these tadpoles. Thus we cannot confirm that the infectious protists identified here are pathogenic, either because: 1) the tadpoles were all sampled in an early phase of disease progression and/or the infectious protists are currently dormant, 2) NAG01 clades A, B or C detected in this study and which branch in a different part of the phylogenetic tree to previously reported disease causing Perkinsea of tadpoles (Fig. 1A and (12)) have a limited or absent disease pathology, and/or 3) disease is only caused in association with other infectious pathogens such as *Ranavirus* (43) or other forms of host stress (44). Other parasitic Perkinsea such as *Perkinsus sensu stricto* (parasites of bivalves), are also widely geographically distributed but infection and catastrophic host population MMEs are localised, with pathogenicity related in part to abiotic factors (33, 45-47).

These data demonstrate that the NAG01 protists detected in tadpole livers in this study are: 1) not a ubiquitous agent or contaminant but instead are liver associated, 2) represent a closely related phylogenetically distinct subgroup within NAG01, 3) lie within the phylum Perkinsea, for which all known taxa are potential parasites (e.g. (18, 48, 49), and 4) are prevalent in a range of tadpole developmental stages (Gosner stages 25-42 - SI Appendix, Table S5). Although frog MMEs associated with a Perkinsea-like parasite has only been recorded in the USA, our results demonstrate that a greater diversity of Perkinsea-like protist infections of tadpoles are widespread. There are increasing efforts to monitor the health of wild amphibian populations (50-52), with justifiable focus on fungal chytrid pathogens and *Ranavirus*, both of which have been identified as causing disease in adult frogs (6-8, 10). Further studies on the aetiology of tadpole (and other larval amphibian) infections are necessary to understand the impact of these protists on amphibian populations and to inform conservation planning.

## Materials and Methods

### Developing group specific primers

Based on a multiple sequence alignment of available Perkinsea SSU rDNA sequences (see SI Appendix Table S4 & S7) assembled using ARB (53) sets of 'NAG01-specific primers' were designed to recover a central portion of the SSU rRNA encoding gene including the variable V4 region (Fig. 1B & SI Appendix, Table S1). Specificity of the PCR primers was checked first *in silico* by submitting sequences to the NCBI nr DNA (using Primer-BLAST - May 2014) and SILVA (TestProbe search - May 2014) databases (54) and, second, *in situ* using an environmental DNA clone library approach.

### Environmental DNA sampling of freshwater environments

Water samples from the surface of the water column were collected from multiple freshwater environments in the UK and French Guiana (details of each sample are given in SI Appendix, Table S2). For the sampling in French Guiana, water samples were pre-filtered through 10  $\mu\text{m}$  polycarbonate filters (Merck Millipore, Darmstadt, Germany) and the filtrate then serially filtered through 5  $\mu\text{m}$  then 2  $\mu\text{m}$  polycarbonate filters to collect size-specific subsections of the microbial community. This process was conducted until each filter became saturated (the volume of water filtered for each sample is reported in SI Appendix, Table S2). Saturated filters were then submerged in RNAlifeGuard (MoBio, Carlsbad, USA) and then stored at  $-20^\circ\text{C}$  for two weeks before being transported back to the UK at ambient temperatures and then finally stored at  $-80^\circ\text{C}$ . The UK samples were processed in a similar manner but serially passed through 20  $\mu\text{m}$  and then 2  $\mu\text{m}$  or 0.2  $\mu\text{m}$  filters and transformed directly to storage in  $-80^\circ\text{C}$ . DNA extraction was performed using the PowerWater DNA Isolation Kit (MoBio, Carlsbad, USA) using the protocol recommended by the manufacturer. Details of filtrations size for positive detection of NAG01 are given in SI Appendix (SI Appendix, Table S2) and indicate presence of this group in both the 0.2-20  $\mu\text{m}$  and 2-5  $\mu\text{m}$  size fraction samples.

### Sampling of tadpole tissue for NAG01 molecular screening

A total of 182 ethanol preserved tadpoles was examined, details of the phylogenetic diversity sampled are given in Fig. 3B and the highest BLAST hit for each sample in the NCBI nr database (accessed October 2014)

is given in SI Appendix Table S8. SI Appendix Table S5 provides details of the environmental provenance of the tadpoles. Tadpoles preserved whole in ethanol were dissected using sterile tools. A piece of liver of each tadpole was removed (taking care not to pierce the gut) and placed in a fresh tube of ethanol. Concurrently, a similarly sized piece of tail was excised and placed in a separate tube of ethanol. Total DNA was extracted from all tadpole tissue samples using the Blood and Tissue DNeasy extraction kit (Qiagen, Limburg, Netherlands) following the manufacturer's protocol with an overnight lysis and incubation step.

#### SSU rDNA clone libraries from environmental DNA and tadpole liver DNA

Environmental DNA, tadpole liver and tadpole tail DNA extractions were used as a source template to construct NAG01-specific SSU rDNA gene clone libraries. Two sets of primers were used including two NAG01-specific forward primers paired with two general eukaryotic reverse primers (see Fig. 1B). For every PCR reaction we included a negative control (distilled H<sub>2</sub>O). All PCR amplification reactions were performed in 25 µl total volume containing 8 ng of DNA and PCR MasterMix (Promega, Madison, USA). Cycling reactions were as follows: 2 min at 95 °C, followed by 35 cycles of 30 s at 95 °C, 30 s at 55 °C, 120 s at 72 °C and with an additional 10 min extension at 72 °C. For each template, three independent PCR reactions were performed, mixed together, purified using the Wizard SV Gel and PCR Clean-Up System kit (Promega, Madison, USA) and cloned using the Strataclone PCR cloning kit (Stratagene, Santa Clara, USA) according to the manufacturer's instructions. Clones were blue/white screened and a subset selected for PCR using M13F and M13R primers, which flank the vector insertion site. Clones showing the presence of an insert of the correct size were sequenced using M13F primers. For each environmental clone library, a minimum of 20 clones was sequenced. All the tadpole tail samples were PCR negative. For each tadpole liver sample with a positive PCR result we sequenced six clones using the M13F primer. All M13F NAG01 SSU sequences that were unique in one or more position were double strand sequenced.

#### Mitochondrial encoded 16S SSU rRNA gene barcoding of tadpoles

Tadpole liver DNA extracts were subject to PCR using the 16Sar-F and 16Sar-R primers (40) which amplify a ~600 bp region of the 16S SSU rRNA encoding gene. PCR amplification reactions were performed in 50 µl total volume containing 8 ng DNA and PCR MasterMix (Promega, Madison, US). Cycling reactions were as follows: 2 min at 95 °C, followed by 25 cycles of 30 s at 95 °C, 30 s at 55 °C, 120 s at 72 °C with an additional 10 min extension at 72 °C. PCR products were checked on 1% agarose gel and purified using the Wizard SV Gel and PCR Clean-up System kit (Promega, Madison, US).

#### DNA extraction and SSU rDNA clone library construction from formalin-fixed, paraffin-embedded liver tissue from the 2003 USA Southern Leopard Frog MMEs

Frozen and/or ethanol preserved tadpole tissue from the 2006 mortality event (12) was unavailable due to theft of copper wiring from the freezers of the Yabsley laboratory resulting in loss of these samples. Thus only formalin-fixed and paraffin-embedded tissues used for microscopy sections were available for molecular analysis. Two liver sections were excised from the paraffin block using sterile scalpel blades. Each section was incubated with xylene at 50 °C for 3 min until the paraffin had dissolved. After this process the liver section still constituted a compact tissue aggregate. The xylene solution was removed by pipetting and the tissue sample washed twice using pure ethanol and dried for 15 min at 37 °C. DNA was then extracted using the DNeasy Blood and Tissue kit (Qiagen, Limburg, Netherlands) protocol with an overnight incubation in 100 µl of lysis buffer at 55 °C. The DNA extractions were conducted twice on the two different formalin-fixed, paraffin-embedded liver samples.

The DNA extractions were checked using a 2100 Bio-analyzer (Agilent Technologies, Santa Clara, US) demonstrating highly fragmented template DNA (similar to ancient DNA samples) with an average fragment size of approximately 160 bp. We thus amended our PCR protocol to target a shorter amplicon using the forward NAG01-specific 300F-B primer with the eukaryotic general reverse primer 600R (see SI Appendix, Table S1 for details). The PCR amplification reaction was performed in 25 µl total volume of PCR MasterMix (Promega, Madison, USA). Cycling reactions were as follows: 2 min at 95 °C, followed by 35 cycles of 30 s at 95 °C, 30 s at 59 °C, 120 s at 72 °C with an additional 10 min extension at 72 °C. For each clone library, three independent PCRs were completed, mixed together and cloned. Two clone libraries were constructed using Strataclone cloning kit (Stratagene, Santa Clara, US) according to the manufacturer's instructions. Twenty independent clones with insertions of appropriate size were selected and sequenced in both directions using M13F and M13R primers.

#### Sequencing and assembly

All sequencing was performed externally by Beckman Coulter Genomics (Hertfordshire, UK). The final sets of sequences were trimmed to regions of high sequencing quality; vector sequences removed, sequence reads assembled into a contiguous sequence using Sequencher (Genecodes, Ann Arbor, USA) and ambiguous sites corrected.

#### Multiple sequence alignment and phylogenetic analysis

Our NAG01 clone library sequencing resulted in 177 sample-specific unique clones from the tadpole liver samples (SI Appendix, Table S6), 46 sample-specific unique clones from the environmental DNA samples and five sample-specific unique clones from formalin-fixed, paraffin-embedded

liver samples (see SI Appendix Table S3 for details). These sequences were assembled into a multiple sequence alignment with 59 Perkinsea-like sequences and *Amoebophrya* sp. SSU rDNA outgroup sequences (HQ658161, HM483395, HM483394, AY208894, AF472555) recovered from NCBI nr database (accessed May 2014 – see SI Appendix, Table S4 & S7). The sequences were aligned using MUSCLE (55) available via the graphical multiple sequence alignment viewer Seaview v4.2.12 (56), employing default settings. The alignment was then checked and masked manually in Seaview resulting in a data matrix of 292 sequences and 776 alignment positions. We note that the alignment included some partial database sequences in order to best sample the diversity of sequences sampled previously. However, Perkinsea-like sequences with the major central portion of the sequence absent were excluded from the final analysis (e.g. EUY162621, EUY162622, EUY162623). This data matrix contained 487 variable alignment positions (excluding alignment positions with gaps) and with 571 parsimony informative sites (including gaps).

All Perkinsea SSU rDNA sequences generated as apart of this study have been deposited in GenBank (see SI Appendix Table S3 & S6 for details). The Perkinsea alignment is available in the Seaview (56) Mase format with the alignment mask information retained and available at DOI: 10.5281/zenodo.12712.

The 182 tadpole 16S SSU rDNA sequences were aligned with a collection of frog 16S SSU rDNA sequences (SI Appendix, Table S9) using the same approach described for the NAG01 sequences and using MUSCLE (55) via Seaview v4.2.12 (56). All tadpole 16S SSU rDNA sequences generated in this study were BLASTn searched against the NCBI nr database (accessed May 2014) and the most similar named frog sequence based on sequence similarity was noted (SI Appendix, Table S8). Preliminary phylogenies were compared to published trees (57) and additional frog 16S SSU rDNA sequences representing intermediate branches (arbitrarily selected) were added to the phylogeny. Sequences of Caudata (salamanders) were chosen as outgroup for the phylogeny: *Pleurodeles waltl* (DQ283445), *Pleurodeles nebulosus* (DQ092266), *Ambystoma mexicanum* (EF107170) and *Ambystoma tigrinum* (DQ283407). The alignment was then checked and masked manually in Seaview resulting in a data matrix of 247 sequences and 440 alignment positions with 225 variable alignment positions (excluding alignment positions with gaps) and with 232 parsimony informative sites (including sites with gaps in the masked data matrix).

All frog SSU rDNA sequences generated as apart of this study have been deposited in GenBank (see SI Appendix Table S5 for details). The amphibian alignment is available in the Seaview (56) Mase format with the mask information retained and available at DOI: DOI: 10.5281/zenodo.12712.

The best-fitting nucleotide substitution model for each alignment was determined using the information criterion and likelihood ratio tests implemented in Modelgenerator v0.85 (Keane et al., 2010). For Perkinsea and amphibian alignments, GTR+Γ and GTR+I+Γ models were selected, respectively. The α parameter for the Γ distributions were 0.38 and 0.30, respectively with eight discrete rate categories while the I parameter for the amphibian alignment was 0.28. These parameters, where possible, were input into a Bayesian analysis using MrBayes v3.1.2 (58), i.e.: lset, nst=6 rates=gamma (or invgamma in the case of the amphibian alignment) and in both cases we included the covarion parameter search. Two independent runs of four Metropolis-Coupled (MCMC) chains (with a heat parameter of 2) were run for 2,000,000 generations. Trees were sampled every 250 generations. In both analyses, the MCMC searches had converged within the first 25% of the generations sampled, as such the first quarter of the search results were discarded (as the burnin). Convergence between the runs and burn-in were assessed using Tracer v1.6 (<http://tree.bio.ed.ac.uk/software/tracer>). The consensus topologies and posterior probabilities of each node were then calculated from the remaining sampled trees.

Support for the Bayesian tree topology was evaluated using two bootstrap methods and the Bayesian posterior probabilities (PP) from the MrBayes runs. Bootstrap support values were estimated using 1) RAXML v8.0.3 (59) with 1,000 pseudo-replicates, and 2) LogDet distance analysis with 1,000 pseudoreplicates using a BioNJ search method (available through Seaview v4.2.1). This second bootstrap analysis was conducted for comparison because it uses a method which minimises artefacts arising from biases in base composition across the alignment (60).

To further investigate the phylogenetic diversity of the NAG01 sequences recovered from the tadpole livers we resampled the alignment mask specifically focusing on sequences recovered from clades A, B and C with the aim of maximising unambiguously aligned sites. This analysis excluded all environmental sequences and only retained tadpole associated Perkinsea sequences from clades A, B and C. This resulted in a data matrix of 177 sequences and 806 alignment positions. This data matrix was analysed with Modelgenerator, selecting the GTR + Γ (α parameter of 1.18) using the standard Akaike information criterion. The phylogeny was then estimated using RAXML v8.0.3 (59) using the GTR + Γ substitution model. For this analysis we aimed to calculate a single tree that best displays the phylogenetic relationships of the NAG01 sequences sampled from tadpole livers. This alignment encompassed little sequence variation, and tree searches did not result in a consensus tree with consistently high/moderate bootstrap support values. This phylogenetic analysis therefore primarily serves to demonstrate the distribution of different SSU-types across the different tissue samples



and not to present a resolved phylogeny. To display co-detection of NAG01 clades A, B and C from specific liver samples we used the Circos tool. Fig. 2 was created using Inkscape (<http://www.inkscape.org>), Circos (61), and the Interactive Tree of Life (iTOL) (<http://embl.itol.de>) (62). Connections highlighting phylogenetically disparate clones from the same frog were plotted and colour-coded by geographic sampling location (see key). This plot was then combined with an inverted circular tree generated by iTOL using Inkscape.

#### Histology of representative tadpole tissue samples

- Wake DB & Vredenburg VT (2008) Are we in the midst of the sixth mass extinction? A view from the world of amphibians. *Proc. Natl. Acad. Sci. USA* 105(Supplement 1):11466-11473.
- Monastersky R (2014) Life - a status report. *Nature* 516:159-161.
- Rohr JR & Raffel TR (2010) Linking global climate and temperature variability to widespread amphibian declines putatively caused by disease. *Proc. Natl. Acad. Sci. USA* 107(18):8269-8274.
- Johnson PTJ, et al. (2007) Aquatic eutrophication promotes pathogenic infection in amphibians. *Proc. Natl. Acad. Sci. USA* 104(40):15781-15786.
- Collins JP & Storer A (2003) Global amphibian declines: sorting the hypotheses. *Diversity and Distributions* 9(2):89-98.
- Pounds AJ, et al. (2006) Widespread amphibian extinctions from epidemic disease driven by global warming. *Nature* 439(7073):161-167.
- Crawford AJ, Lips KR, & Bermingham E (2010) Epidemic disease decimates amphibian abundance, species diversity, and evolutionary history in the highlands of central Panama. *Proc. Natl. Acad. Sci. USA* 107(31):13777-13782.
- Lips KR, et al. (2006) Emerging infectious disease and the loss of biodiversity in a Neotropical amphibian community. *Proc. Natl. Acad. Sci. USA* 103(9):3165-3170.
- Fisher MC, Garner TW, & Walker SF (2009) Global emergence of *Batrachochytrium dendrobatidis* and amphibian chytridiomycosis in space, time, and host. *Annu. Rev. Microbiol.* 63:291-310.
- Daszak P, et al. (1999) Emerging infectious diseases and amphibian population declines. *Emerging Infectious Disease* 5(6):736-748.
- Duffus ALJ, Pauli BD, Wozney K, Brunetti CR, & Berrill M (2008) Frog virus 3-like infections in aquatic amphibian communities. *Journal of Wildlife Diseases* 44(1):109-120.
- Davis AK, Yabsley MJ, Keel MK, & Maerz JC (2007) Discovery of a novel alveolate pathogen affecting Southern Leopard frogs in Georgia: description of the disease and host effects. *EcoHealth* 4:310-317.
- Green DE, Feldman SH, & Wimsatt J (2003) Emergence of a *Perkinsus*-like agent in anuran liver during die-offs of local populations: PCR detection and phylogenetic characterization. *Proceedings of the American Association of Zoo Veterinarians* 2003:120-121.
- Azevedo C (1989) Fine structure of *Perkinsus atlanticus* n. sp. (Apicomplexa, Perkinsae), a parasite of the clam *Ruditapes decussatus* from Portugal. *Int. J. Parasitol.* 19:627-635.
- Lester RJG & Davis GHG (1981) A new *Perkinsus* species (Apicomplexa, Perkinsae) from the abalone, *Haliotis ruber*. *J. Invertebr. Pathol.* 37:181-187.
- Bachvaroff TR, et al. (2014) Dinoflagellate phylogeny revisited: Using ribosomal proteins to resolve deep branching dinoflagellate clades. *Mol. Phylogenet. Evol.* 70(0):314-322.
- Bruggerolle G (2003) Apicomplexan parasite *Cryptophagus* renamed *Rastrimonas* gen. nov. *Eur. J. Protistol.* 39:101.
- Bruggerolle G (2002) *Cryptophagus subtilis*: a new parasite of cryptophytes affiliated with the Perkinsozoa lineage. *Eur. J. Protistol.* 37:379-390.
- Mackin JG (1951) Histopathology of infection of *Crassostrea virginica* (Gmelin) by *Dermocystidium marinum* Mackin, Owen & Collier. *Bull. Mar. Sci.* 1(1):72-87.
- Park MG, Yih W, & Coats DW (2004) Parasites and phytoplankton, with special emphasis on dinoflagellate infections. *J. Euk. Microbiol.* 51(2):145-155.
- Lefranc M, Thénot A, Lepère C, & Debroas D (2005) Genetic diversity of small eukaryotes in lakes differing by their trophic status. *Appl. Environ. Microbiol.* 71(10):5935-5942.
- Lepère C, Domaizon I, & Debroas D (2008) Unexpected importance of potential parasites in the composition of freshwater small-eukaryote community. *Appl. Environ. Microbiol.* 74:2940-2949.
- Mangot JF, Lepère C, Bouvier C, Debroas D, & Domaizon I (2009) Community structure and dynamics of small eukaryotes targeted by new oligonucleotide probes: new insight into the lacustrine microbial food web. *Appl. Environ. Microbiol.* 75(19):6373-6381.
- Brate J, et al. (2010) Freshwater Perkinsae and marine-freshwater colonizations revealed by pyrosequencing and phylogeny of environmental rDNA. *ISME J.* 4(9):1144-1153.
- Chambouvet A, et al. (2014) Diverse molecular signatures for ribosomally 'active' Perkinsae in marine sediments. *BMC Microbiol.* 14(1):110.
- Fredericks DN & Relman DA (1996) Sequence-based identification of microbial pathogens: a reconsideration of Koch's postulates. *Clin. Microbiol. Rev.* 9(1):18-33.
- Relman DA, Schmidt TM, MacDermott RP, & Falkow S (1992) Identification of the uncultured *Bacillus* of Whipple's Disease. *N. Engl. J. Med.* 327(5):293-301.
- Anderson BE, Dawson JE, Jones DC, & Wilson KH (1991) *Ehrlichia chaffeensis*, a new species associated with human ehrlichiosis. *J. Clin. Microbiol.* 29(12):2838-2842.
- Wuyts J, et al. (2000) Comparative analysis of more than 3000 sequences reveals the existence of two pseudoknots in area V4 of eukaryotic small subunit ribosomal RNA. *Nucleic Acids Res.* 28(23):4698-4708.
- Richards TA, Vepritskiy AA, Gouliamova D, & Nierzwicki-Bauer SA (2005) The molecular diversity of freshwater picoeukaryotes from an oligotrophic lake reveals diverse, distinctive and globally dispersed lineages. *Environ. Microbiol.* 7(9):1413-1425.
- Lefèvre E, et al. (2007) Unveiling fungal zooflagellates as members of freshwater picoeukaryotes: evidence from a molecular diversity study in a deep meromictic lake. *Environ. Microbiol.* 9(1):61-71.
- Lefèvre E, Roussel B, Amblard C, & Sime-Ngando T (2008) The molecular diversity of

For eight samples, half of the liver that was not used for DNA extraction was stored in 100% ethanol at 4 °C. Each liver was then embedded in paraffin wax (Sigma-Aldrich, St. Louis, USA), cut into serial sections 5-um-thick using a Shadon tissue processor (Thermo Electron Corporation, Runcorn, UK). Sections were collected onto glass slides, stained with hematoxylin and eosin staining methods (Thermo Fisher, Pittsburgh, USA). The sections of each slide were mounted using Histomount (National Diagnostics, Hull, UK). Sections were examined by light microscopy (Microscope Olympus IX73) for the presence of putative parasites and digital images were obtained using the Infinity 3 camera (Lumenera, Corporation, Ottawa, Canada).

freshwater picoeukaryotes reveals high occurrence of putative parasitoids in the plankton. *PLoS ONE* 3(6):e2324.

- Villalba A, Reece KS, Ordás MC, Casas SM, & Figueras A (2004) Perkinsosis in molluscs: A review. *Aquat. Living Resour.* 17:411-432.
- Stern RF, et al. (2012) Evaluating the ribosomal internal transcribed spacer (ITS) as a candidate dinoflagellate barcode marker. *PLoS ONE* 7(8):e42780.
- Galluzzi L, et al. (2004) Development of a real-time PCR assay for rapid detection and quantification of *Alexandrium minutum* (a dinoflagellate). *Appl. Environ. Microbiol.* 70:1199-1206.
- McCutchan T, et al. (1988) Primary sequences of two small subunit ribosomal RNA genes from *Plasmodium falciparum*. *Mol. Biochem. Parasitol.* 28:63-68.
- Nishimoto Y, et al. (2008) Evolution and phylogeny of the heterogeneous cytosolic SSU rRNA genes in the genus *Plasmodium*. *Mol. Phylogenet. Evol.* 47(1):45-53.
- Jones MEB, Armien AG, Rothermel BB, & Pessier AP (2012) Granulomatous myositis associated with a novel alveolate pathogen in an adult southern leopard frog. *Dis. Aquat. Organ.* 102:163-167.
- Cook J (2008) Transmission and occurrence of *Dermomycoides* sp. in *Rana sevosia* and other Ranids in the North Central Gulf of Mexico States. (University of Southern Mississippi).
- Palumbi SR & Crandall KA (1991) *The simple fool's guide to PCR. A collection of PCR protocols, version 2.* (University of Hawaii).
- Vences M, Thomas M, van der Meijden A, Chiari Y, & Vieites D (2005) Comparative performance of the 16S rRNA gene in DNA barcoding of amphibians. *Front. Zool.* 2(1):5.
- Gahl MK (2007) Spatial and temporal patterns of amphibian disease in acadiana national park wetlands. PhD (The University of Maine, Maine, USA).
- Landsberg JH, et al. (2013) Co-infection by alveolate parasites and frog virus 3-like ranavirus during an amphibian larval mortality event in Florida, USA. *Dis. Aquat. Organ.* 105(2):89-99.
- Rollins-Smith LA (1998) Metamorphosis and the amphibian immune system. *Immunol. Rev.* 166(1):221-230.
- Casas SM, Peyre JFL, Reece KS, Azevedo C, & Villalba A (2002) Continuous in vitro culture of the carpet shell clam *Tapes decussatus* protozoan parasite *Perkinsus atlanticus*. *Dis. Aquat. Organ.* 52(3):217-231.
- Chu FLE & La Peyre JF (1993) *Perkinsus marinus* susceptibility and defense-related activities in Eastern oysters *Crassostrea virginica* - temperature effects. *Dis. Aquat. Organ.* 16:223-234.
- Cook T, Folli M, Klinck J, Ford S, & Miller J (1998) The relationship between increasing sea surface temperature and the northward spread of *Perkinsus marinus* (Dermo) disease epizootics in oysters. *Estuar. Coast. Shelf. Sci.* 40:587-597.
- Andrews JD (1988) Epizootiology of the disease caused by the oyster pathogen *Perkinsus marinus* and its effects on the oyster industry. *Am. Fish. Soc. Spec. Publ.* 18:47-63.
- Garcés E, Alacid E, Bravo I, Fraga S, & Figueras R (2013) *Parvilicifera sinerae* (Alveolata, Myxozoa) is a generalist parasitoid of dinoflagellates. *Protist* 164(2):245-260.
- Young BE, et al. (2001) Population declines and priorities for amphibian conservation in Latin America. *Conserv. Biol.* 15:1213-1223.
- Buckley J, Beebe TJ, & Schmidt BR (2014) Monitoring amphibian declines: population trends of an endangered species over 20 years in Britain. *Anim. Conserv.* 17:27-34.
- Duffus AJ (2009) Chytrid blinders: what other disease risks to amphibians are we missing? *EcoHealth* 6(3):335-339.
- Ludwig W, et al. (2004) ARB: a software environment for sequence data. *Nucleic Acids Res.* 32(4):1363-1371.
- Pruesse E, et al. (2007) SILVA: a comprehensive online resource for quality checked and aligned ribosomal RNA sequence data compatible with ARB. *Nucleic Acids Res.* 35(21):7188-7196.
- Edgar RC (2004) MUSCLE: a multiple sequence alignment method with reduced time and space complexity. *BMC Bioinformatics* 5:113.
- Galtier N, Gouy M, & Gautier C (1996) SEAVIEW and PHYLO-WIN: two graphic tools for sequence alignment and molecular phylogeny. *Comput. Appl. Biosci.* 12(6):543-548.
- Frost DR, et al. (2006) The amphibian tree of life. *Bulletin of the American Museum of Natural History*:1-291.
- Ronquist F & Huelsenbeck JP (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19(12):1572-1574.
- Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30(9):1312-1313.
- Lockhart PJ, Steel MA, Hendy MD, & Penny D (1994) Recovering evolutionary trees under a more realistic model of sequence evolution. *Mol. Biol. Evol.* 11:605-612.
- Krzywinski M, et al. (2009) Circos: An information aesthetic for comparative genomics. *Genome Res.* 19:1639-1645.
- Letunic I & Bork P (2006) Interactive Tree Of Life (iTOL): an online tool for phylogenetic tree display and annotation. *Bioinformatics* 23(1):127-128.

#### Acknowledgements.

This research was funded primarily by the UK Department for Environment, Food and Rural Affairs (DEFRA) through the SYNTAX scheme run by the Linnean Society of London and the Systematics Association



UK awarded to TAR, and by fellowship awards to AC. AC is supported by a Marie Curie Intra-European Fellowship grant (FP7-PEOPLE-2011-IEF - 299815 PARAFROGS) and an EMBO Long-Term fellowship (ATL-1069-2011). TAR is an EMBO Young Investigator and is supported by research grants from the Gordon and Betty Moore Foundation (Grant GBMF3307), NERC, Leverhume Trust, BBSRC, CIFAR and the Royal Society. For permits and approvals for fieldwork in French Guiana, GB, DJG and MW thank Myrian Virevaire (Direction de l'Environnement de l'Aménagement et du Logement) and Le Comité Scientifique Régional du Patrimoine Naturel. Additionally, for assisting our fieldwork we thank Elodie Courtoise, Antoine Fouquet, Philippe Gaucher, Fausto Starace and family, and Jeannot and Odette (Camp Patawa). Cameroon fieldwork was permitted by the Ministry of Forestry & Wildlife (# 5032) funded by the USA Fish & Wildlife's Wildlife Without

This research was funded primarily by the UK Department for Environment, Food and Rural Affairs (DEFRA) through the SYNTAX scheme run by the Linnean Society of London and the Systematics Association UK awarded to TAR, and by fellowship awards to AC. AC is supported by a Marie Curie Intra-European Fellowship grant (FP7-PEOPLE-2011-IEF - 299815 PARAFROGS) and an EMBO Long-Term fellowship (ATL-1069-2011). TAR is an EMBO Young Investigator and is supported by research grants from the Gordon and Betty Moore Foundation (Grant GBMF3307), NERC, Leverhume Trust, BBSRC, CIFAR and the Royal Society. For permits and approvals for fieldwork in French Guiana, GB, DJG and MW thank Myrian Virevaire (Direction de l'Environnement de l'Aménagement et du Logement) and Le Comité Scientifique Régional du Patrimoine Naturel. Additionally, for assisting our fieldwork we thank Elodie Courtoise, Antoine Fouquet, Philippe Gaucher, Fausto Starace and family, and Jeannot and Odette (Camp Patawa). Cameroon fieldwork was permitted by the Ministry of Forestry & Wildlife (# 5032) funded by the USA Fish & Wildlife's Wildlife

Borders – Amphibians in Decline scheme, Royal Geographical Society, and the Zoological Society of London's EDGE Fellowship scheme. DJG, TMD-B and MW thank Marcel Kouete, Solo Ndeme, Gespo Alobwede and many local people, especially in Mamfe Division. For access to Tanzanian material we thank Michele Menegon and Simon Loader. FM is a joint Natural History Museum, London and University College London PhD student. MJ was supported by the Czech Science Foundation (P506/10/2330) and the Institute of Parasitology, BC AS CR (institutional support RVO: 60077344). We thank Hendrik Müller for tadpole expertise, and David Blackburn for assistance in identifying *Xenopus* life cycle stages. We thank Caroline Ware of the NHM Wildlife Garden for assistance with sampling and Anke Lange of Biosciences at the University of Exeter for assistance in histological analysis.

Without Borders – Amphibians in Decline scheme, Royal Geographical Society, and the Zoological Society of London's EDGE Fellowship scheme. DJG, TMD-B and MW thank Marcel Kouete, Solo Ndeme, Gespo Alobwede and many local people, especially in Mamfe Division. For access to Tanzanian material we thank Michele Menegon and Simon Loader. FM is a joint Natural History Museum, London and University College London PhD student. MJ was supported by the Czech Science Foundation (P506/10/2330) and the Institute of Parasitology, BC AS CR (institutional support RVO: 60077344). We thank Hendrik Müller for tadpole expertise, and David Blackburn for assistance in identifying *Xenopus* life cycle stages. We thank Caroline Ware of the NHM Wildlife Garden for assistance with sampling and Anke Lange of Biosciences at the University of Exeter for assistance in histological analysis.

Submission PDF