Title: Multiple facets of biodiversity drive the diversity-stability relationship

Authors

Dylan Craven [1,2,3], Nico Eisenhauer [1,2] , William D. Pearse [5], , Yann Hautier [6], Christiane Roscher [1,7], Forest Isbell [8], John Connolly, Anne Ebeling, John Griffin, Jessica Hines, Anke Jentsch, Nathan Lemoine, Sebastian T. Meyer, Jasper van Ruijven, Melinda Smith, Carl Beierkuhnlein, Gerhard Boenisch, Andy Hector, Jens Kattge, Jürgen Kreyling, Vojtech Lanta, Enrica de Luca, H. Wayne Polley, Peter B. Reich , David Tilman, Alexandra Weigelt, Brian Wilsey, Pete Manning [4]

Author Affiliations

1 German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher Platz 5e , 04103 Leipzig, Germany

2 Institute of Biology, Leipzig University, Johannisallee 21, 04103 Leipzig, Germany

3 Department of Community Ecology, Helmholtz Centre for Environmental Research – UFZ, Theodor-Lieser Straße 4, 06120, Halle (Saale), Germany

4 Senckenberg Gesellschaft für Naturforschung, Biodiversity and Climate Research Centre, 60325 Frankfurt, Germany

5 Department of Biology, Utah State University, Logan,UT 84322, USA

6 Ecology and Biodiversity Group, Department of Biology, Utrecht University, Padualaan 8, Utrecht, CH 3584, The Netherlands

7 Department of Physiological Diversity, Helmholtz Centre for Environmental Research – UFZ, Leipzig, Germany

8 Department of Ecology, Evolution and Behavior, University of Minnesota, Saint Paul, MN, USA

E-mail addresses
dylan.craven@idiv.de, nico.eisenhauer@idiv.de, peter.manning@senckenberg.de, will.pearse@usu.edu, Y.Hautier@uu.nl, christiane.roscher@ufz.de, isbell@umn.edu

Running title: Drivers of biodiversity-stability relationships

Keywords: fast-slow, functional diversity, phylogenetic diversity, species richness, invariability, stability, asynchrony, biodiversity

Article type: Letters

Number of words
Abstract: 150
Main text (excluding abstract, acknowledgements, references, tables, and figures): 4,576
Number of references: 99

Number of figures: 3

Number of tables: 1
Number of text boxes: 0

Corresponding author information:

Dylan Craven

Mailing address:
German Centre for Integrative Biodiversity Research (iDiv)
Deutscher Platz 5e
04103 Leipzig
Germany

Telephone number: +49 341 9733117
Fax number:
E-mail: dylan.craven@aya.yale.edu

Statement of authorship: XX authors contributed functional trait data. YY contributed data from grassland biodiversity experiments. ZZ developed the initial concept at a workshop. WDP built the phylogeny and calculated phylogenetic diversity indices. DC compiled data and analyzed data with input from PM, NE, WDP, and YH. DC and PM wrote the first draft of the manuscript and all co-authors contributed substantially to revisions.

Data accessibility: The authors confirm that data supporting the results of this manuscript will be archived in Dryad and that the corresponding DOI will be included at the end of the article.

Abstract

A significant body of evidence has demonstrated that biodiversity stabilizes ecosystem functioning in grassland ecosystems. However, the relative importance of the biological drivers underlying these relationships remains unclear. Here we utilized data from 39 biodiversity experiments and a structural equation modeling approach to investigate the roles of phylogenetic diversity, functional diversity and community-level averages of 'fast-slow' traits, species richness, and asynchrony in driving the diversity-stability relationship. The structural equation model explained 78% of variation in asynchrony and 58% in ecosystem stability and showed that high species richness and phylogenetic diversity stabilized biomass production via asynchrony and, surprisingly, that low phylogenetic diversity enhanced ecosystem stability directly. The effects of functional diversity and fast-slow traits on ecosystem stability were weak and highly variable across sites, respectively. These results demonstrate that biodiversity influences ecosystem stability via multiple pathways, thus suggesting a more complex role of biodiversity in mediating ecosystem stability than previously recognized.

Introduction

The relationship between the diversity of ecosystems and their stability has long been a fundamental subject of fundamental ecological research (May 1973; McNaughton 1978; Ives & Carpenter 2007), and this research topic has gained new impetus due to global environmental change and biodiversity loss, both of which threaten the stability of ecosystem functions and the ecosystem services they underpin (Balvanera *et al.* 2006; Hautier *et al.* 2015; Isbell *et al.* 2015b; Donohue *et al.* 2016). A substantial body of theoretical and empirical work on this question has focused on the relationship between plant species diversity and biomass production, and this has consistently demonstrated that the productivity of species-rich systems shows less variation over time (e.g., Jiang & Pu 2009; Hector *et al.* 2010; Campbell *et al.* 2011; de Mazancourt *et al.* 2013; Gross *et al.* 2014).

Stability (or invariability) of ecosystem functioning is an integrative measure of short- and long-term responses of populations and communities to environmental variation (Oliver *et al.* 2015). As a result, higher stability of species-rich ecosystems can be attributed to their higher resistance (*i.e.*, biomass shows little deviation from normal during perturbations) and/or resilience (*i.e.*, biomass returns to normal rapidly after perturbations; Tilman & Downing 1994; Van Ruijven & Berendse 2010; Isbell *et al.* 2015a). To date, a considerable number of mechanisms – not necessarily mutually exclusive - have been proposed as the cause of these patterns, primarily asynchrony in species responses to environmental variation, insurance effects, overyielding, and statistical averaging (Doak *et al.* 1998; Yachi & Loreau 1999; Lehman & Tilman 2000; Loreau & Mazancourt 2008; de Mazancourt *et al.* 2013) and tested empirically (*e.g.*, Isbell *et al.* 2009; Hector *et al.* 2010; Roscher *et al.* 2011; de Mazancourt *et al.* 2013; Gross *et al.* 2014; Hautier *et al.* 2014). Here, we explore a suite of potential biodiversity-dependent mechanisms that influence stability. We hypothesize that the relationship between biodiversity and ecosystem stability is mediated by a range of biological

drivers and that we can identify the signal of these mechanisms using community-level properties: functional composition and functional and phylogenetic diversity.

Functional composition may play a key role in stabilizing biomass production in grasslands because growth-related traits strongly influence the production and persistence of plant biomass (Díaz & Cabido 2001). While plants differ greatly in their trait values and strategies, a large proportion of global plant trait variation is correlated along a single axis which distinguishes between 'fast' or exploitative species that are capable of rapid resource uptake, growth, and tissue turnover and 'slow' or conservative species with slower rates of growth, resource uptake, and tissue turnover (Wright *et al.* 2004; Reich 2014). The former typically possess high specific leaf area (SLA), low leaf dry matter content (LDMC), and high leaf nitrogen (N), the latter the opposite (Grime 1977; Reich 2014; Díaz *et al.* 2016). There is growing evidence that variation in functional composition along the fast-slow spectrum has cascading effects on ecosystem stability. For example, high LDMC values have been found to increase ecosystem stability in experimental and semi-natural grassland communities (Polley *et al.* 2013; Májeková *et al.* 2014). We therefore expect that communities dominated by slow species will be more stable than those dominated by fast species. The net effect of functional composition on ecosystem stability, however, may be low because the opposing effects of fast communities, which should be more resilient, and slow communities, which should be more resistant, may cancel each other out.

Variation in plant ecological strategies, quantified as trait diversity, may explain ecosystem stability because higher trait variability should increase temporal niche complementarity, thus reducing variation of productivity over time ("functional insurance" ; Díaz & Cabido 2001). That is, functionally diverse communities maintain biomass production over time because they contain an array of fast and slow species that vary in their ability to compete for and utilise growth-limiting

187 resources, such as water, nutrients, and light (Flynn *et al.* 2011; Roscher *et al.* 2012; Spasojevic &
188 Suding 2012; Fischer *et al.* 2016). As fast species are likely to recover rapidly following
189 disturbance (resilience) and slow species will be able to tolerate environmental stresses and
190 perturbations (resistance ; Grime 1977; Reich 2014), we hypothesize that functionally diverse
191 communities will exhibit both resistance and resilience, thus increasing ecosystem stability.
192
193 The third class of biological drivers that we propose as underlying the diversity-stability
194 relationship are those associated with phylogenetic diversity. Broadly speaking, phylogenetic
195 diversity can be seen as representing the diversity of phylogenetically conserved functional traits.
196 Traits which reflect a shared co-evolutionary history of biotic interactions often show a high degree
197 of phylogenetic conservatism (Gomez *et al.* 2010), such as mycorrhizal tendency and N fixation
198 (Flynn *et al.* 2011; Reinhart *et al.* 2012). Phylogenetically similar species are also known to share
199 pathogens or immune responses via their shared co-evolutionary history (Gilbert *et al.* 2012; Parker
200 *et al.* 2015). Importantly, there is strong evidence showing that phylogenetic diversity has a
201 consistently positive effect on ecosystem stability in grasslands (Flynn *et al.* 2011; Cadotte *et al.*
202 2012; Cadotte 2015). We therefore hypothesize that greater phylogenetic diversity will stabilize
203 biomass production over time by increasing (measured and unmeasured) trait diversity and diluting
204 the effects of pathogen outbreaks and herbivore attacks.
205
206 Plant species richness may affect ecosystem stability via multiple pathways that are independent of
207 functional and phylogenetic diversity. There is evidence for a range of trait-based mechanisms not
208 related to the fast-slow spectrum via which diversity may confer ecosystem stability, such as
209 persistent seedbanks (Pérez-Harguindeguy *et al.* 2013), aerenchyma production that maintains gas
210 exchange (Wright *et al.* 2016), and regrowth from belowground storage organs (Hoover *et al.* 2014)
211 or carbohydrate reserves (McDowell *et al.* 2008), that confer resilience. There are also properties

212 that enhance resistance, such as drought tolerance (Bartlett *et al.* 2012; Craine *et al.* 2013).
213 Furthermore, diversity may also confer ecosystem stability through traits that enable community-
214 level production in the face of environmental variability, such as variation in rooting depth (Weigelt
215 *et al.* 2008) and phenology (Fargione & Tilman 2005). Plant species richness can also directly
216 affect ecosystem stability by modifying environmental conditions. For example, the higher
217 productivity of species-rich communities is associated with more rapid rates of soil organic matter
218 accumulation (Fornara & Tilman 2008; Cong *et al.* 2014; Lange *et al.* 2015) and soil aggregate
219 formation (Gould *et al.* 2016), which result in a more aerobic, mesic soil environment in which
220 plant growth is more constant. We expect that these pathways will be represented statistically by
221 residual effects of species richness on ecosystem stability (Fig. S1).
222
223 Multiple biological drivers (described above) may simultaneously affect ecosystem stability by
224 increasing species asynchrony. Asynchrony, which describes the extent to which species-level
225 productivity is correlated within a community over time, has been identified in a growing number of
226 theoretical and empirical studies as a key mechanism underlying diversity-stability relationships
227 (e.g., Yachi & Loreau 1999; de Mazancourt *et al.* 2013; Gross *et al.* 2014; Hautier *et al.* 2014).
228 Asynchrony, where decreases in the productivity of some species are compensated by increases in
229 the productivity of other species, causes ecosystem stability to increase due to interspecific
230 interactions (e.g., Lehman & Tilman 2000; Gross *et al.* 2014), negative frequency dependence, e.g.
231 due to pathogen outbreaks (Maron *et al.* 2011; Schnitzer *et al.* 2011), and/or the greater likelihood
232 that diverse communities contain a wider range of species' responses to environmental conditions
233 (de Mazancourt *et al.* 2013; Tredennick *et al.* 2017). With the notable exception of species richness,
234 biodiversity-dependent mechanisms have rarely been used to explain the effects of asynchrony (but
235 see Roscher *et al.* 2011). We hypothesize that multiple facets of biodiversity species richness and

functional and phylogenetic diversity will enhance asynchrony by increasing variation in traits that confer resistance and resilience (Mori *et al.* 2013; Aubin *et al.* 2016).

While there is empirical evidence that each of the aforementioned biological drivers contribute to the overall relationship between diversity and stability, the relative importance of these drivers has not been investigated across a range of grassland ecosystems. Here we assessed for the first time the simultaneous contribution of multiple aspects of biodiversity in driving biodiversity-stability relationships by performing a meta-level analysis using data from 39 grassland biodiversity-ecosystem function experiments distributed across North America and Europe. The biological drivers were decoupled using structural equation models, which represented the interrelations described above (Fig. S1). We hypothesized that: *i*) greater plant species richness, functional diversity, and phylogenetic diversity will increase ecosystem stability by increasing asynchrony and *ii*) species-rich communities and those dominated by slow species will increase ecosystem stability directly. We show that high species richness and phylogenetic diversity stabilize biomass production via asynchrony and, surprisingly, that low phylogenetic diversity increases ecosystem stability directly.

Methods

Data preparation

We created a database by combining data from biodiversity experiments that manipulated plant species richness in grasslands and measured community- and species-level aboveground plant biomass for at least three years using 39 studies across North America and Europe from Isbell *et al.* (2015a) and Craven *et al.* (2016) (Table S1). In total, our dataset comprises observations from 1,692 plots and 165 plant species, which were standardized using the Taxonomic Name Resolution Service (http://trns.iplantcollaborative.org ; Boyle *et al.* 2013).

For each plot within these experiments, we quantified ecosystem stability (or ecosystem invariability) as the inverse of the coefficient of variation (μ/σ ; e.g., Isbell *et al.* 2015a), which is the ratio of the mean to the standard deviation of aboveground plant biomass over time. Asynchrony (η) was quantified following Gross *et al.* (2014) as the average correlation across species between the biomass of each species and the total biomass of all other species in a plot:

$$\eta = (1/n) \Sigma_i \text{ corr } (Y_i, \Sigma\, j \neq i\ Y_j), \quad \text{(Eq.1)}$$

where Y_i is the biomass of species *i* in a plot containing *n* species. This measure of asynchrony ranges from -1, where species' aboveground plant biomass is maximally asynchronous, to 1, where species' aboveground plant biomass is maximally synchronized. Further, η is independent of the number of species and individual species' variances (Gross *et al.* 2014).

We selected four 'fast-slow' leaf traits associated with the leaf economic spectrum (Wright *et al.* 2004; Díaz *et al.* 2016)(Wright *et al.* 2004; Díaz *et al.* 2016), specific leaf area (SLA; $mm^2\ mg^{-1}$), leaf dry matter content (LDMC; $g\ g^{-1}$) , foliar N (%), and foliar P (%) and obtained data from the TRY database (Kattge *et al.* 2011) and additional studies that measured traits in our data set (Grime *et al.* 2007; Wacker *et al.* 2009; Roscher *et al.* 2012; Daneshgar *et al.* 2013; Jane A. Catford, unpublished data). Trait values were converted to the same units and outliers were excluded (standard deviation > 4). Values were then averaged by contributor and then by species. Genus-level means were used when species-level data were not available; species-level data for SLA, LDMC, leaf N, and leaf P were available for 98%, 83 %, 92 %, and 62 % of species, respectively. Combining species- and genus-level values, our final trait data set included SLA, LDMC, and foliar N values for more than 96% of the species and leaf P values for 93% of the species.

Functional diversity and composition

We calculated functional diversity in ‘fast-slow’ traits as either functional dispersion (FD; abundance weighted) or functional richness (FRic; not weighted by abundance) to represent complementarity among co-occurring species and volume of trait space, respectively, using the ‘FD’ package (Laliberté & Legendre 2010). Results for both measures of functional diversity were qualitatively similar. Therefore, we present results for functional dispersion in the main text and for functional richness in Supplementary Materials.

We used the first axis of a principal component analysis (PCA) of community-weighted means of SLA, LDMC, leaf N, and leaf P to represent the fast-slow spectrum, where ‘slow’ communities have high community-level averages in trait values that are correlated with slow rates of resource acquisition and growth, and ‘fast’ communities have high community-level averages in trait values that correspond to high rates of resource acquisition and growth (Reich 2014; Salguero-Gómez *et al.* 2016). The first PCA captured 60.4% of variation among the four traits (Figure S2) and represents the ‘fast-slow’ spectrum of communities, from those dominated by ‘slow’ species with low SLA and leaf N and P and high LDMC to those dominated by ‘fast’ species with high SLA and leaf N and P and low LDMC. Trait measures were calculated annually for each plot and then averaged across years.

Phylogenetic diversity

We used the molecular phylogeny from Zanne *et al.* (2014) as a backbone to build a phylogeny of all species within the experiments, conservatively binding species into the backbone using dating information from congeners in the tree (using *congeneric.merge* ; Pearse *et al.* 2015). We then calculated abundance-weighted phylogenetic diversity as mean nearest taxon distance (eMNTD ; Webb *et al.* 2002; Pearse *et al.* 2014). eMNTD captures competitive differences among species in previous studies (Godoy *et al.* 2014) and the sharing of specialized pathogens tends to be confined

311 to closely related species (Gilbert *et al.* 2012; Parker *et al.* 2015). eMNTD, therefore, is a good
312 metric to test our hypotheses about the mechanisms that explain variation in species asynchrony and
313 ecosystem stability.
314
315 *Climate*
316 As previous studies have reported strong impacts of inter-annual variation in weather conditions on
317 plant productivity over time (Huxman *et al.* 2004; Sala *et al.* 2012), we included site-level climate
318 data in order to explain across-site variation in ecosystem stability. To describe environmental
319 conditions during each study in a consistent manner across sites, we calculated mean annual
320 precipitation (MAP), mean annual temperature (MAT), inter-annual variation in precipitation
321 (coefficient of variation of MAP), and inter-annual variation in temperature (coefficient of variation
322 of MAT) using data from CRU TS 3.2.3 (Harris *et al.* 2014). We tested for the individual effects of
323 each climatic variable on mean temporal stability using a linear regression model and found that
324 inter-annual variation in precipitation best explained variation in mean temporal stability (ΔAICc =
325 3.68). This variable was therefore selected for use in later analyses.
326
327 *Data analysis*
328 To explore the bi-variate relationships between each of our hypothesized drivers and ecosystem
329 stability, we first fitted separate linear mixed-effects models that tested for the effects of plant
330 species richness, asynchrony, phylogenetic diversity, functional diversity, and the fast-slow
331 spectrum on ecosystem stability. Multiple random effect structures were tested for each model;
332 random effects were included for a study factor and interactions of study with plant species richness
333 and other predictor variables. Random effect structures allowed the intercepts and slopes to vary
334 among studies if their retention was supported by model selection. We used AICc, a second-order
335 bias correction to Akaike's information criterion for small sample sizes, to select the most

parsimonious model (Burnham & Anderson 2003). Models were fit using the 'nlme' package and model assumptions were checked by visually inspecting residual plots for homogeneity and quantile-quantile plots for normality. Intra-class correlation (ICC) was calculated to compare the variability within a study to variability across studies.

To estimate the direction and strength of relationships between plant species richness, functional and phylogenetic diversity, the fast-slow spectrum, and asynchrony, Pearson's correlation coefficients and sampling variance were calculated for each study. Using the 'metafor' package, mean effect sizes of the relationships among the aforementioned variables were estimated using random effects models and restricted maximum likelihood (Viechtbauer 2010). The Knapp-Hartung adjustment was used to account for the uncertainty in the estimation of residual heterogeneity (Knapp & Hartung 2003).

To test the relative importance of the fast-slow spectrum, functional and phylogenetic diversity, climate, and asynchrony in driving temporal stability, we fit piecewise structural equation models (SEM ; Lefcheck 2016) using 'piecewiseSEM'. Testing the aforementioned effects on resistance and resilience (as in Isbell *et al.* 2015a) was not possible because of the unequal distribution of extreme climate events across sites, which prevented fitting a general SEM. We formulated a hypothetical causal model (Fig. S1) based on a priori knowledge of the system, which we used to test the fit of the model to the data. This model mirrors the relationships. We also included direct paths from species richness, functional and phylogenetic diversity to ecosystem stability to represent other potential biodiversity-dependent mechanisms that influence ecosystem stability. Finally, we included a direct path from inter-annual variation in precipitation to ecosystem stability. We included direct paths from species richness to functional and phylogenetic diversity and the fast-slow spectrum because variation in these variables can be directly attributed to the

experimental manipulation of species richness in all studies (Flynn *et al.* 2011). All initial models contained correlated errors between functional diversity, phylogenetic diversity, and the fast-slow spectrum. Paths were added to the initial model if they significantly improved model fit using modification indices ($P < 0.05$). This resulted in the addition of a direct path between phylogenetic diversity and ecosystem stability in the final model. Model fit was assessed using Fisher's C statistic ($P > 0.10$). SEMs were fitted using linear mixed-effects models where study was treated as a random group factor and species richness as a random slope. In all analyses, plant species richness and ecosystem stability were $\log_2$ transformed. Model assumptions of normality and homogeneity of variance were inspected visually and collinearity was assessed by estimating variance inflation (VIF < 2 ; Zuur *et al.* 2010). All analyses were performed using R 3.3.1 (R Core Team 2016).

Results

Our analysis confirms that species richness, phylogenetic and functional diversity, and asynchrony each demonstrated a significant and positive relationship with ecosystem stability that was generally consistent across experiments (Figs. 1 and 2). Individual fixed effects of these drivers explained low amounts of variation in ecosystem stability (Table 1, marginal R^2), with a larger proportion being explained by the random effects (Tables 1 and S3, conditional R^2). In contrast, there was not a consistent effect of the fast-slow spectrum on ecosystem stability ($P > 0.10$; Fig. 2c). While there was evidence that communities dominated by 'slow' species stabilized productivity to a greater extent than those dominated by 'fast' species at certain experimental sites (Fig. 2c) and vice-versa, the high intra-class correlation for this model (Tables 1 and S3) indicates that the CWM fast-slow effect was highly variable across all experimental sites.

Across experimental sites, all measures of diversity were significantly positively correlated with one another ($r = 0.67 – 0.94$; Fig. S2, Table S2). CWM of the fast-slow spectrum also varied significantly in response to species richness and functional richness. With increasing species and functional richness, communities became increasingly dominated by 'fast' species (Fig. S2, Table S2). With increasing species richness, phylogenetic diversity and functional diversity, and asynchrony increased significantly (Fig. S2, Table S2).

Our structural equation model provided clear evidence that asynchrony is a key mechanism mediating the biodiversity-stability relationships and that it is is driven by multiple aspects of biodiversity (Fig. 3). Overall, the data fit our model well (Fisher's $C = 11.84$, $df = 12$, $P = 0.51$; AICc = 72.96, K = 30, n = 1,692). Fixed effects explained 19% of variation in ecosystem stability (marginal R^2), which increased to 58% (conditional R^2) when accounting for fixed and random effects. In total, species richness, phylogenetic and functional diversity, and the fast-slow spectrum explained 53% of variation in asynchrony (marginal R^2), which increased to 78% when random effects were accounted for (conditional R^2).

The strongest pathway of influence on ecosystem stability was from plant species richness via asynchrony (standardized indirect effect = 0.21). This effect was larger and more consistent across experimental sites than the marginally significant direct effect of species richness (standardized path coefficient of direct effect = 0.08, $P = 0.099$). This suggests that much of the effects of plant species richness on stability are explained by asynchrony. Phylogenetic diversity also had strong yet opposing effects on ecosystem stability. Phylogenetic diversity indirectly increased ecosystem stability via its effect on asynchrony (standardized path coefficient of indirect effect = 0.12). Conversely, the unexpected post hoc direct pathway between phylogenetic diversity and stability was negative (standardized path coefficient of direct effect = - 0.10; $P < 0.001$) but weaker, thus

explaining the overall positive relationship between phylogenetic diversity and ecosystem stability (Fig. 2a). Interannual variation in precipitation also had strong, direct, and negative effect on ecosystem stability. Independent of factors related to biodiversity, less variable environmental conditions stabilized plant productivity. Contrary to our expectations, the SEM revealed that neither the functional diversity nor mean of fast-slow leaf traits consistently stabilized (or destabilized) ecosystem productivity ($P > 0.05$). This lack of relationship held when an alternative measure of of functional diversity, functional richness, was used (Fig. S4).

Discussion

The results support our overall hypothesis that multiple components of biodiversity mediate the diversity-stability relationship, principally via their effects on asynchrony. However, the relative importance of certain biological drivers, e.g. functional diversity and fast-slow leaf traits, varied substantially across studies.

The strongest and most consistent driver of stability in the 39 experiments of our study was that of species richness, operating via asynchrony. This is likely to reflect functional niche differences among species that affect their relative performance over time in a temporally variable environment (Yachi & Loreau 1999; Allan *et al.* 2011; Isbell *et al.* 2011; Turnbull *et al.* 2016). However, these niche differences were not captured by our measures of functional (fast-slow) diversity. Instead, the species richness-asynchrony-stability relationship points to a role of traits unrelated to the fast-slow spectrum that stabilize stabilize productivity. Such traits may include rooting strategies, photosynthetic pathways, and regeneration traits (e.g., Edwards *et al.* 2010; Hoover *et al.* 2014; Schroeder-Georg i *et al.* 2016). Data for such traits is relatively sparse (Iversen *et al.* 2017) and the collection of such information should be a priority in addressing the current question and those

related to the components of temporal stability, i.e. resistance and resilience (e.g., Mori *et al.* 2013; Aubin *et al.* 2016).

Species richness also affected ecosystem stability directly, albeit via a weak path that was marginally statistically significant (P = 0.099). We suggest that these effects operated via the greater accumulation of soil organic matter and nutrient stocks in diverse communities (Fornara & Tilman 2008; Oelmann *et al.* 2011; Cong *et al.* 2014), which may be further enhanced by positive effects of diversity on the abundance of soil macro- and micro-organisms, such as earthworms and mycorrhiza, that improve the physical structure of soils (Van Der Heijden *et al.* 2006; Eisenhauer *et al.* 2010, 2012). Further, greater root biomass – which also increases with species richness (e.g., Fornara & Tilman 2008; Mueller *et al.* 2012) – has been found to stabilize ecosystem productivity (Tilman *et al.* 2006) by enhancing nutrient storage and carbohydrate reserves. It is unlikely that these positive feedback effects between plant species richness and environmental conditions operate via asynchrony.

The next most important driver of diversity-stability relationships was phylogenetic diversity. Interestingly, phylogenetic diversity affected ecosystem stability via two different pathways, one positive and operating via asynchrony and one negative and operating directly. The hypothesized indirect pathway was the stronger of these, resulting in a weakly positive overall effect (Fig. 2a) and is likely to be due to a range of phylogenetically conserved traits. Those traits may contain pathogen and herbivore outbreaks to just a few species and therefore a small proportion of community biomass. The direct negative effect was not hypothesized. We suggest that this may reflect a habitat filtering effect possibly related to climatic variability, where a subset of closely related species are better suited to typical site conditions making them more consistently productive over time (Bai *et*

460 *al.* 2004). This path may also reflect experimental communities that are dominated by more
461 inherently stable plant functional groups, such as grasses (Hoover *et al.* 2015; Shi *et al.* 2016).
462
463 Evidence for fast-slow leaf traits affecting ecosystem stability, as either an overall strategy (CWM)
464 or in terms of their functional diversity, was weak. Individual site-level relationships between the
465 CWM of fast-slow traits and stability were often very strong, but extremely variable across sites,
466 ranging from strongly positive to strongly negative and resulting in an overall weak effect. This
467 suggests that the relationship between the fast-slow spectrum and ecosystem stability is heavily
468 dependent upon environmental conditions and the 'matching' of appropriate functional strategies to
469 a site. For example, fast traits may confer ecosystem stability at sites subject to repeated
470 disturbances due to their ability to allow fast recovery, while slow traits may confer ecosystem
471 stability in the face of chronic environmental stresses, such as low nutrient availability or aridity,
472 e.g. the dry grasslands of the experimental sites in Texas included in our study (Wilsey & Polley
473 2004; Wilsey *et al.* 2009). However, we did not detect significant interactions between CWM of the
474 fast-slow spectrum and the multiple descriptors of climate on ecosystem stability (results not
475 presented). Site-level information detailing disturbance regimes and the constancy of soil water
476 availability and nutrient supply may clarify in which environmental conditions particular plant
477 strategies stabilize (or destabilize) biomass production.
478
479 The final driver of ecosystem stability in our models was climate. Inter-annual variation in climate
480 conditions – but not mean annual climate conditions – negatively affected ecosystem stability. This
481 is likely to represent the strong annual variation in the timing and intensity of aboveground biomass
482 production in such environments, e.g. inter-annual variation in the timing and intensity of seasonal
483 rains, and provides evidence that inter-annual variation in climate may be a key fundamental driver
484 of ecosystem stability. As mentioned above, a better characterization of site conditions may provide

485 a more complete understanding of the drivers of ecosystem stability (Ives & Carpenter 2007).
486 Furthermore, initial investigations indicate a powerful interactive role between environmental
487 conditions and biotic community properties (e.g., Xu *et al.* 2015; Yang *et al.* 2017), as abiotic and
488 management factors not only control diversity and productivity but also influence the capacity for
489 diversity to stabilize ecosystem function (Hautier *et al.* 2014; Craven *et al.* 2016). This means that
490 in natural conditions changes in diversity are not the ultimate cause of ecosystem stability, but are
491 an intermediate property of ecosystem response to global change drivers that might also influence
492 ecosystem stability via other pathways. A greater understanding of these interactions and how they
493 operate in natural ecosystems is required to improve both our fundamental understanding of
494 ecosystem stability and to integrate knowledge of diversity-stability relationship into agroecosystem
495 management. With respect to this, our results indicate that the promotion of certain components of
496 diversity (e.g. phylogenetic diversity) would play a greater role than others (e.g. functional diversity
497 of fast-slow traits) in promoting the stability of fodder production. However, the effect of such
498 management on other ecosystem functions and services and their ecosystem stability (e.g.
499 productivity) would also need to be considered (Donohue *et al.* 2016). Threshold-based measures of
500 stability (Oliver *et al.* 2015) may also be more relevant to such applications than the variability
501 measures employed here, as a threshold-based view of ecosystem stability allows under- and
502 overproduction to be viewed differently.
503
504 In conclusion, our study is the first to relate multiple components of biodiversity to ecosystem
505 stability and to estimate their relative importance in driving the diversity-stability relationship.
506 Doing this showed that the role of biodiversity in stabilizing grassland biomass productivity
507 operated via numerous pathways, and therefore that it is more complex and nuanced than has been
508 previously demonstrated. By accounting for multiple drivers of stability, we were also able explain
509 a large amount of variation in both synchrony (conditional R^2 = 78 %) and ecosystem stability

(conditional R^2 = 58 %). In an era of increased climate instability (Goodess 2013; Stott 2016), it is important to gain a greater understanding of each of these component processes so that the functional benefits of biodiversity may be effectively conserved and promoted.

Acknowledgements

This paper is a joint effort of the sTABILITY group funded by sDiv (www.idiv.de/stability), the Synthesis Centre of the German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig (DFG FZT 118). The Jena Experiment is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation; FOR 1451). The Cedar Creek biodiversity experiments are supported by the US National Science Foundation (LTER Award 1234162). The study has been supported by the TRY initiative on plant traits (http://www.try-db.org). The TRY initiative and database is hosted, developed, and maintained by J. Kattge and G. Boenisch (Max Planck Institute for Biogeochemistry, Jena, Germany). TRY is currently supported by DIVERSITAS/Future Earth and the German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig. The authors would also like to thank Jon Lefcheck for his help in revising the structure equation model.

536

537

538

539 **References**

540

541 Allan, E., Weisser, W., Weigelt, A., Roscher, C., Fischer, M. & Hillebrand, H. (2011). More diverse
542 plant communities have higher functioning over time due to turnover in complementary dominant
543 species. *Proc. Natl. Acad. Sci.*, 108, 17034–17039

Aubin, I., Munson, A.D., Cardou, F., Burton, P.J., Isabel, N., Pedlar, J.H., *et al.* (2016). Traits to stay, traits to move: a review of functional traits to assess sensitivity and adaptive capacity of temperate and boreal trees to climate change. *Environ. Rev.*, 24, 164–186

Bai, Y., Han, X., Wu, J., Chen, Z. & Li, L. (2004). Ecosystem stability and compensatory effects in the Inner Mongolia grassland. *Nature*, 431, 181–184

Balvanera, P., Pfisterer, A.B., Buchmann, N., He, J.-S., Nakashizuka, T., Raffaelli, D., *et al.* (2006). Quantifying the evidence for biodiversity effects on ecosystem functioning and services. *Ecol. Lett.*, 9, 1146–1156

Bartlett, M.K., Scoffoni, C. & Sack, L. (2012). The determinants of leaf turgor loss point and prediction of drought tolerance of species and biomes: a global meta-analysis. *Ecol. Lett.*, 15, 393–405

Boyle, B., Hopkins, N., Lu, Z., Raygoza Garay, J.A., Mozzherin, D., Rees, T., *et al.* (2013). The taxonomic name resolution service: an online tool for automated standardization of plant names. *BMC Bioinformatics*, 14, 16

Burnham, K. & Anderson, D. (2003). *Model Selection and Multimodel Inference: a practical information-theoretic approach*. 2nd edn. Springer-Verlag

Cadotte, M.W. (2015). Phylogenetic diversity and productivity: gauging interpretations from experiments that do not manipulate phylogenetic diversity. *Funct. Ecol.*, 29, 1603–1606

Cadotte, M.W., Dinnage, R. & Tilman, D. (2012). Phylogenetic diversity promotes ecosystem stability. *Ecology*, 93, S223–S233

Campbell, V., Murphy, G. & Romanuk, T.N. (2011). Experimental design and the outcome and interpretation of diversity–stability relations. *Oikos*, 120, 399–408

Cong, W.-F., Ruijven, J., Mommer, L., De Deyn, G.B., Berendse, F. & Hoffland, E. (2014). Plant species richness promotes soil carbon and nitrogen stocks in grasslands without legumes. *J. Ecol.*, 102, 1163–1170

Craine, J.M., Ocheltree, T.W., Nippert, J.B., Towne, E.G., Skibbe, A.M., Kembel, S.W., *et al.* (2013). Global diversity of drought tolerance and grassland climate-change resilience. *Nat. Clim Change*, 3, 63–67

Craven, D., Isbell, F., Manning, P., Connolly, J., Bruelheide, H., Ebeling, A., *et al.* (2016). Plant diversity effects on grassland productivity are robust to both nutrient enrichment and drought. *Philos. Trans. R. Soc. Lond. B Biol. Sci.*, 371

Daneshgar, P.P., Polley, H.W. & Wilsey, B.J. (2013). Simple plant traits explain functional group diversity decline in novel grassland communities of Texas. *Plant Ecol.*, 214, 231–241

Díaz, S. & Cabido, M. (2001). Vive la différence: plant functional diversity matters to ecosystem processes. *Trend Ecol. Evol.*, 16, 646–655

Díaz, S., Kattge, J., Cornelissen, J.H.C., Wright, I.J., Lavorel, S., Dray, S., *et al.* (2016). The global spectrum of plant form and function. *Nature*, 529, 167–171

Doak, D.F., Bigger, D., Harding, E., Marvier, M., O'malley, R. & Thomson, D. (1998). The statistical inevitability of stability-diversity relationships in community ecology. *Am. Nat.*, 151, 264–276

Donohue, I., Hillebrand, H., Montoya, J.M., Petchey, O.L., Pimm, S.L., Fowler, M.S., *et al.* (2016). Navigating the complexity of ecological stability. *Ecol. Lett.*, 19, 1172–1185

Edwards, E.J., Osborne, C.P., Strömberg, C.A.E. & Smith, S.A. (2010). The Origins of C4 Grasslands: Integrating Evolutionary and Ecosystem Science. *Science*, 328, 587–591

Eisenhauer, N., Bebler, H., Engels, C., Gleixner, G., Habekost, M., Milcu, A., *et al.* (2010). Plant diversity effects on soil microorganisms support the singular hypothesis. *Ecology*, 91, 485–496

Eisenhauer, N., Reich, P.B. & Isbell, F. (2012). Decomposer diversity and identity influence plant diversity effects on ecosystem functioning. *Ecology*, 93, 2227–2240

Fargione, J. & Tilman, D. (2005). Niche differences in phenology and rooting depth promote coexistence with a dominant C4 bunchgrass. *Oecologia*, 143, 598–606

Fischer, F.M., Wright, A.J., Eisenhauer, N., Ebeling, A., Roscher, C., Wagg, C., *et al.* (2016). Plant species richness and functional traits affect community stability after a flood event. *Philos. Trans. R. Soc. Lond. B Biol. Sci.*, 371

Flynn, D.F.B., Mirotchnick, N., Jain, M., Palmer, M.I. & Naeem, S. (2011). Functional and phylogenetic diversity as predictors of biodiversity-ecosystem-function relationships. *Ecology*, 92, 1573–1581

Fornara, D. & Tilman, D. (2008). Plant functional composition influences rates of soil carbon and nitrogen accumulation. *J. Ecol.*, 96, 314–322

Gilbert, G.S., Magarey, R., Suiter, K. & Webb, C.O. (2012). Evolutionary tools for phytosanitary risk analysis: phylogenetic signal as a predictor of host range of plant pests and pathogens. *Evol. Appl.*, 5, 869–878

Godoy, O., Kraft, N.J.B. & Levine, J.M. (2014). Phylogenetic relatedness and the determinants of competitive outcomes. *Ecol. Lett.*, 17, 836–844

Gomez, J.M., Verdu, M. & Perfectti, F. (2010). Ecological interactions are evolutionarily conserved across the entire tree of life. *Nature*, 465, 918–921

Goodess, C.M. (2013). How is the frequency, location and severity of extreme events likely to change up to 2060? *Environ. Sci. Policy*, 27, Supplement 1, S4–S14
Gould, I.J., Quinton, J.N., Weigelt, A., De Deyn, G.B. & Bardgett, R.D. (2016). Plant diversity and root traits benefit physical properties key to soil function in grasslands. *Ecol. Lett.*, 19, 1140–1149

Grime, J.P. (1977). Evidence for the existence of three primary strategies in plants and its relevance to ecological and evolutionary theory. *Am. Nat.*, 111, 1169–1194

Grime, J.P., Hodgson, J.G. & Hunt, R. (2007). *Comparative plant ecology: a functional approach to common British species*. Springer

Gross, K., Cardinale, B.J., Fox, J.W., Gonzalez, A., Loreau, M., Polley, H.W., *et al.* (2014). Species Richness and the Temporal Stability of Biomass Production: A New Analysis of Recent Biodiversity Experiments. *Am. Nat.*, 183, 1–12

Hautier, Y., Seabloom, E.W., Borer, E.T., Adler, P.B., Harpole, W.S., Hillebrand, H., *et al.* (2014). Eutrophication weakens stabilizing effects of diversity in natural grasslands. *Nature*, 508, 521–525

Hautier, Y., Tilman, D., Isbell, F., Seabloom, E.W., Borer, E.T. & Reich, P.B. (2015). Anthropogenic environmental changes affect ecosystem stability via biodiversity. *Science*, 348, 336–340

Hector, A., Hautier, Y., Saner, P., Wacker, L., Bagchi, R., Joshi, J., *et al.* (2010). General stabilizing effects of plant diversity on grassland productivity through population asynchrony and overyielding. *Ecology*, 91, 2213–2220

Hoover, D.L., Duniway, M.C. & Belnap, J. (2015). Pulse-drought atop press-drought: unexpected plant responses and implications for dryland ecosystems. *Oecologia*, 179, 1211–1221

Hoover, D.L., Knapp, A.K. & Smith, M.D. (2014). Resistance and resilience of a grassland ecosystem to climate extremes. *Ecology*, 95, 2646–2656

Huxman, T.E., Smith, M.D., Fay, P.A., Knapp, A.K., Shaw, M.R., Loik, M.E., *et al.* (2004). Convergence across biomes to a common rain-use efficiency. *Nature*, 429, 651–654

Isbell, F., Calcagno, V., Hector, A., Connolly, J., Harpole, W.S., Reich, P.B., *et al.* (2011). High plant diversity is needed to maintain ecosystem services. *Nature*, 477, 199–202

Isbell, F., Craven, D., Connolly, J., Loreau, M., Schmid, B., Beierkuhnlein, C., *et al.* (2015a). Biodiversity increases the resistance of ecosystem productivity to climate extremes. *Nature*, 526, 574–577

Isbell, F., Tilman, D., Polasky, S. & Loreau, M. (2015b). The biodiversity-dependent ecosystem service debt. *Ecol. Lett.*, 18, 119–134

Isbell, F.I., Polley, H.W. & Wilsey, B.J. (2009). Biodiversity, productivity and the temporal stability of productivity: patterns and processes. *Ecol. Lett.*, 12, 443–451

Iversen, C.M., McCormack, M.L., Powell, A.S., Blackwood, C.B., Freschet, G.T., Kattge, J., *et al.* (2017). A global Fine-Root Ecology Database to address below-ground challenges in plant ecology. *New Phytol.*, in press (accepted)

Ives, A.R. & Carpenter, S.R. (2007). Stability and Diversity of Ecosystems. *Science*, 317, 58–62

Jiang, L. & Pu, Z. (2009). Different effects of species diversity on temporal stability in single-trophic and multitrophic communities. *Am. Nat.*, 174, 651–659

Kattge, J., Diaz, S., Lavorel, S., Prentice, I.C., Leadley, P., BÖNisch, G., *et al.* (2011). TRY - a global database of plant traits. *Glob. Change Biol.*, 17, 2905–2935

Knapp, G. & Hartung, J. (2003). Improved tests for a random effects meta-regression with a single covariate. *Stat. Med.*, 22, 2693–2710

Laliberté, E. & Legendre, P. (2010). A distance-based framework for measuring functional diversity from multiple traits. *Ecology*, 91, 299–305

Lange, M., Eisenhauer, N., Sierra, C.A., Bessler, H., Engels, C., Griffiths, R.I., *et al.* (2015). Plant diversity increases soil microbial activity and soil carbon storage. *Nat. Commun.*, 6, 6707

Lefcheck, J.S. (2016). piecewiseSEM: Piecewise structural equation modelling in r for ecology, evolution, and systematics. *Methods Ecol. Evol.*, 7, 573–579

Lehman, C.L. & Tilman, D. (2000). Biodiversity, Stability, and Productivity in Competitive Communities. *Am. Nat.*, 156, 534–552

Loreau, M. & Mazancourt, C. de. (2008). Species synchrony and its drivers: neutral and nonneutral community dynamics in fluctuating environments. *Am. Nat.*, 172, E48–E66

Májeková, M., de Bello, F., Doležal, J. & Lepš, J. (2014). Plant functional traits as determinants of population stability. *Ecology*, 95, 2369–2374

Maron, J.L., Marler, M., Klironomos, J.N. & Cleveland, C.C. (2011). Soil fungal pathogens and the relationship between plant diversity and productivity. *Ecol. Lett.*, 14, 36–41

May, R.M. (1973). *Stability and complexity in model ecosystems*. Princeton University Press

de Mazancourt, C., Isbell, F., Larocque, A., Berendse, F., De Luca, E., Grace, J.B., *et al.* (2013). Predicting ecosystem stability from community composition and biodiversity. *Ecol. Lett.*, 16, 617–625

McDowell, N., Pockman, W.T., Allen, C.D., Breshears, D.D., Cobb, N., Kolb, T., *et al.* (2008). Mechanisms of plant survival and mortality during drought: why do some plants survive while others succumb to drought? *New Phytol.*, 178, 719–739

McNaughton, S.J. (1978). Stability and diversity of ecological communities. *Nature*, 274, 251–253

Mori, A.S., Furukawa, T. & Sasaki, T. (2013). Response diversity determines the resilience of ecosystems to environmental change. *Biol. Rev.*, 88, 349–364

Mueller, K.E., Tilman, D., Fornara, D.A. & Hobbie, S.E. (2012). Root depth distribution and the diversity–productivity relationship in a long-term grassland experiment. *Ecology*, 94, 787–793

Oelmann, Y., Buchmann, N., Gleixner, G., Habekost, M., Roscher, C., Rosenkranz, S., *et al.* (2011). Plant diversity effects on aboveground and belowground N pools in temperate grassland ecosystems: Development in the first 5 years after establishment. *Glob. Biogeochem. Cycles*, 25

Oliver, T.H., Heard, M.S., Isaac, N.J.B., Roy, D.B., Procter, D., Eigenbrod, F., *et al.* (2015). Biodiversity and Resilience of Ecosystem Functions. *Trends Ecol. Evol.*, 30, 673–684

Parker, I.M., Saunders, M., Bontrager, M., Weitz, A.P., Hendricks, R., Magarey, R., *et al.* (2015). Phylogenetic structure and host abundance drive disease pressure in communities. *Nature*, 520, 542–544

Pearse, W.D., Cadotte, M.W., Cavender-Bares, J., Ives, A.R., Tucker, C.M., Walker, S.C., *et al.* (2015). pez: phylogenetics for the environmental sciences. *Bioinformatics*, 31, 2888–2890

Pearse, W.D., Purvis, A., Cavender-Bares, J. & Helmus, M.R. (2014). Metrics and models of community phylogenetics. In: *Modern phylogenetic comparative methods and their application in evolutionary biology*. Springer Berlin Heidelberg, pp. 451–464

Pérez-Harguindeguy, N., Díaz, S., Garnier, E., Lavorel, S., Poorter, H., Jaureguiberry, P., *et al.* (2013). New handbook for standardised measurement of plant functional traits worldwide. *Aust. J. Bot.*, 61, 167–234

Polley, H.W., Isbell, F.I. & Wilsey, B.J. (2013). Plant functional traits improve diversity-based predictions of temporal stability of grassland productivity. *Oikos*, 122, 1275–1282

R Core Team. (2016). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria

Reich, P.B. (2014). The world-wide "fast–slow" plant economics spectrum: a traits manifesto. *J. Ecol.*, 102, 275–301

Reinhart, K.O., Wilson, G.W.T. & Rinella, M.J. (2012). Predicting plant responses to mycorrhizae: integrating evolutionary history and plant traits. *Ecol. Lett.*, 15, 689–695

Roscher, C., Schumacher, J., Gubsch, M., Lipowsky, A., Weigelt, A., Buchmann, N., *et al.* (2012). Using Plant Functional Traits to Explain Diversity–Productivity Relationships. *PLoS ONE*, 7, e36760

Roscher, C., Weigelt, A., Proulx, R., Marquard, E., Schumacher, J., Weisser, W.W., *et al.* (2011). Identifying population- and community-level mechanisms of diversity–stability relationships in experimental grasslands. *J. Ecol.*, 99, 1460–1469

Sala, O.E., Gherardi, L.A., Reichmann, L., Jobbágy, E. & Peters, D. (2012). Legacies of precipitation fluctuations on primary production: theory and data synthesis. *Philos. Trans. R. Soc. Lond. B Biol. Sci.*, 367, 3135–3144

Salguero-Gómez, R., Jones, O.R., Jongejans, E., Blomberg, S.P., Hodgson, D.J., Mbeau-Ache, C., *et al.* (2016). Fast–slow continuum and reproductive strategies structure plant life-history variation worldwide. *Proc. Natl. Acad. Sci.*, 113, 230–235

Schnitzer, S.A., Klironomos, J.N., HilleRisLambers, J., Kinkel, L.L., Reich, P.B., Xiao, K., *et al.* (2011). Soil microbes drive the classic plant diversity–productivity pattern. *Ecology*, 92, 296–303

Schroeder-Georgi, T., Wirth, C., Nadrowski, K., Meyer, S .T., Mommer, L. & Weigelt, A. (2016). From pots to plots: hierarchical trait-based prediction of plant performance in a mesic grassland. *J. Ecol.*, 104, 206–218

Shi, Z., Xu, X., Souza, L., Wilcox, K., Jiang, L., Liang, J., *et al.* (2016). Dual mechanisms regulate ecosystem stability under decade-long warming and hay harvest. *Nat Commun*, 7

Spasojevic, M.J. & Suding, K.N. (2012). Inferring community assembly mechanisms from functional diversity patterns: the importance of multiple assembly processes. *J. Ecol.*, 100, 652–661

Stott, P. (2016). How climate change affects extreme weather events. *Science*, 352, 1517–1518

Tilman, D. & Downing, J.A. (1994). Biodiversity and stability in grasslands. *Nature*, 367, 363–365

Tilman, D., Reich, P.B. & Knops, J.M.H. (2006). Biodiversity and ecosystem stability in a decade-long grassland experiment. *Nature*, 441, 629–632

Tredennick, A.T., de Mazancourt, C., Loreau, M. & Adler, P.B. (2017). Environmental responses, not species interactions, determine synchrony of dominant species in semiarid grasslands. *Ecology*, in press (accepted)

Turnbull, L.A., Isbell, F., Purves, D.W., Loreau, M. & Hector, A. (2016). Understanding the value of plant diversity for ecosystem functioning through niche theory. *Proc. R. Soc. Lond. B Biol. Sci.*, 283

Van Der Heijden, M.G.A., Streitwolf-Engel, R., Riedl, R., Siegrist, S., Neudecker, A., Ineichen, K., *et al.* (2006). The mycorrhizal contribution to plant productivity, plant nutrition and soil structure in experimental grassland. *New Phytol.*, 172, 739–752

Van Ruijven, J. & Berendse, F. (2010). Diversity enhances community recovery, but not resistance, after drought. *J. Ecol.*, 98, 81–86

Viechtbauer, W. (2010). Conducting Meta-Analyses in R with the metafor Package. *J. Stat. Softw.*, 36, 1–48

Wacker, L., Baudois, O., Eichenberger-Glinz, S. & Schmid, B. (2009). Diversity effects in early- and mid-successional species pools along a nitrogen gradient. *Ecology*, 90, 637–48

Webb, C.O., Ackerly, D.D., McPeek, M.A. & Donoghue, M.J. (2002). Phylogenies and Community Ecology. *Annu. Rev. Ecol. Syst.*, 33, 475–505

Weigelt, A., Schumacher, J., Roscher, C. & Schmid, B. (2008). Does biodiversity increase spatial stability in plant community biomass? *Ecol. Lett.*, 11, 338–347

Wilsey, B.J. & Polley, H.W. (2004). Realistically Low Species Evenness Does Not Alter Grassland Species-Richness-Productivity Relationships. *Ecology*, 85, 2693–2700

Wilsey, B.J., Teaschner, T.B., Daneshgar, P.P., Isbell, F.I. & Polley, H.W. (2009). Biodiversity maintenance mechanisms differ between native and novel exotic-dominated communities. *Ecol. Lett.*, 12, 432–442

Wright, A.J., Kroon, H. de, Visser, E.J.W., Buchmann, T., Ebeling, A., Eisenhauer, N., *et al.* (2016). Plants are less negatively affected by flooding when growing in species-rich plant communities. *New Phytol.*

Wright, I.J., Reich, P.B., Westoby, M., Ackerly, D.D., Baruch, Z., Bongers, F., *et al.* (2004). The worldwide leaf economics spectrum. *Nature*, 428, 821–827

Xu, Z., Ren, H., Li, M.-H., van Ruijven, J., Han, X., Wan, S., *et al.* (2015). Environmental changes drive the temporal stability of semi-arid natural grasslands through altering species asynchrony. *J. Ecol.*, 103, 1308–1316

Yachi, S. & Loreau, M. (1999). Biodiversity and ecosystem productivity in a fluctuating environment: The insurance hypothesis. *Proc. Natl. Acad. Sci.*, 96, 1463–1468

Yang, Z., Zhang, Q., Su, F., Zhang, C., Pu, Z., Xia, J., *et al.* (2017). Daytime warming lowers community temporal stability by reducing the abundance of dominant, stable species. *Glob. Change Biol.*, 23, 154–163

Zanne, A.E., Tank, D.C., Cornwell, W.K., Eastman, J.M., Smith, S.A., FitzJohn, R.G., *et al.* (2014). Three keys to the radiation of angiosperms into freezing environments. *Nature*, 506, 89–92

Zuur, A.F., Ieno, E.N. & Elphick, C.S. (2010). A protocol for data exploration to avoid common statistical problems. *Methods Ecol. Evol.*, 1, 3–14