

THE COMPLEX LORENZ EQUATIONS

A. C. FOWLER

Department of Mathematics, Massachusetts Institute of Technology, Cambridge, Mass. 02139, USA

J. D. GIBBON

Department of Mathematics, Imperial College of Science & Technology, London SW7 2BZ, GB

and

M. J. McGUINNESS

Department of Mathematical Physics, University College Dublin, Belfield, Dublin 4, Republic of Ireland

Received 11 March 1981

We have undertaken a study of the complex Lorenz equations

$$\begin{aligned}\dot{x} &= -\sigma x + \sigma y, \\ \dot{y} &= (r - z)x - ay, \\ \dot{z} &= -bz + \frac{1}{2}(x^*y + xy^*),\end{aligned}$$

where x and y are complex and z is real. The complex parameters r and a are defined by $r = r_1 + ir_2$; $a = 1 - ie$ and σ and b are real. Behaviour remarkably different from the real Lorenz model occurs. Only the origin is a fixed point except for the special case $e + r_2 = 0$. We have been able to determine analytically two critical values of r_1 , namely r_{1c} and r_{1c} . The origin is a stable fixed point for $0 < r_1 < r_{1c}$, but for $r_1 > r_{1c}$ a Hopf bifurcation to a limit cycle occurs. We have an *exact* analytic solution for this limit cycle which is always stable if $\sigma < b + 1$. If $\sigma > b + 1$ then this limit is only stable in the region $r_{1c} < r_1 < r_{1c}$. When $r_1 > r_{1c}$, a transition to a finite amplitude oscillation about the limit cycle occurs. The nature of this bifurcation is studied in detail by using a multiple time scale analysis to derive the Stuart–Landau amplitude equation from the original equations in a frame rotating with the limit cycle frequency. This latter bifurcation is either a sub- or super-critical Hopf-like bifurcation to a doubly periodic motion, the direction of bifurcation depending on the parameter values. The nature of the bifurcation is complicated by the existence of a zero eigenvalue.

1. Introduction

In a previous paper [1], the so-called “complex” Lorenz equations were derived as a generalization of the original equations first derived by Lorenz [2]. The complex equations are written in the form

$$\dot{x} = -\sigma x + \sigma y, \quad (1.1a)$$

$$\dot{y} = -xz + rx - ay, \quad (1.1b)$$

$$\dot{z} = -bz + \frac{1}{2}(x^*y + xy^*). \quad (1.1c)$$

The Rayleigh number r and the parameter a are

complex numbers defined by

$$r = r_1 + ir_2, \quad (1.2a)$$

$$a = 1 - ie, \quad (1.2b)$$

and σ , b , r_1 , r_2 and e are real and positive. The form of (1.1) shows that x and y are complex but z is real.

The intention of this paper is to make a mathematical study of eqs. (1.1) in order to show that significantly different behaviour occurs in the bifurcation sequence than in the

real Lorenz equations. The real Lorenz model is embedded in (1.1) and can be recovered when $r_2 = e = 0$ and x and y are real. The physical motivation for studying the complex Lorenz model comes from work by two of the three present authors who derived a set of amplitude equations near criticality for a class of dispersively unstable, weakly nonlinear, weakly damped physical systems [1, 4]. This was based on work which considered the undamped systems first [3] and then weak damping was added afterwards.

When only temporal variation was included, the relevant set of amplitude equations turned out to be of the form

$$\frac{d^2 A}{dT^2} + \Delta_1 \frac{dA}{dT} = \alpha A - \beta AB, \quad (1.3a)$$

$$\frac{dB}{dT} + \Delta_2 B = \frac{d}{dT} |A|^2 + \Delta_3 |A|^2. \quad (1.3b)$$

In the circumstances discussed in [1], the parameters β , Δ_2 and Δ_3 were always real and positive. If only weak damping is added to the original system then α and Δ_1 are also real and positive. This is the case for the laser equations [1, 4]. If extra weak dispersive effects are also added then it turns out that α and Δ_1 become complex. This circumstance occurs in the 2-layer and Eady models of baroclinic instability [1, 5] when a "weak" beta-effect is included. Eqs. (1.3) can be transformed into eqs. (1.1) by the following transformations:

$$t = \Omega T, \quad (1.4a)$$

$$\Omega = \text{Re}(\Delta_1) - \Delta_1/2, \quad (1.4b)$$

$$x = (2\beta)^{1/2} \Omega^{-1} A, \quad (1.4c)$$

$$z = 2\beta \Omega^{-1} \Delta_3^{-1} B, \quad (1.4d)$$

where the variables r , α , a and b in (1.1) are given by

$$\alpha = \Delta_3/2\Omega,$$

$$b = \Delta_3/\Omega,$$

$$r_1 = 1 + 2\text{Re}(\alpha)/(\Delta_3\Omega),$$

$$r_2 = [2\text{Im}(\alpha) + \Delta_1 \text{Im}(\Delta_1)]/(\Omega\Delta_3),$$

$$e = -\text{Im}(\Delta_1)/\Omega.$$

The complex Lorenz equations also form the basic model for bistable optical systems of two level atoms. Hassan, Drummond and Walls [19] and Ikeda [20] have shown that in a single mode high-Q ring cavity, the semi-classical equations of motion are a set of damped Maxwell-Bloch type equations. By observation we can show that these can be transformed into eqs. (1.1), but with an external driving field. In this case, $r_2 = 0$. But $e \neq 0$. Refs. 19 and 20 extend earlier results by Bonifacio and Lugiato [21]. As a physical problem, the driving field is necessary and obviously the inclusion of this adds an extra degree of complication to the problem. A study of the system (1.1) on its own is therefore important as a first step towards understanding the more general problem, in which a forcing term is present.

However, the essence of this paper is not a discussion of the physical derivation of (1.1) since this was performed in [1] but rather we seek to undertake a mathematical and simple numerical analysis of these equations in their own right. A consideration of this more general system may also cast more light on the real Lorenz model.

Our approach is analytical, but we should point out that the analytical results obtained here were strongly motivated by the results of numerical computations. An example of this is the exact periodic solution displayed in section 2, the form of which was suggested by numerical computations which gave a perfect ellipse in the $\text{Re}(x) - \text{Re}(y)$ phase plane with z quickly reaching a constant equilibrium value. The analysis of the bifurcation of the limit cycle to doubly periodic solutions given in section 3 was also suggested by numerical results which showed fast and slow oscillations, thereby indicating that a multiple time scale calculation

was in order. It is interesting, in our view, that so much information can be derived by analytical means for what is essentially a fifth order problem. A qualitative difference between the first two bifurcations is that while the first is a supercritical Hopf bifurcation of the origin into a limit cycle, the nature of the second is blurred by the existence of a $\lambda = 0$ eigenvalue which occurs because of rotational symmetry. For this reason, the aims of sections 2 and 3 are different. In section 2, following Lorenz in which we use r_1 as the main bifurcation parameter, we calculate the two critical values of r_1 at which firstly, the origin bifurcates to the limit cycle and secondly, at which this limit cycle becomes unstable. Section 3 is devoted to understanding the nature of this bifurcation using multiple scales to obtain the Stuart–Landau equation in the rotating frame,

$$\frac{dA}{dt} = k_1 A + k_2 A |A|^2. \quad (1.5)$$

Following McLaughlin and Martin [6], the criterion for determining whether the bifurcation is super- or sub-critical is determined by whether $\text{Re}(k_2)$ is negative or positive respectively and consequently whether the limit cycle undergoes ‘soft’ bifurcation to a doubly periodic solution, or ‘hard’ bifurcation to some other type of motion.

2. An exact periodic solution

In studying (1.1) we shall follow Lorenz’ analysis of the nature and stability of solutions. Equilibrium solutions in which time derivatives are absent are given by the origin $x = y = z = 0$, or (from (1.1a)) $x = y$ whence $z = r - a$ from (1.1b). Eq. (1.1c) thus implies that

$$|x|^2 = b(r - a). \quad (2.1)$$

Since z is real, it follows that such points can

only exist if $\text{Im}(r - a) = 0$; that is,

$$e + r_2 = 0. \quad (2.2)$$

In this case, there is a *continuum* of steady states given by

$$z = r_1 - 1, \quad (2.3a)$$

$$|x| = |y| = [b(r_1 - 1)]^{1/2}. \quad (2.3b)$$

This rather pathological possibility already reveals the special nature of (1.1). In [1] it was found for the baroclinic two layer model with weak dissipation and weak beta-effect that

$$e = 3r_2, \quad (2.4)$$

$$\sigma = 2. \quad (2.5)$$

The special condition (2.2) is not satisfied and so only the origin is a fixed point. In the case $e = r_2 = 0$ we technically do not return to the full real Lorenz equations as x and y can still remain complex. The two fixed points of the real Lorenz equations (in addition to the origin) are replaced by the continuum of points (2.3) although there is very little difference in this intermediate case from the real case.

2.1. Stability of the origin

We examine the stability of the steady state $(0, 0, 0)$ by linearising (1.1) about this point. To do so we simply neglect quadratic terms, thus

$$\begin{pmatrix} \dot{x} \\ \dot{y} \\ \dot{z} \end{pmatrix} = \begin{pmatrix} -\sigma & \sigma & 0 \\ r & -a & 0 \\ 0 & 0 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix}. \quad (2.6)$$

We note that $z = 0$ is always a stable manifold ($\dot{z} = -bz$). All solutions to (2.6) are proportional to $\exp(\lambda t)$, where the eigenvalues λ (assumed distinct) are given by $\lambda = -b$, and $(\sigma + \lambda)(a + \lambda) - r\sigma = 0$, whence

$$\lambda = \frac{1}{2}[-(\sigma + a) \pm \{(\sigma + a)^2 + 4\sigma(r - a)\}^{1/2}]. \quad (2.7)$$

If either value of λ in (2.7) has $\text{Re}(\lambda) > 0$, then the origin is said to be linearly unstable. Note that the two values of λ in (2.7) are not generally complex conjugates; this is because the characteristic equation for λ does not have real coefficients. If (as can easily be done) the eqs. (2.1) are first written out as a five by five system for the variables

$$x_R = \text{Re}(x); \quad x_I = \text{Im}(x); \quad y_R = \text{Re}(y); \\ y_I = \text{Im}(y); \quad z,$$

then the resulting matrix equation analogous to (2.6) would have eigenvalues $-b, \lambda$ given by (2.7) together with λ^* .

Let us define

$$p + iq = \{(\sigma + a)^2 + 4\sigma(r - a)\}^{1/2}; \quad p > 0 \quad (2.8)$$

(this can be done without loss of generality). Then (2.7) gives the eigenvalues as

$$\lambda = \frac{1}{2}[-(\sigma + a) \pm (p + iq)], \quad (2.9)$$

so that

$$\text{Re}(\lambda) = \frac{1}{2}[-\sigma - a \pm p]. \quad (2.10)$$

It follows (remember $p > 0$) that one eigenvalue always has negative real part, and the other is negative or positive depending as $p \leq \sigma + 1$. The critical stability limit is when

$$p = \sigma + 1, \quad (2.11)$$

and this relation determines a corresponding relation between the parameters which, plotted as a curve in parameter space, divides regions of stability from those of instability. From (2.8), we find

$$p^2 - q^2 = 2ipq = (\sigma + a)^2 + 4\sigma(r - a),$$

whence a little algebra shows that

$$p^2 - q^2 = (\sigma + 1)^2 + 4\sigma(r_1 - 1) - e^2, \\ pq = 2\sigma(e + r_2) - e(\sigma + 1). \quad (2.12)$$

Following Lorenz, we use r_1 as the bifurcation parameter, and denote its value at the stability limit by r_{1c} ; thus when $r_1 = r_{1c}$, $p = \sigma + 1$, so (2.12) implies

$$e^2 - q^2 = 4\sigma(r_1 - 1), \\ e + q = \frac{2\sigma(e + r_2)}{\sigma + 1}. \quad (2.13)$$

Note immediately that the frequency ω of the critically stable eigenmode is given, from (2.9), by

$$\omega = \text{Im} \lambda = \frac{1}{2}(e + q).$$

Eq. (2.13) implies

$$\omega = \frac{\sigma(e + r_2)}{\sigma + 1}. \quad (2.14)$$

We may observe that if $e + r_2 \neq 0$, then the origin becomes oscillatorily unstable, so that the conditions for a Hopf bifurcation will occur (provided also $d(\text{Re} \lambda)/dr_1 \neq 0$ at $r_1 = r_{1c}$): thus we may expect a limit cycle to bifurcate from the origin at $r_1 = r_{1c}$, with approximate frequency ω ; hence as $e + r_2 \rightarrow 0$, the frequency tends to zero, and so the continuum of equilibrium points (2.3) may be interpreted as the limit of a limit cycle in which the frequency has decreased to zero. We will generally suppose $\omega \neq 0$.

Eliminating q in (2.13) we obtain

$$r_{1c} = 1 + \frac{(e + r_2)(e - \sigma r_2)}{(\sigma + 1)^2} \quad (2.15)$$

as the critical value of r_1 . It is easy to see from (2.12) that if $r_1 < r_{1c}$, then $p < \sigma + 1$ (and vice versa), so that the origin is linearly stable for $r_1 < r_{1c}$, and linearly unstable for $r_1 > r_{1c}$.

2.2. Limit cycle

From (2.12), it is also easy to check that $d(\text{Re } \lambda)/dr_1 > 0$ at $r_1 = r_c$, so that (provided (1.1) is written as a five by five system, and considering the previous remarks about eigenvalues in this case) the Hopf theorem is applicable, and thus a limit cycle does bifurcate from the origin at $r = r_{lc}$, provided $\omega \neq 0$. Approximate techniques for giving the form of this exist (for $|r_1 - r_{lc}| \ll 1$). However, numerical computation (fig. 1) shows that even for $r_1 - r_{lc} \approx 1$, the limit

cycle appears to be an ellipse in the (x_R, y_R) plane (R denoting real part). Also in the (x_R, z) plane it is evident that z quickly reaches a constant value.

This suggests that we look for an *exact* solution to the equations in which z is constant, and x and y are sinusoidal. Since (1) is linear in x and y if z is constant, it is clear that such a solution is possible. We put

$$x = A e^{i\Omega t}, \quad y = B e^{i\Omega t}, \quad z = M. \quad (2.16)$$




Fig. 1. (a) Stable elliptical limit cycle in the x_R - y_R plane with parameter values $b = 4/3$, $\sigma = 2$, $r_1 = 2$, $r_2 = 1$, $e = 3$.




Fig. 1. (b) Plot of z versus y_R for the limit cycle in (a) showing z reaching a constant value.

Then (1.1) implies

$$(a + if)\left(1 + \frac{if}{\sigma}\right) = r - |A|^2/b. \quad (2.19)$$

$$ifA \approx -\sigma A + \sigma B,$$

$$ifB = (r - M)A - aB, \quad (2.17)$$

$$0 = \frac{1}{2}(AB^* + A^*B) - bM,$$

Equating real and imaginary parts of (2.19) shows that

so that we find

$$f = \omega = \frac{\sigma(e + r_2)}{\sigma + 1}, \quad (2.20)$$

$$B = \left(1 + \frac{if}{\sigma}\right)A; \quad M = |A|^2/b, \quad (2.18)$$

and

and f and $|A|$ are given by

$$|A|^2 = b(r_1 - r_{1c}). \quad (2.21)$$

(2.20) and (2.21) conform with the dictates of Hopf's theorem, but show that the limit cycle can be determined exactly up to a phase factor in A . The amplitude increases with $(r_1 - r_{1c})^{1/2}$ so that the bifurcation is supercritical and hence stable (at least initially) and the frequency of the oscillation remains constant. The solution can be written

$$x = A e^{i\omega t}; \quad y = \left(1 + \frac{i\omega}{\sigma}\right) A e^{i\omega t}; \quad z = |A|^2/b. \quad (2.22)$$

We note that such exact limit cycle solutions may exist for other ordinary differential systems, and in this context we draw attention to the papers of Fujisaka and Yamada [15] and Escher [16].

2.3. Stability of the limit cycle

Normally, one cannot explicitly study the stability of a limit cycle, since the basic state is rarely susceptible to analysis. It is therefore very fortunate in the present case that the stability may be explicitly examined, since the limit cycle is exactly known. It is possible to calculate the critical value of r_1 where the limit cycle becomes unstable, which we denote by r_{1c} . This calculation however does not enlighten us as to the qualitative nature of this bifurcation and so this qualitative study is left to section 3 and we confine ourselves to purely calculating r_{1c} .

Firstly we change variables to those with respect to a rotating frame, which reduces the limit cycle to a fixed point. That is, we put

$$x = X e^{i\omega t}; \quad y = Y e^{i\omega t}; \quad z = Z, \quad (2.23)$$

where ω is given by (2.20). X, Y, Z satisfy

$$\begin{aligned} \dot{X} &= -(\sigma + i\omega)X + \sigma Y, \\ \dot{Y} &= (r - Z)X - (a + i\omega)Y, \\ \dot{Z} &= \frac{1}{b}(XY^* + X^*Y) - bZ. \end{aligned} \quad (2.24)$$

which have the fixed points $X = Y = Z = 0$ (unstable for $r_1 > r_{1c}$), and also

$$X = A; \quad Y = \left(1 + \frac{i\omega}{\sigma}\right) A; \quad Z = |A|^2/b \quad (2.25)$$

where $|A|$ is given by (2.21). Thus (2.24) does indeed have a continuum of equilibrium points (each corresponds to the limit cycle), as was the case for (1.1) when $\omega = 0$.

We observe that since (2.25) gives an equilibrium point independent of the phase, we may choose without any loss of generality that A is real. Now a steady state solution of (2.24) exists, in which A is replaced by $A \exp(i\epsilon) = A + i\epsilon A + O(\epsilon^2)$. It follows that the linearised equations about (2.25) must have a *neutrally stable solution* $(iX_0, iY_0, 0)$ (where 0 denotes the steady state) for all values of $r_1 > r_{1c}$. In other words, one eigenvalue of the matrix equation governing such perturbations will be zero. This physically signifies stability, since it is only a phase shift in the limit cycle, but ensures that the bifurcation does not satisfy the Hopf criterion in its entirety. The nature of this bifurcation and particularly the rôle of the $\lambda = 0$ eigenvalue is discussed in more detail in section 3. Furthermore, it is apparent that this discussion should in principle apply to any genuine limit cycle: examination of the present model thus gives an opportunity of explicit comparison of the theory with numerical experiment.

To analyse perturbations about the equilibrium point, we take A real (as above) and set

$$X = A + \xi,$$

$$Y = \left(1 + \frac{i\omega}{\sigma}\right) A + \eta,$$

$$Z = A^2/b + \zeta.$$

The linearised equations for ξ, η and ζ are then

$$\begin{aligned}
\dot{\xi} &= -(\sigma + i\omega)\xi + \sigma\eta, \\
\dot{\eta} &= (r - A^2/b)\xi - (a + i\omega)\eta - A\zeta, \\
\dot{\zeta} &= \frac{1}{2}A\left(1 - \frac{i\omega}{\sigma}\right)\xi + \frac{1}{2}\left(1 + \frac{i\omega}{\sigma}\right)A\xi^* \\
&\quad + \frac{1}{2}A\eta + \frac{1}{2}A\eta^* - b\zeta.
\end{aligned} \quad (2.26)$$

Because of the presence of ξ^* and η^* , these equations cannot immediately be written in matrix form, but must be supplemented by equations for ξ^* and η^* :

$$\begin{aligned}
\dot{\xi}^* &= -(\sigma - i\omega)\xi^* + \sigma\eta^*, \\
\dot{\eta}^* &= (r^* - A^2/b)\xi^* - (a^* - i\omega)\eta^* - A\zeta.
\end{aligned} \quad (2.27)$$

We define

$$\begin{aligned}
L &= a + i\omega, \\
N &= 1 + i\omega/\sigma, \\
P &= r - A^2/b,
\end{aligned} \quad (2.28)$$

and recalling (2.19), we observe that

$$LN = P. \quad (2.29)$$

(2.26) and (2.27) may now be written in the form

$$\begin{pmatrix} \dot{\xi} \\ \dot{\eta} \\ \dot{\xi}^* \\ \dot{\eta}^* \\ \dot{\zeta} \end{pmatrix} = \begin{pmatrix} -\sigma N & \sigma & 0 & 0 & 0 \\ P & -L & 0 & 0 & -A \\ 0 & 0 & -\sigma N^* & \sigma & 0 \\ 0 & 0 & P^* & -L^* & -A \\ \frac{1}{2}AN^* & \frac{1}{2}A & \frac{1}{2}AN & \frac{1}{2}A & -b \end{pmatrix} \begin{pmatrix} \xi \\ \eta \\ \xi^* \\ \eta^* \\ \zeta \end{pmatrix} \quad (2.30)$$

and as before, there exist solutions proportional to $\exp(\lambda t)$. At this point we observe that such solutions will be such that ξ^* is *not* the conjugate of ξ ; this is analogous to having complex

exponents λ in matrix equations governing real variables. In such cases the linearity of the equations enables suitable superpositions to be chosen so that the variables *are* real. Similarly, we would choose a superposition in the present case such that ξ^* is the conjugate of ξ .

This may be verified by writing (2.30) as a five by five *real* matrix equation for x_R, y_R , etc. Since the solutions proportional to $\exp(\lambda t)$ give complex conjugate pairs, or real values of λ , the same must be true of (2.30) (as is shown below). Thus solutions proportional to $\exp(\lambda t)$ exist when λ is an eigenvalue of (2.30); that is, when

$$\begin{vmatrix} -(\sigma N + \lambda) & \sigma & 0 & 0 & 0 \\ P & -(L + \lambda) & 0 & 0 & -A \\ 0 & 0 & -(\sigma N^* + \lambda) & \sigma & 0 \\ 0 & 0 & P^* & -(L^* + \lambda) & -A \\ \frac{1}{2}AN^* & \frac{1}{2}A & \frac{1}{2}AN & \frac{1}{2}A & -(b + \lambda) \end{vmatrix} = 0. \quad (2.31)$$

This is easily evaluated by multiplying the elements of the fifth column by its cofactors. We obtain

$$\begin{aligned}
0 &= (b + \lambda)[(\sigma N + \lambda)(L + \lambda) - \sigma P] \\
&\quad \times [(\sigma N^* + \lambda)(L^* + \lambda) - \sigma P^*] \\
&\quad + \frac{1}{2}A^2(\sigma N^* + \lambda + \sigma N)[(\sigma N + \lambda)(L + \lambda) \\
&\quad - \sigma P + (\sigma N^* + \lambda)(L^* + \lambda) - \sigma P^*],
\end{aligned} \quad (2.32)$$

Using $LN = P$ from (2.29), we find that λ is a factor (as predicted): thus $\lambda = 0$, or

$$\begin{aligned}
&\lambda(\lambda + b)[\lambda + (L + N\sigma)][\lambda + (L^* + N^*\sigma)] \\
&\quad + \frac{1}{2}A^2(\lambda + \sigma N + \sigma N^*) \\
&\quad \times [(\lambda + L + N\sigma) + (\lambda + L^* + N^*\sigma)] = 0,
\end{aligned} \quad (2.33)$$

which is a quartic polynomial in λ . We evaluate an explicit stability criterion as follows. Define

$$\begin{aligned}
\alpha &= (L + \sigma N)(L^* + \sigma N^*) = (\sigma + 1)^2 + (2\omega - e)^2, \\
\beta &= \frac{1}{2}[L + \sigma N + L^* + \sigma N^*] = \sigma + 1, \\
\gamma &= \sigma(N + N^*) = 2\sigma.
\end{aligned} \quad (2.34)$$

Eq. (2.33) is thus

$$\lambda(\lambda + b)(\lambda^2 + 2\beta\lambda + \alpha) + A^2(\lambda + \gamma)(\lambda + \beta) = 0. \quad (2.35)$$

We have $\alpha, \beta, \gamma > 0$; thus the constant term is positive for all $A^2 > 0$. Hence $\lambda = 0$ is not a solution of (2.35) and instability can only occur if conjugate roots of (2.35) cross the imaginary axis. For $A^2 > 0$, $A^2 \rightarrow 0$, the roots of (2.35) all have negative real part. Therefore the limit cycle is stable for $r_1 > r_{lc}$ until a critical value $r_1 = r'_{lc}$, at which (2.35) has roots $\lambda = \pm i\Omega$, say. At this point (if it exists), (2.35) may be written

$$(\lambda^2 + \Omega^2)(\lambda^2 + p\lambda + q) = 0. \quad (2.36)$$

Multiplying out (2.35) and (2.36), we find, on identifying coefficients of powers of λ , that we must have

$$\begin{aligned} p &= b + 2\beta, \\ \Omega^2 + q &= \alpha + 2b\beta + A^2, \\ p\Omega^2 &= \alpha b + (\beta + \gamma)A^2, \\ q\Omega^2 &= \beta\gamma A^2. \end{aligned} \quad (2.37)$$

Eliminating p, q and Ω^2 gives a critical value of A^2 , and hence of r_1 , at which instability sets in. Since α, b, β and γ are all positive, p, q , and Ω are also, and thus (2.36) implies that Ω is real, and the other two roots of (2.36) have negative real parts. From (2.37), we have

$$\Omega^2 = [\alpha b + (\beta + \gamma)A^2]/(b + 2\beta), \quad (2.38)$$

whence

$$q = \frac{\beta\gamma A^2(b + 2\beta)}{[\alpha b + (\beta + \gamma)A^2]}, \quad (2.39)$$

so that

$$\begin{aligned} \alpha + 2b\beta + A^2 &= \frac{\alpha b + (\beta + \gamma)A^2}{(b + 2\beta)} \\ &+ \frac{\beta\gamma A^2(b + 2\beta)}{[\alpha b + (\beta + \gamma)A^2]} \end{aligned} \quad (2.40)$$

is the stability criterion for A^2 . This is evidently a quadratic equation for A^2 ,

$$Q(A^2) \equiv Q_1 A^4 + Q_2 A^2 + Q_3 = 0, \quad (2.41)$$

where some simplification shows that

$$\begin{aligned} Q_1 &= (\beta + \gamma)(\gamma - b - \beta) \\ &= (3\sigma + 1)(\sigma - b - 1), \\ Q_2 &= \beta(b + 2\beta)(2\beta\gamma - 2b\beta - b\gamma) \\ &\quad - \alpha(-b\gamma + b^2 + \beta b + 2\beta^2 + 2\beta\gamma), \\ Q_3 &= -2\alpha\beta b[\alpha + b^2 + 2\beta b]. \end{aligned} \quad (2.42)$$

We have $Q_3 < 0$. If $\sigma > b + 1$, then $Q_1 > 0$ and (2.41) has a unique positive root, which determines the critical value of A^2 . This turns out to be

$$\rho = \frac{-Q_2 + [Q_2^2 - 4Q_1Q_3]^{1/2}}{2Q_1}. \quad (2.43)$$

In this case the critical value of r_1 where the limit cycle becomes unstable is given by

$$r'_{lc} = r_{lc} + \rho/b, \quad (2.44)$$

where ρ is given by (2.43), Q_1, Q_2 and Q_3 by (2.42), α, β and γ by (2.34) and ω by (2.20). This is a useable, though messy criterion.

If $\sigma < b + 1$, then $Q_1 < 0$ and Q may attain positive values for a finite range of A^2 . A necessary condition that this occurs is that $Q_2 > 0$. Using the definitions of β and γ in (2.34), together with the fact that $\alpha \geq \beta^2$ and $b > \sigma - 1$, evaluation of Q_2 in (2.42) shows that necessarily $Q_2 < 0$. It follows that the limit cycle is always stable if $\sigma < b + 1$, which is identical to the corresponding real case.

To summarise, we have shown that the limit cycle, in the form of the exact solution (2.22), is always stable if $\sigma < b + 1$. If $\sigma > b + 1$ then it is also stable for values of r_1 in the range $r_{lc} < r_1 < r'_{lc}$ but becomes unstable when $r_1 > r'_{lc}$. Due to the presence of the zero eigenvalue of the linearised matrix equation the bifurcation is not

completely of Hopf type although without this eigenvalue a Hopf bifurcation would occur. The nature of this bifurcation thus requires further consideration.

3. Bifurcation of the limit cycle in the complex Lorenz equations

3.1. General case

Let us first consider the general case of bifurcation of a limit cycle. Straightforward linearised perturbation analysis may be carried out, leading to a linear matrix equation for the perturbations in which the matrix is periodic of period T , say (where T is the period of the underlying limit cycle). Floquet theory (see e.g., Coddington and Levinson [7]) then tells us that the solutions of this linear equation may be written as the product of a function of period T and the function $\exp(\mu t)$. Stability then rests on the magnitude of the Floquet multipliers $\exp(\mu T)$, and thus on the nature of the Floquet exponents μ . In the case of our analysis of the complex Lorenz limit cycle, the exponents μ are essentially the same as the eigenvalues λ .

Various possibilities for bifurcation now occur, and are discussed by Ruelle and Takens [8], Joseph [9] and Lanford [10]. Particularly, if λ crosses the imaginary axis at $i\Omega$ (and so also λ^* at $-i\Omega$), then the limit cycle bifurcates to motion on a 2-torus. Generally this motion will be doubly periodic (corresponding to a "Hopf bifurcation" in a frame rotating with the limit cycle, or of the associated Poincaré map); for particular values of λ , subharmonic periodic solutions may also occur (Joseph [9]).

However, Floquet theory is not generally useful for explicit calculations, and in addition the limit cycle solution is not usually known explicitly. Thus practical applications of such theorems as exist do not seem prevalent in the literature: particularly, we are unaware of a method for computation of the stability of the

bifurcating torus. In the Hopf case, there are numerous "different" methods which can be used, which probably amount to the same idea: the center manifold theorem, Hopf's theorem, the Poincaré-Lindstedt method, the Krylov-Bogoliubov-Mitropolsky method of averaging, and the Cole-Kevorkian method of multiple scales, as originally developed for fluid flows by Stuart [11]. The last-named method derives an amplitude equation for marginally stable oscillatory perturbations of the form

$$\dot{x} = x_0 + [\epsilon A(\tilde{t}) e^{i\Omega \tilde{t}} u_0 + \text{c.c.}] + \dots, \quad (3.1)$$

wherein ϵ is a measure of the amplitude A ; $\tilde{t} = \epsilon^2 t$ is a slow time variable, and Ω is the marginal frequency. Uniformly valid expansions of the form (3.1) require A to satisfy an equation of the form

$$\frac{dA}{d\tilde{t}} = k_1 A + k_2 A |A|^2, \quad (3.2)$$

a result obtained by McLaughlin and Martin [6] in their classification of the bifurcations in the real Lorenz model. Along with the various different methods mentioned above, our calculation is essentially equivalent to theirs.

If $\text{Re } k_2 > 0$, then the bifurcation is subcritical, and the limit cycle is unstable; if $\text{Re } k_2 < 0$, the bifurcation is supercritical and stable. Calculation of k_2 is straightforward but messy.

The distinction between super- and sub-critical bifurcations is an important one in the context of turbulence and chaotic trajectories of differential equations. Ruelle and Takens [8] proposed that the trajectories on higher dimensional tori (corresponding to further bifurcations of the system under consideration) would not generally be of periodic or almost periodic type; rather, they could approach "strange" attractors, which for all practical purposes would appear turbulent, or chaotic. It is clear that this scheme is only viable if the intermediate bifurcations are supercritical, so that the trajectories

on the various tori are stable (in that they approach the torus). It is for this reason that it is useful to analyse the direction of bifurcation in real cases. In fluid flow along pipes, bifurcation (when it occurs) is of subcritical type, and so the Ruelle-Takens ideas are presumably irrelevant. In the Bénard problem, the first bifurcation, and in the Taylor column, the first two bifurcations are supercritical and stable, and turbulence in these cases may be along the lines of the abstract theory.

For the above reasons, we wish to examine the direction of bifurcation of the 2-torus for the complex Lorenz equations. Our approach will be constructive; that is, we will use the formal method of multiple scales to find amplitude equations satisfied by the perturbations of the limit cycle. An analysis of these gives conditions of stability and direction of bifurcation of the 2-torus. Since the bifurcation is subcritical in the real case (McLaughlin and Martin [6]), we expect this to be also true in the present case, at least for sufficiently small ϵ and r_2 .

3.2. Rotationally invariant systems

The procedure we adopt is straightforward. A similar general method is given by Haken [12], who follows somewhat the approach of Eckhaus [13]. This determines an infinity of amplitude functions, of which only a finite set are relevant; an unstable mode and a set of "slaved" modes, in Haken's terminology. We prefer to adopt the method of multiple scales (e.g., Nayfeh [14]), since then the approximate expansions adopted are made explicit from the outset, and the nature of the equations is then apparent.

Let us consider the system of real-valued ordinary differential equations

$$\frac{dx_i}{dt} = f_i(x; \mu). \quad (3.3)$$

We shall suppose that this system is derived via a change of variables from another system, in

which a limit cycle exists for a certain range of μ ; we shall refer to this as the underlying limit cycle. For the complex Lorenz equations, we obtain (3.3) via the change of variables $(x, y, z) \rightarrow (xe^{i\omega t}, ye^{i\omega t}, z)$; generally, we can change into a rotating frame in this way for any system with an underlying limit cycle, but we expect to obtain a non-autonomous system unless the oscillation is exactly sinusoidal. In this case, it is reasonable to adopt the following assumptions. Let x_0 be a fixed point of (3.3) corresponding to an underlying limit cycle. We define a rotation matrix $R(t)$ which satisfies $R(\alpha)R(\beta) = R(\alpha + \beta)$, and the underlying limit cycle is given by $R(t)x_0$ (i.e., (3.3) is obtained from the original system via the change of variables $x \rightarrow Rx$). Since then we have that

$$f[Rx_0; \mu] = 0, \quad R = R(\alpha), \quad (3.4)$$

for all μ and α , (i.e., (3.3) has a continuum of equilibria), it is reasonable to assume that (3.3) is invariant under rotation; that is, if $y = R(\alpha)x$, then $dy/dt = f(y; \mu)$. This implies that f satisfies

$$Rf(x; \mu) = f(Rx; \mu) \quad (3.5)$$

for all x and α . This condition is satisfied by the complex Lorenz equations, for example (see below). Differentiation of (3.5) with respect to α yields

$$R'_{ij}(\alpha)f_j(x; \mu) = f_{i,j}(Rx; \mu)R'_{jk}(\alpha)x_k; \quad (3.6)$$

putting $x = x_0$ immediately shows, using (3.5), that

$$f_{i,j}(Rx_0; \mu)R'_{jk}x_{0k} = 0. \quad (3.7)$$

In other words the Jacobian matrix $Df = (f_{i,j}(Rx_0; \mu))$ has, for every value of μ (and α), a right eigenvector u_0 with corresponding eigenvalue zero, where u_0 is given by

$$u_0 = R'(\alpha)x_0. \quad (3.8)$$

Since linear perturbations to x_0 are proportional to $\exp(\lambda t)$, where λ is an eigenvalue of Df , it follows that x_0 is (at least) neutrally stable for all μ . However, this is by virtue of the fact that the underlying limit cycle may have its phase perturbed, but yet be arbitrarily stable; thus we discount $\lambda = 0$ as representing a state of marginal stability, since if all other $\text{Re } \lambda$ are less than zero, then the underlying limit cycle is structurally stable.

If we differentiate (3.5) with respect to x , we find

$$R(Df(x; \mu)) = (Df(Rx; \mu))R. \quad (3.8a)$$

Now suppose that U is a right eigenvector of $Df(x; \mu)$ with eigenvalue λ : then pre-multiplying U by (3.8a) shows that RU is the corresponding eigenvector of $Df(Rx; \mu)$. The use of this is that U is independent of α in $R(\alpha)$. Particularly, (3.8) implies (with an obvious notation)

$$U_0 = R'(0)x_0, \quad (3.8b)$$

whence $u_0 = R(\alpha)R'(0)x_0$, which is consistent with (3.8), since consideration of $R(\alpha + \beta) = R(\alpha)R(\beta)$ shows that $R'(\alpha) = R(\alpha)R'(0) = R'(0)R(\alpha)$.

Let us denote $f_{i,j}(x_0; \mu)$ by $f_{i,j}^0$; we will now assume that (apart from the zero eigenvalue), all other eigenvalues of $f_{i,j}^0$ are such that $\text{Re}(\lambda) < 0$, but that at $\mu = \mu_c$, a pair of eigenvalues $\pm i\Omega$, $\Omega > 0$, cross the imaginary axis in such a way that $d(\text{Re } \lambda)/d\mu|_{\mu_c} > 0$: the corresponding (critical) eigenvectors of $f_{i,j}^0$ are U_n and U_n^* . These conditions (apart from the zero eigenvalue) resemble those of the Hopf bifurcation, and are valid in the complex Lorenz case. From the paragraph above, it follows that $f_{i,j}(Rx_0; \mu)$ has corresponding eigenvectors $u_n = RU_n$ and $u_n^* = RU_n^*$ with eigenvalues $+i\Omega$ and $-i\Omega$, respectively.

We now seek an approximate solution when $|\mu - \mu_c| \ll 1$; accordingly we define (with some

foresight)

$$\mu = \mu_c \pm \epsilon^2, \quad 0 < \epsilon \ll 1, \quad (3.9)$$

where the plus and minus signs refer respectively to super- and sub-critical states (weakly unstable and weakly stable). We also define the slow time scale

$$\tau = \epsilon^2 t, \quad (3.10)$$

and will seek solutions to (3.3) in the form

$$x = x^{(0)} + \epsilon x^{(1)} + \epsilon^2 x^{(2)} + \dots \quad (3.11)$$

where $x^{(i)} = x^{(i)}(t, \tau)$. The procedure as usual is straightforward, but there are one or two subtleties which distinguish the expansion from the more conventional Hopf case. Substituting (3.9), (3.10) and (3.11) into (3.3), we obtain

$$\begin{aligned} \left(\frac{\partial}{\partial t} + \epsilon^2 \frac{\partial}{\partial \tau} \right) (x^{(0)} + \epsilon x^{(1)} + \epsilon^2 x^{(2)} + \epsilon^3 x^{(3)} + \dots) \\ = f[x^{(0)} + \epsilon x^{(1)} + \dots; \mu_c \pm \epsilon^2], \end{aligned} \quad (3.12)$$

whence we derive

$$\begin{aligned} \left[\frac{\partial x_1^{(0)}}{\partial t} + \epsilon \frac{\partial x_1^{(1)}}{\partial t} + \epsilon^2 \frac{\partial x_1^{(2)}}{\partial t} + \epsilon^3 \frac{\partial x_1^{(3)}}{\partial t} + \dots \right] \\ + \left[\epsilon^2 \frac{\partial x_1^{(0)}}{\partial \tau} + \epsilon^3 \frac{\partial x_1^{(1)}}{\partial \tau} + \dots \right] \\ = [f_1^c + \{\epsilon x_1^{(1)} + \epsilon^2 x_1^{(2)} + \epsilon^3 x_1^{(3)} + \dots\} f_{1,j}^c] \\ + \{\epsilon x_1^{(1)} + \epsilon^2 x_1^{(2)} + \dots\} \{\epsilon x_1^{(1)} + \epsilon^2 x_1^{(2)} + \dots\} f_{1,k}^c \\ + \{\epsilon x_1^{(1)} + \epsilon^2 x_1^{(2)} + \dots\} \{\epsilon x_1^{(1)} + \epsilon^2 x_1^{(2)} + \dots\} f_{1,\mu}^c + \dots] \\ \pm \epsilon^2 \left[\frac{\partial f_1^c}{\partial \mu} + \{\epsilon x_1^{(1)} + \dots\} \frac{\partial f_{1,j}^c}{\partial \mu} + \dots \right] + \dots, \end{aligned} \quad (3.13)$$

where $f_{i,j}^c$ denotes $f_{i,j}$ evaluated at $x^{(0)}$ and μ_c , etc. Equating terms of $O(1)$, we have

$$\frac{\partial x_1^{(0)}}{\partial t} = f_1(x^{(0)}; \mu_c). \quad (3.14)$$

The relevant t -independent solution of (3.14) is given, from (3.4), by

$$\mathbf{x}^{(0)} = R(\phi)\mathbf{x}_0, \quad (3.15)$$

where $\phi = \phi(\tau)$ is a slowly-varying function of time; essentially, it is the phase shift in the underlying limit cycle. At $\mathcal{O}(\epsilon)$, we have the linearised equation

$$\frac{\partial \mathbf{x}_i^{(1)}}{\partial t} - f_{i,j}^c \mathbf{x}_j^{(1)} = 0, \quad f_{i,j}^c = f_{i,j}[R(\phi)\mathbf{x}_0; \mu_c], \quad (3.16)$$

Neglecting initial transients, this equation has solutions proportional to u_0 , $u_\Omega e^{i\Omega t}$ and $u_\Omega^* e^{-i\Omega t}$; however, notice that if, in (3.15) $\phi = \phi_0 + \epsilon\phi_1 + \dots$, then

$$\mathbf{x}^{(0)} = R(\phi_0)\mathbf{x}_0 + \epsilon iR'(\phi_0)\mathbf{x}_0\phi_1 + \dots$$

Since $u_0 = R'(\phi_0)\mathbf{x}_0$, it is clear that we may absorb the term proportional to u_0 in $\mathbf{x}^{(1)}$ into ϕ . In fact we can do this at each stage of the solution, provided we let ϕ depend on ϵ as well as τ . Then the solution of (3.16), neglecting transients, may be written

$$\mathbf{x}^{(1)} = A(\tau)u_\Omega e^{i\Omega t} + (*), \quad (3.17)$$

where $(*)$ denotes the complex conjugate.

At $\mathcal{O}(\epsilon^2)$, we have, since $\partial f_i^c / \partial \mu = 0$ (from (3.4)),

$$\begin{aligned} \frac{\partial \mathbf{x}_i^{(2)}}{\partial t} - f_{i,j}^c \mathbf{x}_j^{(2)} &= -\frac{\partial \mathbf{x}_i^{(0)}}{\partial \tau} + \frac{1}{2} \mathbf{x}_k^{(1)} \mathbf{x}_l^{(1)} f_{i,jk}^c \\ &= -R'_{ij}(\phi) \frac{d\phi}{d\tau} x_{0j} + f_{i,jk}^c |A|^2 u_{\Omega j} u_{\Omega k}^* \\ &\quad + \left[\frac{1}{2} f_{i,jk}^c A^2 e^{2i\Omega t} u_{\Omega j} u_{\Omega k} + (*) \right], \end{aligned} \quad (3.18)$$

since $f_{i,jk} = f_{i,jk}$. The term in $e^{2i\Omega t}$ gives a particular solution proportional to $e^{2i\Omega t}$; however (since zero is an eigenvalue of $f_{i,j}^c$) the constant terms may be secular, and it is the elimination of these secular terms which determines the phase $\phi(\tau)$. Specifically, if a (complex-valued)

matrix A has an eigenvalue zero, then the equation $A\mathbf{x} = \mathbf{c}$ has a solution if and only if $\bar{v}^T \mathbf{c} = 0$, where a bar denotes the complex conjugate, for all v such that $\bar{A}^T v = 0$. Thus if λ is an eigenvalue of $Df = (f_{i,j}^c)$, then the equation

$$\frac{\partial \mathbf{x}}{\partial t} - (Df)\mathbf{x} = \mathbf{c} e^{\lambda t} \quad (3.19)$$

has a solution proportional to $e^{\lambda t}$ if and only if $v^T \mathbf{c} = 0$ where $[(Df) - \lambda I]^T \bar{v} = 0$: i.e. $(Df)^T v^* = \lambda^* v^*$ ($*$ denotes complex conjugate) since Df is real. Now let v_0 and v_Ω be the left eigenvectors of Df , corresponding to the eigenvalues zero and $i\Omega$ respectively, thus

$$\begin{aligned} v_0^T (Df) &= 0, \quad \text{i.e. } v_{0j} f_{i,j}^c = 0, \\ v_\Omega^T (Df) &= i\Omega v_\Omega^T, \quad \text{i.e. } v_{\Omega j} f_{i,j}^c = i\Omega v_{\Omega i}; \end{aligned} \quad (3.20)$$

then $(Df)^T v_0^* = 0$, $(Df)^T v_\Omega^* = -i\Omega v_\Omega^*$, and thus it follows that the equation

$$\frac{\partial \mathbf{x}}{\partial t} - (Df)\mathbf{x} = \mathbf{c}_0 + \mathbf{c}_\Omega e^{i\Omega t} \quad (3.21)$$

can only have a solution of the form $\mathbf{a}_0 + \mathbf{a}_\Omega e^{i\Omega t}$ if the constraints

$$v_0 \cdot \mathbf{c}_0 = v_\Omega \cdot \mathbf{c}_\Omega = 0 \quad (3.22)$$

are satisfied; otherwise, secular terms t , $te^{i\Omega t}$ will occur. Applying this to (3.18), we require ϕ to satisfy, recalling (3.8),

$$\frac{d\phi}{d\tau} = \left[\frac{v_{0j} f_{i,jk}^c u_{\Omega j} u_{\Omega k}^*}{u_{0i} v_{0i}} \right] |A|^2. \quad (3.23)$$

If (3.23) is satisfied, then a particular solution of (3.18) can be written

$$\mathbf{x}^{(2)} = \mathbf{a}^{(2)} |A|^2 + \left[\mathbf{a}^{(2)} A^2 e^{2i\Omega t} + (*) \right], \quad (3.24)$$

where

$$f_{i,j}^c \mathbf{a}_0^{(2)} = \left[\frac{v_{0j} f_{i,jk}^c u_{\Omega j} u_{\Omega k}^*}{u_{0k} v_{0k}} \right] u_{0i} - f_{i,jk}^c u_{\Omega j} u_{\Omega k}^*, \quad (3.25)$$

and

$$2i\Omega a_{ij}^{(2)} - f_{i,j}^c a_{ij}^{(2)} = \frac{1}{2} f_{i,k}^c u_{nj} u_{nk}. \quad (3.24)$$

We have already decided to incorporate constant solutions proportional to u_0 into ϕ ; similarly, there is no loss in generality in absorbing the solutions proportional to $u_0 e^{i\Omega t}$ into $A(\tau)$, provided also we understand $A = A(\tau; \epsilon)$. Then (3.24) is the complete solution at $O(\epsilon^2)$, and we turn to $O(\epsilon^3)$ terms. These are

$$\begin{aligned} \frac{\partial x_j^{(3)}}{\partial \epsilon} - f_{i,j}^c x_j^{(3)} = & -\frac{\partial x_i^{(1)}}{\partial \tau} + \frac{1}{2} \{x_j^{(1)} x_k^{(2)} + x_j^{(2)} x_k^{(1)}\} f_{i,k}^c \\ & + \frac{1}{6} x_j^{(1)} x_k^{(1)} x_l^{(1)} f_{i,kl}^c \pm x_j^{(1)} \frac{\partial f_{i,j}^c}{\partial \mu}, \end{aligned} \quad (3.27)$$

and we again have to eliminate terms on the right-hand side which are secular. From (3.17) and (3.24), it is apparent that there are no constant secular terms, and thus we only need choose A such that the $e^{i\Omega t}$ terms are non-secular; we in fact obtain the Landau–Stuart equation (3.2), as previously discussed. The coefficients are determined as follows. The coefficient of $e^{i\Omega t}$ in (3.27) is, recalling that $u_0 = R(\phi)U_0$ and that $R'(\phi) = R'(0)R(\phi)$,

$$\begin{aligned} -\frac{dA}{d\tau} u_{0i} \pm A \frac{\partial f_{ij}^c}{\partial \mu} u_{nj} - AR'_{ij}(0) u_{nj} \frac{d\phi}{d\tau} \\ + \frac{1}{2} f_{i,k}^c \{u_{nj} a_{jk}^{(2)} + u_{nj}^* a_{jk}^{(2)} + u_{nk} a_{ij}^{(2)} + u_{nk}^* a_{ij}^{(2)}\} |A|^2 A \\ + \frac{1}{6} \{u_{nj} u_{nk} u_{nl}^* + u_{nj} u_{nk}^* u_{nl} + u_{nj}^* u_{nk} u_{nl}\} f_{i,kl}^c |A|^2 A, \end{aligned}$$

so it follows from (3.22), using (3.23), that the Landau equation for A is

$$\frac{dA}{d\tau} = \pm k_1 A + k_2 |A|^2 A, \quad (3.28)$$

where

$$k_1 = v_{0i} \frac{\partial f_{ij}^c}{\partial \mu} u_{nj} / u_{0i} v_{0i}, \quad (3.29)$$

$$\begin{aligned} k_2 = & [v_{0i} f_{i,k}^c \{u_{nj} a_{jk}^{(2)} + u_{nj}^* a_{jk}^{(2)}\} - \beta v_{0i} R'_{ij}(0) u_{nj} \\ & + \frac{1}{6} v_{0i} \{u_{nj} u_{nk} u_{nl}^* + u_{nj} u_{nk}^* u_{nl} \\ & + u_{nj}^* u_{nk} u_{nl}\} f_{i,kl}^c] / v_{0i} u_{0i}, \end{aligned} \quad (3.30a)$$

$$\beta = \frac{v_{0i} f_{i,k}^c u_{nj} u_{nk}^*}{u_{0i} v_{0i}}, \quad (3.30b)$$

$k_1 = d\lambda/d\mu|_{\mu=\mu_c}$ is the linear growth rate ($\text{Re } k_1 > 0$ by assumption). Therefore, the bifurcation is supercritical if $\text{Re } k_2 < 0$, and subcritical if $\text{Re } k_2 > 0$ since $|A|^2$ satisfies

$$\frac{d|A|^2}{d\tau} = \pm 2(\text{Re } k_1)|A|^2 + 2(\text{Re } k_2)|A|^4. \quad (3.31)$$

Thus the stability of the bifurcating solutions is determined in exactly the same manner as in a Hopf bifurcation. It is not obvious (though we might suspect) that β , k_1 and k_2 are independent of τ , since u_0 , u_0 , v_0 and v_0 are functions of τ . However, using $u_0 = RU_0$, $u_0 = RU_0$, and the equivalent formulae for v_0 and v_0 , it is not difficult to check, using x -differentials of (3.5), that this is indeed the case. For example, pre-multiplication of (3.8a) by v , a left eigenvector of $Df(Rx; \mu)$, easily shows (if V is the corresponding eigenvector of $Df(x; \mu)$) that $V^T = v^T R$. It easily follows that we have $v_0^T = V_0^T R^{-1}$, $v_0^T = V_0^T R^{-1}$. We then have $v_{0i} u_{0i} = v_0^T u_0 = V_0^T R^{-1} R U_0 = V_0^T U_0$, which is independent of ϕ , and hence of τ . Similar considerations show that β , k_1 and k_2 are all independent of τ .

3.3. Nature of the solutions

The solution can be written as

$$x = R[\phi(\tau)]x_0 + \epsilon[A(\tau)u_0 e^{i\Omega t} + (*)] + O(\epsilon^2), \quad (3.32)$$

where

$$\frac{d\phi}{d\tau} = \beta |A|^2, \quad \frac{dA}{d\tau} = k_1 A + k_2 |A|^2 A. \quad (3.33)$$

If the bifurcation is supercritical, then for $\text{Re } k_1 > 0$, i.e., $\mu > \mu_c$, the solution for x tends to

a doubly periodic motion in the rotating plane. This consists of a fast ($t \sim 1$) oscillation of amplitude order $\sqrt{\mu - \mu_c}$ about the equilibrium point, which itself precesses periodically around the underlying limit cycle. In the rotating plane, time-plots of x_r versus time will thus consist of a fast oscillation which (if $u_{0i} \neq 0$) is superimposed by a larger scale slowly oscillating solution. This is observed in the complex Lorenz equations (see fig. 2).

In the fixed frame, we have $x_F = Rx$, therefore (3.32) is, using the property that $R(\alpha)R(\beta) = R(\alpha + \beta)$, and that $u_n = RU_n$,

$$x_F = R(t + \phi(\tau))[x_0 + \epsilon\{A(\tau)U_{0i}e^{i\Omega t} + (*)\} + \mathcal{O}(\epsilon^2)]. \quad (3.34)$$

In the limit as $\tau \rightarrow \infty$ (with $\text{Re } k_2 < 0$, $\text{Re } k_1 > 0$), we have

$$|A|^2 \rightarrow -\left(\frac{\text{Re } k_1}{\text{Re } k_2}\right) = -\frac{k_{1R}}{k_{2R}}, \quad (3.35)$$

thus

$$\begin{aligned} \phi &\sim -\frac{\beta k_{1R}}{k_{2R}}\tau, \\ A &\sim \exp\left[\left(k_1 - k_2 \frac{k_{1R}}{k_{2R}}\right)\tau\right] \\ &= \exp\left[i\left\{\frac{k_{1I}k_{2R} - k_{2I}k_{1R}}{k_{2R}}\right\}\tau\right]. \end{aligned} \quad (3.36)$$

It follows that the solutions in the fixed frame can be written in component form as the sum of products of two functions of distinct periods, thus

$$\begin{aligned} x_{Fi} &= R_{ij}[t + \phi(\tau)]x_{0j} \\ &\quad + \epsilon\{A(\tau)R_{ij}(t + \phi(\tau))U_{0j}e^{i\Omega t} + (*)\} + \mathcal{O}(\epsilon^2) \\ &= \sum_j p_j^{(i)}(t)q_j(t), \end{aligned} \quad (3.37)$$

where

$$\begin{aligned} p_j^{(i)}(t) &= R_{ij}(t + \phi(\tau)), \\ q_j(t) &= x_{0j} + \epsilon\{A(\tau)U_{0j}e^{i\Omega t} + (*)\} + \mathcal{O}(\epsilon^2), \end{aligned}$$




Fig. 2. Plot of y_R versus time at $b = 0.8$, $\sigma = 2$, $r_1 = 220$, $r_2 = 1$ and $\epsilon = 3r_2$, at which there is a stable finite amplitude motion. Note the slow and fast periods in the oscillations.

and $p_j^{(n)}$ and q_j have periods ω_1 and ω_2 respectively, where

$$\begin{aligned}\omega_1 &= \omega \left[1 - \epsilon^2 \frac{\beta k_{1R}}{k_{2R}} \dots \right], \\ \omega_2 &= \Omega + \epsilon^2 \left\{ \frac{k_{11}k_{2R} - k_{21}k_{1R}}{l_{2R}} \right\} \dots\end{aligned}\quad (3.38)$$

The presence of two distinct, generally incommensurate frequencies characterise these solutions as doubly periodic. A spectral analysis of the solutions should thus reveal a peak at the fundamental frequency ω_1 , together with smaller peaks, at $|\omega_1 \pm \omega_2|$, etc. Such behaviour is found by Fenstermacher, Swinney and Gollub [17] in an experimental analysis of Taylor vortex flow.

3.4. Application to complex Lorenz equations

In the rotating plane, the complex Lorenz equations are

$$\begin{aligned}\dot{X} &= -(\sigma + i\omega)X + \sigma Y, \\ \dot{Y} &= (r - Z)X - (a + i\omega)Y, \\ \dot{Z} &= \frac{1}{b}[XY^* + X^*Y] - bZ,\end{aligned}\quad (3.39)$$

where

$$\omega = \frac{\sigma(e + r_2)}{\sigma + 1}, \quad (3.40)$$

The non-zero equilibria are

$$X = A, \quad Y = A \left(1 + \frac{i\omega}{\sigma} \right), \quad Z = |A|^2/b, \quad (3.41)$$

$$|A|^2 = b(r_1 - r_{1c}), \quad (3.42)$$

Note that the equations are invariant under the rotation $(X, Y, Z) \rightarrow (X \exp(i\alpha), Y \exp(i\alpha), Z)$. Putting

$$X = A(x_1 + ix_2), \quad Y = A(x_3 + ix_4), \quad Z = \frac{|A|^2}{b}x_5, \quad (3.43)$$

we get

$$\begin{aligned}\dot{x}_1 + ix_2 &= -(\sigma + i\omega)(x_1 + ix_2) + \sigma(x_3 + ix_4), \\ \dot{x}_3 + ix_4 &= [r_1 + ir_2 - (r_1 - r_{1c})x_5](x_1 + ix_2) \\ &\quad - (1 - ie + i\omega)(x_3 + ix_4), \\ \dot{x}_5 &= b \left[\frac{1}{2} \{ (x_1 + ix_2)(x_3 - ix_4) \right. \\ &\quad \left. + (x_1 - ix_2)(x_3 + ix_4) \} - x_5 \right],\end{aligned}$$

whence

$$\begin{aligned}\dot{x}_1 &= -\sigma x_1 + \omega x_2 + \sigma x_3, \\ \dot{x}_2 &= -\omega x_1 + \sigma x_2 + \sigma x_4, \\ \dot{x}_3 &= [r_1 - (r_1 - r_{1c})x_5]x_1 - r_2 x_2 - x_3 + (\omega - e)x_4, \\ \dot{x}_4 &= r_2 x_1 + [r_1 - (r_1 - r_{1c})x_5]x_2 - (\omega - e)x_3 - x_4, \\ \dot{x}_5 &= b[x_1 x_3 + x_2 x_4 - x_5],\end{aligned}\quad (3.44)$$

with an equilibrium point

$$x_1 = 1, \quad x_2 = 0, \quad x_3 = 1, \quad x_4 = \frac{\omega}{\sigma}, \quad x_5 = 1. \quad (3.45)$$

By rotation through an angle α of X and Y , other equilibria are given by

$$\begin{aligned}x_1 &= \cos \alpha, \\ x_2 &= \sin \alpha, \\ x_3 &= \cos \alpha - \frac{\omega}{\sigma} \sin \alpha, \\ x_4 &= \frac{\omega}{\sigma} \cos \alpha + \sin \alpha, \\ x_5 &= 1.\end{aligned}\quad (3.46)$$

The transformation $X \rightarrow X \exp(i\alpha)$, $Y \rightarrow Y \exp(i\alpha)$, $Z \rightarrow Z$ corresponds to $x_1 \rightarrow x_1 \cos \alpha - x_2 \sin \alpha$, $x_2 \rightarrow x_1 \sin \alpha + x_2 \cos \alpha$, $x_3 \rightarrow x_3 \cos \alpha - x_4 \sin \alpha$, $x_4 \rightarrow x_3 \sin \alpha + x_4 \cos \alpha$, $x_5 \rightarrow x_5$; thus the rotation matrix in this case can be written down as

$$R(\alpha) = \begin{pmatrix} \cos \alpha & -\sin \alpha & 0 & 0 & 0 \\ \sin \alpha & \cos \alpha & 0 & 0 & 0 \\ 0 & 0 & \cos \alpha & -\sin \alpha & 0 \\ 0 & 0 & \sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix} \quad (3.47)$$

and the methods previously described can be used to study the bifurcation of the limit cycle solution of (3.44): the algebra involved is somewhat tedious, and is not pursued here. However, we can already draw some conclusions from (3.47), (3.32) and (3.37).

In the rotating frame, the components of x will bifurcate to a small amplitude oscillation about a large scale oscillation of much smaller frequency. This follows from (3.32), and is essentially observed in fig. 2. In the fixed frame, we will see the same picture, but with the discrepancy in frequencies less visible. However, note that since, from (3.47), we must have $R_{5j}x_{0j} = x_{05}$, and similarly for U_0 , the z -component of x is given in either fixed or rotating frame by

$$Z = x_5 = x_{05} + \epsilon \{A(\tau) e^{i\Omega\tau} + (*)\} + \mathcal{O}(\epsilon^2); \quad (3.48)$$

therefore, the solution for z is singly periodic at bifurcation, and no slow variation of large amplitude should exist: this is seen in fig. 6.

4. Numerical results and conclusions

In the previous sections, we have shown that, except for the singular case $e + r_2 = 0$, only the origin is a fixed point. The two further fixed points of the real Lorenz equations which exist when $r > 1$ are replaced by the limit cycle which has frequency ω . Much of this paper has been devoted to the stability of this limit cycle and the nature of the bifurcation when it becomes unstable. The analysis of section 3 showed that the sign of $\text{Re}(k_2)$ effectively determines whether this bifurcation is of a subcritical

($\text{Re}(k_2) > 0$) or supercritical nature ($\text{Re}(k_2) < 0$) with a transition to doubly periodic motion in the latter case. As McLaughlin and Martin [6] have shown for the real Lorenz equations, the case $\text{Re}(k_2) > 0$ is a sub-critical Hopf bifurcation. In our problem the actual determination of the criterion $\text{Re}(k_2) = 0$, giving the exact delimitation between the two types of behaviour, is extremely messy to calculate. Instead we have performed a limited number of numerical integrations to examine the kind of bifurcations which can occur.

There are five parameters in the equations (1.1): σ , b , r_1 , r_2 , and e . We will concern ourselves mainly with holding these fixed, except for r_1 , which is thus the bifurcation parameter. Clearly a complete numerical analysis of (1.1) is out of the question in a paper of the present nature, and we can only give a brief idea of the types of bifurcation which occur.

In fig. 3, we show a stability diagram in (r_1, r_2) space which shows the dependence of r_{1c} and r_{2c} on r_2 at $\sigma = 2$, $e = 3r_2$, $b = 0.8$. These represent the analytic results of sections 2 and 3. Thus to the left of A, the origin is linearly stable; to the right, it is unstable and the stable solution is a limit cycle whose frequency decreases to zero




Fig. 3. Stability diagram in (r_1, r_2) space showing r_{1c} , r_{2c} , and also the approximate region of subcritical transition. Parameter values are $b = 0.8$, $\sigma = 2$, $e = 3r_2$.

as $r_2 \rightarrow 0$. This limit cycle is linearly stable between A and C, and linearly unstable below C. Fig. 4 shows the decay of a trajectory in the rotating $(\text{Re } x, \text{Re } y)$ phase space at a value of $r_1 = 254$ (here $r_{1c} = 255.67 \dots$ at $r_2 = 1$); motion is from top to bottom, and consists of a fast oscillation of slowly decaying amplitude, which slowly precesses around the underlying limit cycle (set of equilibria).

At a value of $r_1 = 256 > r_{1c}$, a small perturbation to the limit cycle initially grows as predicted by the analysis, but the final resultant motion is hardly of small amplitude, though in other respects it resembles the result of a

supercritical bifurcation. Later stages in the evolution of a small perturbation to the limit cycle are shown in fig. 5. The motion is anti-clockwise. In fig. 5a, we see that from $t = 835$ to $t = 935$, the solution precesses slowly around the underlying limit cycle, with a precession rate that slowly increases. Fig. 5b shows the solution from $t = 900$ to 1,000: the bottom of 5a is visible on the inside (though the scale is different); we see that the precession rate increases dramatically (more than three complete precessions are visible) and the solution rapidly attains a steady form, which is shown in fig. 5c. Corresponding plots of $\text{Re}(x)$, $\text{Re}(y)$ and z are shown in fig. 6.



Fig. 4. Slowly decaying trajectory in (x_R, y_R) phase space at $r_1 = 254 < r_{1c} = 255.67 \dots$, $r_2 = 1$; other parameters as for fig. 3.

Fig. 6 shows that between $t = 50$ and $t = 100$ (different initial values were chosen so that the final steady solution was approached rapidly, in $t < 50$) ten full precessions occur, and there are about eight to nine fast oscillations per precession. Close examination of the time plots suggests that (as fig. 5c also indicates) the solutions are periodic (or very nearly so) with a period of forty-two fast oscillations, or five precessions.

The large amplitude of this oscillation suggests that the bifurcation at $r_1 = r'_{lc} = 255.67 \dots$ is sub-critical, and so $\text{Re } k_2 > 0$

there. In this case, we suggest that at values $r_1 < r'_{lc}$, there may exist an unstable and a (larger) stable torus, for which the amplitude of the smaller tends to zero as $r_1 \rightarrow r'_{lc}$. Generally, the nature of the stable large-amplitude solution is inaccessible to analysis, but we suggest on the basis of fig. 5c (and other numerical evidence) that it also consists of motion on a torus which one might expect to be doubly periodic. As r_1 decreases to a value $r''_{lc} < r'_{lc}$, the stable and unstable forms may coalesce and vanish: or, they may both be unstable. The dashed line marked B in fig. 3 represents (roughly) the





Fig. 5. Evolution of a small disturbance to the limit cycle at $r_1 = 256 > r'_{lc}$; other parameters as for fig. 4; (a) $t = 835$ to 935.


Fig. 5. (b) $t = 900$ to $1,000$.

position of the graph of r_1^* as a function of r_2 . It has been determined on the basis of a limited number of integrations. As long as B lies above C, the bifurcation will then be subcritical. If B intersects C, then at this point $\text{Re}(k_2) = 0$, and for greater values we should expect $\text{Re}(k_2) < 0$, and a supercritical bifurcation: we have not determined this coefficient, however. Although the motion exhibited in fig. 6 is not of small amplitude, the nature of the numerical solution nevertheless suggests a multiple time scales analysis. One possibility for such an analysis is that the parameters are 'close' to the values at which $k_{2R} = 0$: an analysis near this point (and for $|r_1 - r_1^*| \ll 1$) would describe the *outer* stable

oscillation as well: on the other hand its form may be due to the largeness of r_1 .

Fig. 7 exhibits the finite amplitude stable torus at a subcritical value of $r_1 = 220 < r_1^*$. As far as can be seen, the motion is doubly periodic (frequency locking does not occur): the occurrence of precisely two incommensurate frequencies may be (and has been) checked by spectral analyses, and by computing phase plots of $|X|^2$ versus $|Y|^2$ (which appear as limit cycles in a doubly periodic motion).

We have found further bifurcations of familiar type, by decreasing r_2 . Of the two possibilities, we have found no further instability of the 2-torus to a higher dimensional attractor, but

Fig. 5. (c) Final motion, $t \rightarrow \infty$.

we have found an approach to 'chaos' by 'period-doubling' of the torus; specifically, $|X|^2$ versus $|Y|^2$ (or Z) phase plots exhibit period-doubling limit cycles as r_2 is reduced at, for example, $r_1 = 40$. This is analogous to the period-doubling approach to chaos in the real Lorenz equations as r_1 is reduced from infinity (Robbins, 1979). Fig. 8 shows an example of a slightly supercritical ($r_1 > r_{1c}$) chaotic motion (in the rotating frame) at values $r_1 = 60$, $r_2 = 0.02$, close to the real Lorenz transition to chaos (at $r_1 = 58$, $r_2 = 0$).

Generally speaking, the effect of the com-

plexification is to convert oscillatory states into 'doubly' oscillatory ones: limit cycles into tori, fixed points into limit cycles. Thus increasing the dimension of the system effectively increases the dimension of the attractors, and there is no particular reason to suppose the rich behaviour of the real Lorenz equations is otherwise modified, except in this manner: fuller numerical investigation is necessary for the corroboration of this statement. The chaotic behaviour in the real Lorenz model does not, however, appear to extend very far into the parameter space of the complex Lorenz equa-






Fig. 6. x_R , y_R and z plots versus time for fig. 5c.




Figure 7. Nonperiodic solution at $r_1 < r_{1c}$: same parameters as fig. 2.




Fig. 8. Chaotic motion at $r_1 = 60$, $r_2 = 0.02$; other parameters as fig. 3.

tions, at least when $e = 3r_2$. The general tendency of complexifying the equations at fixed r_1 is thus to replace chaotic behaviour by motion on a torus or a limit cycle.

Acknowledgements

We are grateful to Joel Smoller and Bob Williams for some very interesting conversations. One of us (MJM) was supported by the Department of Education in Dublin on a Postdoctoral Fellowship. We would also like to express our gratitude to colleagues at Dublin Institute for Advanced Studies School of Theoretical Physics where we held some 'teach yourself' seminars and where this work was originally started.

References

- [1] J.D. Gibbon and M.J. McGuinness; preprint (1980): to appear in *Physica D*.
- [2] E.N. Lorenz, *J. Atmos. Sci.* 20 (1963) 130.
- [3] J.D. Gibbon, M.J. McGuinness and I.N. James, Amplitude equations at the critical points of unstable dispersive systems, to appear in *Proc. Roy. Soc. Series A* (1981).
- [4] J.D. Gibbon and M.J. McGuinness, *Phys. Letts.* 77A (1980) 295.
- [5] J. Brindley and I.M. Moroz, preprint 1980; also *Phys. Letts.* 77A (1980) 441.
- [6] J.B. McLaughlin and P.C. Martin, *Phys. Rev. A.* 12 (1975) 186.
- [7] E.A. Coddington and N. Levinson, *Theory of Ordinary Differential Equations* (McGraw-Hill, New York, 1955).
- [8] D. Ruelle and F. Takens, *Comm. Math. Phys.* 20 (1971) 167.
- [9] D.D. Joseph, The bifurcation of T-periodic solutions into nT-periodic solutions and TORI, in *Synergetics-A Workshop*, H. Haken, ed. (Springer, New York, 1977).
- [10] O. Lanford III, Bifurcation of periodic solutions into invariant tori: the work of Ruelle and Takens, in *Non-linear Problems in the Physical Sciences and Biology, Lecture Notes in Mathematics* (Springer Verlag, New York) 322 (1977) 159.
- [11] J.T. Stuart, *J. Fluid Mech.* 9 (1960) 353.
- [12] H. Haken, *Z. Physik B29* (1978) 61.
- [13] W. Eckhaus, *Studies in Nonlinear Stability Theory*, (Springer, New York, 1965).
- [14] A.H. Nayfeh, *Perturbation Methods* (Wiley, New York, 1973).
- [15] H. Fujisaka and T. Yamada, *Phys. Lett.* 66A (1978) 450.
- [16] C. Escher, *Ber. Buns. Gesell. Phys. Chem.* 84 (1980) 387.
- [17] P.R. Fenstermacher, H.L. Swinney and J.P. Gollub, *J. Fluid Mech.* 94 (1979) 103.
- [18] K.A. Robbins, *Siam J. Appl. Math.* 36 (1979) 457.
- [19] S.S. Hassan, P.D. Drummond and D.F. Walls, *Optics Comm.* 27, (1979) 480.
- [20] K. Ikeda, *Optics Comm.* 30 (1979) 257.
- [21] R. Bonifacio and L.A. Lugiato, *Phys. Rev. A* 18 (1978) 1129.