

Cauldron subsidence and subglacial floods

G. W. Evatt and A. C. Fowler

Mathematical Institute, Oxford University
24-29 St Giles', Oxford OX1 3LB, England

October 23, 2006

Abstract

Ice cauldrons are depressions which form at the surface of ice sheets when an underlying subglacial lake empties. Notable examples of such cauldrons occur on the surface of the Vatnajökull ice cap in Iceland, and in particular are formed when subglacial volcanic eruptions occur. More generally, cauldrons will form when a subglacial lake empties during a jökulhlaup. The rate of subsidence of the ice surface is related to the rate at which the subglacial water empties from the lake. We use a viscous version of classical beam theory applied to the ice sheet to determine the relation between the subsidence rate and flood discharge, and we use the results to make inferences concerning ring fracture spacings in cauldrons, the consequent effect on flood discharge dynamics, and the likely nature of subsidence events in the Antarctic Ice Sheet.

Keywords: Ice cauldrons, jökulhlaups.

1 Introduction

Ice cauldrons are depressions which form on ice caps when basal ice is melted to form subglacial lakes which then drain (Björnsson, 1988). Two well-known examples are the Skaftá cauldrons on the ice cap Vatnajökull in Eastern Iceland, which lie above subglacial lakes which drain every year. A dramatic example of cauldron formation occurred during the 1996 eruption at Gjálp, also under Vatnajökull. Figure 1 shows a forming cauldron in its early stages. The ice surface subsided at a rate of some 12 metres per hour, as the basal fissure eruption melted the subglacial ice. The resultant meltwater (some three km³) ran off towards the subglacial lake Grímsvötn, whence a jökulhlaup eventually emerged. Another less dramatic example has recently been observed at the surface of the Antarctic Ice Sheet, where a subsidence of the order of four metres has occurred over a period of some years above an inferred subglacial lake, whose area is likely to be hundreds of km² (Wingham and others, 2006).

The subsidence is not passive, because ice is viscous, and the drawdown of the ice which is necessary to accommodate the discharge of the subglacial floodwater requires the floodwater to be underpressured, in order that it can suck down the ice.




Figure 1: An ice cauldron forming after the Gjalp eruption under Vatnajökull in 1996. The cauldron is about two kilometres in diameter and hundreds of metres deep, and ring shear fractures can be seen, indicating yield of the ice. The subsidence rate of the ice surface was initially about 12 metres per hour. Photo by Magnús Tumi Guðmundsson, University of Iceland, obtained from <http://www.hi.is/~mmh/gos/photos.html>. For further information see Guðmundsson and others (2004).

The underpressure competes with the viscosity of the ice, and the consequence is that there must be a dynamic relationship between the underpressure and the rate of subsidence of the ice.

Our purpose in this paper is to propose the nucleus of a theory that will predict how the rate at which the ice surface deforms is related to the water pressure in the basal lake. This theory will apply to cauldrons such as that in Figure 1, and also to the smaller depressions which have recently been observed on the surface of the Antarctic Ice Sheet (Wingham and others, 2006). The basis for our theory lies in the approximate treatment of the ice overlying the lake as a viscous ‘beam’. The theory is essentially that of classical beam theory in elasticity, except that viscosity replaces elasticity, thus introducing an extra time derivative into the problem. The approximation is based on the large aspect ratio of the ice, and is thus a version of the shallow ice approximation; it is similar to the description of ice shelf dynamics, insofar as normal stresses are dominant, but there the similarity ends. For reasons of space, our theoretical treatment is sketchy. For more details, see Evatt (2006).




Figure 2: Schematic diagram of cauldron geometry.

2 Viscous beam theory

2.1 Basic equations

The geometry of the situation we wish to consider is shown in Figure 2. For simplicity and pedagogy we work in two dimensions x and z . We let the corresponding components of velocity be u and w , and we let the deviatoric longitudinal stress be τ_1 , and the shear stress be τ_3 . The equations of slow flow (that is, flow at zero Reynolds number, also known as Stokes flow) can then be written in the form (where here and subsequently, subscripts x and z and later t denote partial derivatives)

$$\begin{aligned}
u_x + w_z &= 0, \\
0 &= -p_x + \tau_{1x} + \tau_{3z}, \\
0 &= -p_z + \tau_{3x} - \tau_{1z} - \rho_i g, \\
u_z + w_x &= A\tau^{n-1}\tau_3, \\
2u_x &= A\tau^{n-1}\tau_1, \\
\tau^2 &= \tau_1^2 + \tau_3^2,
\end{aligned} \tag{1}$$

where p is pressure, τ is the second stress invariant, A is the flow law rate coefficient, typically dependent on temperature, n ($= 3$) is the exponent in Glen's law, ρ_i is ice density, and g is the acceleration due to gravity. The boundary conditions at the top and bottom surfaces are those of stress continuity and material integrity (manifested as the kinematic condition). These boundary conditions take the following forms: on $z = s$,

$$\begin{aligned}
(-p + \tau_1)s_x - \tau_3 &= 0, \\
\tau_3 s_x + p + \tau_1 &= 0, \\
w &= s_t + u s_x - a,
\end{aligned} \tag{2}$$

while at the lake roof $z = h$,

$$(p - \tau_1)h_x + \tau_3 = p_w h_x,$$

$$\begin{aligned} p + \tau_1 + \tau_3 h_x &= p_w, \\ w &= h_t + u h_x - m, \end{aligned} \quad (3)$$

where a is the surface accumulation rate and m is the basal melt rate.

In addition, we suppose that the water pressure in the lake is hydrostatic, which implies

$$[p_w + \rho_w g h]_x = 0 \quad \text{on} \quad z = h. \quad (4)$$

Finally, if the net volume flux out of the lake is Q , then

$$Q = - \int_{A_L} w \, dS, \quad (5)$$

where A_L is the lake area. Our two-dimensional theory assumes the lake is rectangular in shape.

2.2 Non-dimensionalisation

Let us suppose that the lake is of width l and the ice is of depth d . We define the aspect ratio to be

$$\nu = \frac{d}{l}. \quad (6)$$

The basis of viscous beam theory (Howell, 1996, Ribe, 2001) is the assumption that ν is small. We suppose that a suitable scale for the rate of subsidence is w_s , and that a suitable scale for the longitudinal stress is τ_A . In beam theory, the greatest stresses are normal stresses, and if we consider also the variation of the flow rate coefficient A with temperature, then the rate-controlling value of A is its minimal value, obtained at the surface. We let A_s denote this surface value, and suppose for simplicity that it is constant. The drainage from the lake is controlled by the effective pressure of the lake (Nye, 1976), which is defined as

$$N = \rho_i g(s - h) - p_w, \quad (7)$$

and we suppose that a typical value of this is N_0 , and that the time scale appropriate to channel drainage is t_0 .

It is appropriate to scale the variables as

$$\begin{aligned} z, h, s &\sim d, \quad x \sim l, \quad p - \rho_i g(s - z) \sim \tau_A, \quad \tau_1 \sim \tau_A, \quad \tau_3 \sim \nu \tau_A, \\ A &\sim A_s, \quad w \sim w_s, \quad u \sim \nu w_s, \quad t \sim t_0, \quad Q \sim A_L w_s, \quad N \sim N_0. \end{aligned} \quad (8)$$

The beam stress is chosen to be

$$\tau_A = \left(\frac{\nu^2 w_s}{d A_s} \right)^{1/n}, \quad (9)$$

in order to balance terms in the definition of longitudinal strain rate.

With these scales, the equations become

$$\begin{aligned}
\nu^2 u_x + w_z &= 0, \\
0 &= -\beta s_x + \tau_{3z} - p_x + \tau_{1x}, \\
0 &= -p_z - \tau_{1z} + \nu^2 \tau_{3x}, \\
u_z + w_x &= \nu^2 A \tau^{n-1} \tau_3, \\
2u_x &= A \tau^{n-1} \tau_1, \\
\tau^2 &= \nu^2 \tau_3^2 + \tau_1^2,
\end{aligned} \tag{10}$$

and we retain the now dimensionless A in case we might want at some point to have it as a function of temperature and thus depth: in the present paper, only the case $A = 1$ is considered.

The boundary conditions become, on the surface $z = s$,

$$\begin{aligned}
\tau_3 + (p - \tau_1) s_x &= 0, \\
\nu^2 \tau_3 s_x + p + \tau_1 &= 0, \\
s_t &= \mu (w - \nu^2 u s_x + a^*),
\end{aligned} \tag{11}$$

where

$$\mu = \frac{w_s t_0}{d} \tag{12}$$

and

$$a^* = \frac{a}{w_s}, \tag{13}$$

and on the lake roof $z = h$,

$$\begin{aligned}
\tau_3 + (p - \tau_1) h_x &= -\gamma \beta N h_x, \\
\nu^2 \tau_3 h_x + p + \tau_1 &= -\gamma \beta N, \\
h_t &= \mu (w - \nu^2 u h_x + m^*),
\end{aligned} \tag{14}$$

where

$$m^* = \frac{m}{w_s}. \tag{15}$$

The parameters γ and β are defined by

$$\beta = \frac{\rho_i g d}{\tau_A}, \quad \gamma = \frac{N_0}{\rho_i g d}. \tag{16}$$

The lake hydrostatic condition (4) becomes

$$[s + \delta h - \gamma N]_x = 0, \tag{17}$$

where

$$\delta = \frac{\rho_w - \rho_i}{\rho_i}, \tag{18}$$

and the lake refilling equation (5) becomes in dimensionless terms

$$Q = - \int_{x_-}^{x_+} w \, dx, \quad (19)$$

where x_- and x_+ are the upstream and downstream positions of the lake margins.

The deflation rate scale w_s must be determined, if the effective pressure scale is known. The way it is chosen is by selecting the parameter μ to balance terms in the hydrostatic equation, and we do this by choosing

$$\mu = \frac{\gamma}{1 + \delta}, \quad (20)$$

and we will see that this is appropriate for closed subsurface lakes.

The three other parameters involved in finding an approximate solution to this set of equations are γ , β and ν . Generally, we can assume $\gamma \lesssim 1$, $\nu \ll 1$ and $\beta \gtrsim 1$. We therefore begin by seeking approximate solutions for $\nu \ll 1$, and then subsequently considering possible choices for γ and β . The limit $\nu \rightarrow 0$ is the classical approximation associated with beam theory.

2.3 The beam equation

With $\nu \ll 1$, we have $w \approx w(x, t)$, $\tau \approx |\tau_1|$ and $u_z + w_x \approx 0$, whence

$$\begin{aligned} 2u_x &\approx A|\tau_1|^{n-1}\tau_1, \\ u &\approx v - zw_x, \end{aligned} \quad (21)$$

where $v(x, t)$ is to be determined. From the equations and boundary conditions (10), (11) and (14), we see that $p + \tau_1 \sim \nu^{21}$, and therefore we define

$$p + \tau_1 = -\nu^2 \sigma_3. \quad (22)$$

From (21), we find

$$\tau_1 = \left(\frac{2}{A} \right)^{1/n} \frac{v_x - zw_{xx}}{|v_x - zw_{xx}|^{(n-1)/n}}. \quad (23)$$

We now define the three quantities M , the bending moment, S , the shear force, and T , the tension, as

$$M = - \int_h^s 2z\tau_1 \, dz, \quad S = \int_h^s \tau_3 \, dz, \quad T = \int_h^s \tau_1 \, dz. \quad (24)$$

We also define the secondary quantities U , the uplift force, and L , the lifting torque, to be

$$U = \int_h^s \sigma_3 \, dz, \quad L = \int_h^s z\sigma_3 \, dz. \quad (25)$$

¹Strictly this requires $\gamma\beta \sim \nu^2$, but for the moment we only use this to motivate the definition in (22). See also the comment after (28).

Integrating the force balance equations and applying the boundary conditions on $z = s$ and $z = h$, we then find that

$$\begin{aligned} 2T_x + \nu^2 U_x &= -\gamma\beta N h_x + \beta(s - h)s_x, \\ M_x - \nu^2 L_x + S &= \gamma\beta N h h_x - \frac{1}{2}\beta(s^2 - h^2)s_x, \\ S_x &= \frac{\gamma\beta N}{\nu^2}. \end{aligned} \tag{26}$$

Subject to suitable boundary conditions, these equations will enable us to provide a closed solution which determines the surface depression rate $-w$ in terms of the underlying effective pressure N .

The simplest assumption for the boundary conditions is to assume that the positions of the lake margins are fixed, so that the values x_{\pm} are constant. The ice at the margins does not subside, so we put

$$w = 0 \quad \text{at} \quad x = x_{\pm}. \tag{27}$$

In reality, the margins will move as the ice sags, forming a moving grounding line problem whose solution in the case of ice sheet/ice shelf transition is itself non-trivial to solve (Schoof, 2006).

The second condition at a margin is less easy to motivate. The horizontal velocity scale in the beam is νw_s (see (8)), so that for the rapid subsidence in the Vatnajökull cauldrons, we would expect to prescribe $u = 0$ and thus $v = w_x = 0$ at the margin (clamped conditions). For the much smaller rates of subsidence appropriate to the Antarctic sub-glacial lake (Wingham and others, 2006), the horizontal velocity scale is much smaller than that of the surrounding ice sheet ice, and a more appropriate boundary condition may be to prescribe zero longitudinal stress, thus $\tau_1 = 0$. Another argument in favour of this is that the beam stress τ_A will typically be much larger than the normal stress scale in the ice sheet.

Another consideration is the effect of temperature in the ice. The rate of subsidence in the beam depends on the surface rate constant A_s and the beam stress τ_A . The effective beam viscosity is thus $\eta = \frac{1}{2A_s\tau_A^{n-1}}$. Generally the beam stresses are higher than the basal shear stress beyond the lake, while the surface rate coefficient may be much lower, for example in Antarctica. If we suppose that the beam ice is much stiffer than the ice sheet ice, then this also suggests conditions of zero longitudinal stress at the margins, i.e., $\tau_1 = 0$.

None of these arguments is watertight, and the issue of the correct choice of effective boundary conditions is one of some difficulty. We do not pursue the issue further in this paper, but to be specific we will choose conditions of zero longitudinal stress at the margins, based on the idea that the ice sheet ice is very weak in comparison to the beam ice.

The neglect of terms in ν makes it unlikely that we can satisfy this pointwise, but we can certainly suppose that this condition can be applied in a vertically integrated sense, whence we prescribe

$$M = T = 0 \quad \text{at} \quad x = x_{\pm}. \tag{28}$$

We solve (26) by formally assuming $\gamma\beta \sim \nu^2$. This is useful for exposition, but in fact the resulting approximations are still valid even if $\gamma\beta \gg \nu^2$, as is easily checked. The only term we retain on the right hand sides of (26) is therefore the forcing term for S . Neglecting terms of relative $O(\nu)$, we then have that $T = T(t)$, and the boundary conditions (28) imply that $T = 0$. A consequence of this from (23) is that

$$v_x = z_m w_{xx}, \quad (29)$$

where z_m is the mid-depth point,

$$z_m = \frac{1}{2}(s + h), \quad (30)$$

and from this it follows that

$$M = \frac{D_n w_{xx}}{|w_{xx}|^{(n-1)/n}}, \quad (31)$$

where

$$D_n = \frac{nH^{(2n+1)/n}}{(2n+1)A^{1/n}}, \quad H = s - h, \quad (32)$$

and the beam equation takes the resultant form

$$\left\{ \frac{D_n w_{xx}}{|w_{xx}|^{(n-1)/n}} \right\}_{xx} = -\frac{\gamma\beta N}{\nu^2}, \quad (33)$$

subject to

$$w = w_{xx} = 0 \quad \text{at} \quad x = x_{\pm}. \quad (34)$$

An expression for N results from the hydrostatic equation (17), together with the approximate kinematic equations

$$s_t \approx \mu(w + a^*), \quad h_t \approx \mu(w + m^*). \quad (35)$$

Let us suppose for simplicity that a^* and m^* are negligible. Then the depth $H = s - h$ depends only on x . If we suppose that it is constant, then $s_x = h_x$, and the integral of (17) is (using the definition of μ in (20))

$$s = s_+ + \mu\{N - N_+(t)\}, \quad (36)$$

where $s = s_+$ (constant, since $w = 0$ at x_+) and $N = N_+$ at $x = x_+$. Differentiating, we finally have

$$N_t = \dot{N}_+ + w. \quad (37)$$

By assumption of the choice of scales t_0 , N_0 and w_s , the terms in (37) are all $O(1)$, and it is this which motivates the choice (20).

2.4 An approximate solution

An approximate solution to the beam equation (33) and the lake refilling equation (37) can be obtained using a boundary layer theory. Details are presented by Evatt (2006). Here we summarise the results. We first estimate the size of the term $\gamma\beta/\nu^2$ for the cauldron in Figure 1. The ice depth is about 500 m and the cauldron width some 2 km, so we take $\nu \approx 0.2$. The product $\gamma\beta = N_0/\tau_A$ is the ratio of the channel effective pressure scale to the beam stress. Typical values of $w_s \approx 10^5$ m y⁻¹ (12 metres per hour), $A_s \approx 10^{-2}$ bar⁻³ y⁻¹ (based on a surface temperature of -10° C, see Paterson, 1994), $n = 3$; with these values, we find $\tau_A \approx 20$ bar. This is comparable to typical estimates of channel effective pressure (Evatt, 2006), and then $\gamma\beta \sim O(1)$, and $\gamma\beta/\nu^2 \gg 1$. In this case, the solution becomes of singular perturbation type, and supports boundary layers near the lake margins.

There is a central flat portion of the ice beam in which $N \approx 0$, the ice is essentially floating. The refilling equation together with (19) then implies that the boundary condition for the flow in the channel is

$$\dot{N}_+ = \frac{Q}{x_+ - x_-}, \quad (38)$$

which is essentially the same refilling condition that has been used in the past (Fowler, 1999). This is a first conclusion of the present study.

Near the margins, there are boundary layers of thickness

$$\sigma = \left(\frac{\nu^2 D_n}{\gamma\beta} \right)^{\frac{n}{2(n+1)}}, \quad (39)$$

and the governing boundary layer equation has a similarity solution if we assume that the rising limb of the flood hydrograph is given by a power law of the form

$$N_+ = \frac{c}{(t^* - t)^\alpha}, \quad Q = \frac{\alpha c}{(t^* - t)^{\alpha+1}}, \quad (40)$$

where c and α are positive constants; this was shown by Nye (1976), with $\alpha = 3$; here t is time, and t^* a constant chosen to fit the numerically computed form of the hydrograph. An essentially analytic solution for w can thus be obtained, and from this the surface deformation history can be computed by integrating (35). The result of doing this is shown in Figure 3, which shows a theoretical subsidence sequence above a lake whose discharge follows the Nye rising hydrograph (40). The parameter choices for this lake are chosen to be appropriate for a sub-Antarctic lake.

3 Cauldrons and ring fractures

It is of interest to calculate the maximum stress in the beam. From (23), the maximum stress is at the surface and base of the ice, and is given by

$$|\tau_1|_{\max} = \left| \frac{H w_{xx}}{A} \right|^{1/n}. \quad (41)$$




Figure 3: Solution of (35) (starting at $t = 0$ d) at times $t = 360$ d, $t = 371$ d, $t = 382$ d, $t = 393$ d using (40) for N_+ , with $t^* = 1$, $c = 2.5 \times 10^{-3}$, $\alpha = 3$, $\gamma\beta/\nu^2 = 15.8$, and a time scale of $t_0 = 416$ d. The units of vertical subsidence are in metres, and the units of horizontal distance are km. The results in the figure correspond to a choice of $\mu d = N_0/\rho_i g = 3$ m, and thus $N_0 = 0.3$ bars. The results are plotted for a lake of width 20 km, in which case if $d = 4$ km, $\nu = 0.2$ and thus $\gamma\beta = 0.63$, corresponding to a value of $\tau_A = 0.48$ bars. No surface fracture is expected in this case.

We can use the boundary layer theory to estimate the maximum stress. After some algebra, we find that the maximum dimensional stress is

$$\tau_{\max} \sim \left(\frac{\gamma\beta}{\nu^2} N_+ \dot{N}_+ \right)^{1/4} \tau_A, \quad (42)$$

where we take $n = 3$, and this occurs at a distance from the margin of

$$\Delta x \sim \left(\frac{\dot{N}_+}{N_+^3} \right)^{1/8} \frac{\nu^{1/2} l}{(\gamma\beta)^{1/4}}, \quad (43)$$

where l is the lake width. If τ_{\max} reaches the yield stress τ_c of ice, then the ice will fracture, forming a crevasse, and the overlake ice will reset itself to the application of effective boundary conditions at the position of this ring fracture. Thereafter, continued rise of effective pressure will allow a new ring fracture to occur in-lake of the old fracture, and in this way a sequence of such fractures may form, with a spacing indicated by (43), as seen in Figure 1. If we equate $\tau_{\max} = \tau_c$, and use (42)

to eliminate \dot{N}_+ , we find that the fracture spacing should be

$$\Delta x \sim \left(\frac{\tau_c}{\tau_A}\right)^{1/2} \left(\frac{\nu^2}{\gamma\beta}\right)^{3/8} \frac{l}{N_+^{1/2}}. \quad (44)$$

If $\tau_A \gg \tau_c$ and $\gamma\beta/\nu^2 \gg 1$, as we surmise for the cauldron of Figure 1, then we see that $\Delta x \ll l$, as is clearly observed in Figure 1. Having said that, it is not easy to get good quantitative agreement with observed fracture spacing, for which a consideration of viscoelastic effects may be necessary.

4 Floods in the Antarctic

Recently, Wingham and others (2006) have observed subsidences of the Antarctic ice sheet surface of the order of four metres occurring over a period of some three years. They ascribe these local deflations to small lake flooding events (one lake floods into another 300 km away, above which the ice surface rises several metres). Our theory can be applied to these floods provided the value of μ in (35) is appropriate; specifically, we need $\mu d \approx 4$ m. Since from (20) and (16), $\mu \approx \gamma = N_0/\rho_i g d$, we see that the depth of subsidence is of order $N_0/\rho_i g$. Assuming this is 4 metres, we infer that $N_0 \approx 0.4$ bar. This is essentially the value that was inferred for the effective pressure under ice stream B (Whillans ice stream) (Blankenship and others 1987), where the ice overlies marine sediment (Kamb, 2001). The lake described by Wingham and others (2006) lies in the Adventure subglacial trench, which was presumably a large lake even before Antarctic glaciation (Drewry, 1983, sheet 6), and thus likely to be bedded with glaciolacustrine sediments. We interpret the small subsidence observed by Wingham and others as indicative of a high pressure canal-type drainage system (Walder and Fowler, 1994), although jökulhlaups in such drainage systems have not previously been described. Other interpretations are possible, for example that the lake was small and simply emptied.

Figure 3 shows a sequence of profiles of the surface profile s as time increases obtained by solving the surface kinematic equation (35). The two features of these shapes are their depth and the sharpness of the boundary layers. The depth is controlled by the parameter $\mu \approx \gamma$, while the boundary layer thickness is $(\gamma\beta/\nu^2)^{1/4}$. The depression of the surface by about 3 metres (over an ice depth of 4000 metres) suggests that $\mu \sim 10^{-3}$ and $N_0 \sim 0.3$ bars. To calculate $\gamma\beta = N_0/\tau_A$, we use the definition of τ_A in (9). With $\nu \sim 0.2$, $w_s \sim 3$ m y⁻¹, $d \sim 4000$ m, $A_s \sim 10^{-4}$ bar⁻³ y⁻¹ (Paterson, 1994, p. 97, at -50° C), we find $\tau_A \sim 0.67$ bars, and thus $\gamma\beta \sim 0.45$. With $\nu = 0.2$ (a 20 km wide lake), we have the boundary layer width $(\nu^2/\gamma\beta)^{1/4} \sim 0.55$. In this case the boundary layers are quite wide, as indicated in Figure 3, where they essentially extend to the centre of the depression.

5 Conclusions

The formation of cauldrons on the surface of ice caps is associated with the emptying of subglacial lakes. The rate of subsidence is directly related to the rate of lake discharge, but requires an underpressure in the lake in order to draw down the overlying ice. Therefore the underpressure in the lake is related to the rate of subglacial lake discharge. This has consequences for theories of jökulhlaups, since this relationship is an essential part of the flood hydrograph theory. Previous theories (e. g., Nye, 1976)) have in effect simply assumed a flotation condition for the ice over the lake, but it is not obvious that this will apply when the lake is isolated from the atmosphere. In this paper we have presented the elements of a theory which can be used to predict the relationship between ice surface subsidence and lake underpressure. The theory is that of a viscous beam (Howell, 1996, Ribe, 2001), and is based on the assumption of small aspect ratio (depth/length) of the ice above the lake. Providing a further parameter $\gamma\beta = N_0/\tau_A$ is not too small, where N_0 is the scale of the effective pressure at the entrance to the lake drainage channel and τ_A is the size of the deviatoric normal stress in the ice, a boundary layer theory shows that in effect the bulk of the ice can be considered to be approximately in flotation, thus vindicating the Nye assumption for closed subglacial lakes.

For rapid channel drainage, such as occurs following subglacial volcanic eruptions, the induced beam stresses may be very large (of order ten bars), and will cause closely spaced ring fractures to occur in the beam as it collapses. The present theory can be used to predict an approximate fracture spacing, but does not seem to give good quantitative agreement with observation, perhaps because the model is not sophisticated enough.

The recent observation of small subsidence events (of the order of metres) on Antarctica is indicative of small floods. The duration of these floods (with a time scale of a year) is comparable to those of simulations using Nye's (1976) theory (Evatt and others, 2006), but the slight depressions either indicate very shallow lakes, or may imply that such floods occur in channelised flow over soft sediments at low effective pressure. We consider this latter (tentative) explanation to be more robust. It raises further questions concerning the nature of hydraulic drainage under ice sheets.

Acknowledgements

G. W. E. thanks W. H. R. Evatt for financial support, and the Centre for Polar and Ocean Modelling at University College, London for the award of a research assistantship. A. C. F. thanks the University of Limerick for their continued support of his research.

References

- Björnsson, H. 1988. *Hydrology of ice caps in volcanic regions*. Reykjavik, Societas Scientarium Islandica, University of Iceland.

- Blankenship, D. D., C. R. Bentley, S. T. Rooney and R. B. Alley. 1987. Till beneath ice stream B. 1. Properties derived from seismic travel times. *J. Geophys. Res.* **92** (B9), 8903–8911.
- Drewry, D. J. 1983. *Antarctica: glaciological and geophysical folio*. Cambridge, U. K., Scott Polar Research Institute.
- Evatt, G. W. 2006. *Jökulhlaups and sub-glacial floods*. D. Phil. thesis, Oxford University (*sub judice*).
- Evatt, G. W., A. C. Fowler, C. D. Clark and N. Hulton. 2006. Subglacial floods under ice sheets. *Phil. Trans. Roy. Soc.* **364** (1844), 1769–1794.
- Fowler, A. C. 1999. Breaking the seal at Grímsvötn. *J. Glaciol.* **45** (151), 506–516.
- Guðmundsson, M. T., F. Sigmundsson, H. Björnsson and T. Högnadóttir. 2004. The 1996 eruption at Gjálp, Vatnajökull ice cap, Iceland: efficiency of heat transfer, ice deformation and subglacial water pressure. *Bull. Volcanol.* **66** (1), 46–65.
- Howell, P. D. 1996 Models for thin viscous sheets. *Eur. J. Appl. Math.* **7** (4), 321–343.
- Kamb, B. 2001. Basal zone of the West Antarctic ice streams and its role in lubrication of their rapid motion. In: *The West Antarctic Ice Sheet: behavior and environment*, eds. R. B. Alley and R. A. Bindshadler, pp. 157–199. Washington, D. C., Antarctic Research Series, A. G. U.
- Nye, J. F. 1976. Water flow in glaciers: jökulhlaups, tunnels, and veins. *J. Glaciol.* **17** (76), 181–207.
- Paterson, W. S. B. 1994. *The physics of glaciers*, 3rd edition. Oxford, Pergamon.
- Ribe, N. M. 2001. Bending and stretching of thin viscous sheets. *J. Fluid Mech.* **433**, 135–160.
- Schoof, C. 2006. Marine ice sheet dynamics. Part I. The case of rapid sliding. *J. Fluid Mech.*, in press.
- Wingham, D. J., M. J. Siegert, A. Shepherd and A. S. Muir. 2006. Rapid discharge connects Antarctic subglacial lakes. *Nature* **440** (7087), 1,033–1,037.