

Title: Regulatory B cells in transplantation: Roadmaps to clinic

Running title: Bregs: translation to clinic

Authors: Joseph Beckett¹, Joanna Hester¹, Fadi Issa¹, Sushma Shankar¹

Affiliations: ¹ Transplant Research and Immunology Group, Nuffield Department of Surgical Sciences, University of Oxford, Oxford, United Kingdom

Corresponding Author:

Sushma Shankar

sushma.shankar@nds.ox.ac.uk

Transplant Research and Immunology Group

Nuffield Department of Surgical Sciences

University of Oxford

Level 6 John Radcliffe Hospital

Oxford

United Kingdom OX3 9DU

Funding Sources: No funding

Conflict of Interest Statement: No conflict of interest

Keywords: T-cells, B-cells, macrophages; Tolerance strategies and mechanisms; Novel immunosuppressants; Immunosuppression; Tolerance induction; Immunosuppression Experimental

Regulatory B cells in transplantation: Roadmaps to clinic

Beckett J, Hester J, Issa F, Shankar S

Abstract

Over the last two decades, an additional and important role for B cells has been established in immune regulation. Pre-clinical studies demonstrate that regulatory B cells (Breg) can prolong allograft survival in animal models and induce regulatory T cells. Operationally tolerant human kidney transplant recipients demonstrate B-cell associated gene signatures of immune tolerance, and novel therapeutic agents can induce Bregs in Phase I clinical trials in transplantation. Our rapidly expanding appreciation of this novel B cell subtype has made the road to clinical application a reality. Here, we outline several translational pathways by which Bregs could soon be introduced to the transplant clinic.

Introduction

Several cellular therapies have now been introduced in clinical trials of human kidney transplant recipients. The landmark ONE study explored the protective role of several regulatory cell types including regulatory T cells (Tregs), T regulatory type 1 cells (Tr1), dendritic cells (DC) and regulatory macrophages in transplantation [1, 2]. Over the last decade, evidence has emerged of an additional and novel regulatory cell type, regulatory B cells (Bregs), in transplantation. Adoptive transfer studies have demonstrated the protective capacity of mouse Bregs in prolonging allograft survival [3, 4]. Operationally tolerant, human kidney transplant recipients exhibit increased frequencies of circulating Breg populations and B cell signatures have been shown to be sensitive predictors of a tolerogenic state [5, 6]. Multiple Breg populations at several stages of B cell maturation have been identified, with a range of phenotypes and suppressive mechanisms. Bregs can suppress Th1 and Th17 responses and DCs, whilst promoting regulatory populations such as Tr1 cells and Treg. These subsets and their mechanistic functions have been extensively reviewed elsewhere, and include mouse marginal zone (MZ), transitional-2 marginal zone precursor (T2 MZP) and B10 Bregs, as well as human CD24^{hi}CD38^{hi}, CD24^{hi}CD27⁺ and TIM-1⁺ Bregs (Table 1) [7, 8]. Whilst much of the initial experimental work to characterise Breg has been in the context of autoimmunity, most Breg populations identified in transplantation are characterised by their ability to produce IL-10, and IL-10 itself can be induced in Bregs and their precursors by multiple stimuli (Table 1). In this review, we discuss recent advances in IL-10⁺ Bregs, and explore how this cell population may be harnessed therapeutically in clinical transplantation.

Immune Monitoring

The majority of transplant recipients require the use of potent immunosuppressants to prevent acute allograft rejection. However, not all recipients who take immunosuppression will need such high dosage regimens. In renal transplantation, patient weight determines calcineurin

inhibitor (CNI) and sirolimus doses, whilst azathioprine and mycophenolate doses are dictated largely by population-based data [9]. Many patients may be over-immunosuppressed, accounting for increased mortality from cardiovascular complications, cancer or infection in patients with functioning grafts. However, chronic rejection remains a leading cause of long-term graft loss, demonstrating the importance of adequate immunosuppression over time.

To date, there is no robust definition of immunological tolerance in humans; descriptions of ‘clinically operational tolerance,’ rely on the stability of graft function in absence of immunosuppression for a minimum period. Accurate identification of such recipients will be critical to guide the tailored reduction of immunosuppression and its associated co-morbidity in this cohort. Whilst blood may not necessarily be the best compartment by which to measure clinical pathology across all organ-types in transplantation, disease gene profiles showed good concordance between blood and solid tissues [10]. Evidence is now emerging of potential blood-based biomarkers of operational tolerance and predictors of rejection that focus upon Bregs. These may offer personalised regimens for patients who do not require as potent a regimen as the current standard, as well as for patients who are at risk of developing chronic rejection and losing their grafts (Fig. 1).

B cell-related tolerance signatures

The importance of Bregs in transplantation was highlighted by the identification of B cell signatures of tolerance in human kidney transplant recipients in the European Indices of Tolerance (IoT) and American Immune Tolerance Network (ITN) studies [11, 12]. Tolerant kidney transplant recipients (‘TOL’), who were defined by stable graft function with no immunosuppression for over 1 year, were compared to control cohorts including recipients with stable graft function who were receiving immunosuppression (‘SI’). Tolerant kidney transplant recipients demonstrated increased frequencies of naive CD19⁺ B cells and CD24^{hi}CD38^{hi}IL-10⁺ B cells in peripheral blood, and reduced donor-specific antibody (DSA) levels when compared to control groups other than healthy subjects, in whom levels were equivalent. Gene microarray analysis revealed a bias towards differential expression of B cell-related genes. The ITN study identified three B-cell-related genes, *IGKV4-1*, *IGLL1*, and *IGKVID-13* that were predictive of tolerance. These observations resulted in specific and sensitive cross-platform signatures that could predict tolerance in human kidney transplant recipients. Other groups described similar enrichments of IL-10⁺ Breg in tolerant kidney transplant recipients, as well as a decrease in differentiated plasma cells when compared to SI cohorts [13, 14].

Critically however, tolerance signatures in these studies were compared to those of patients still taking immunosuppression, and so may be confounded by effects of immunosuppressants upon circulating B cells. Moreover, these B cell signatures were similar to those described in healthy individuals, so may instead represent a signature of health, rather than of tolerance. The IoT authors repeated their analysis in three independent kidney transplant recipient cohorts [15]: the original IoT dataset; CNI-free dataset; dataset of kidney transplant recipients being weaned off steroids for clinical reasons. Using this approach, the authors established

that previous tolerance signatures were biased by immunosuppression. Prednisolone affected the IoT gene dataset [12], whilst prednisolone and azathioprine adversely affected the frequency of CD24^{hi}CD38^{hi} B cells. These findings were reinforced by work within our group, which described a decrease in transitional and naive B cell numbers in association with azathioprine, within peripheral blood of kidney transplant recipients [16].

However, the IoT re-analysis revealed a novel 9-gene set that was highly predictive of tolerance and which distinguished operationally tolerant kidney transplant recipients from SI recipients [17]. When applied to 173 SI kidney transplant recipients, this new gene set predicted that almost 12% were ‘probably tolerant.’ The signature importantly discriminated tolerant patients from healthy controls and B cell-associated pathways were still prominently represented in 3 of the 9-gene set: BCL2A1, EEF1A1 and IGLC1. The ITN also addressed the immunosuppression question, and found that expression of 2 of the 3-gene ITN B cell tolerance signature [11] increased over time in SI kidney transplant recipients, despite largely unchanged immunosuppression [18]. 14% of 124 recipients on varying immunosuppressive regimens consistently displayed the tolerance signature of IGKV1D-13 and IGKV4-1 over 2 years at multiple time-points [6]. Despite increased frequencies of all circulating B cell subsets in peripheral blood, these patients demonstrated a clinically significant improvement in renal function that increased over time, and persistently lower DSA levels when compared to patients who did not express this signature. It seems unlikely that immunosuppressive drugs prevented development of a B cell tolerance signature; moreover, the absence of immunosuppression was not a prerequisite for developing a B cell-correlated tolerogenic state. Notably, predicted tolerance was greatest in patients receiving a CNI-based maintenance immunosuppressive regimen and reduced in those receiving corticosteroids, mycophenolate, or thymoglobulin induction. Whether these associations reflect direct effects of immunosuppressants on B cells and are independent of a tolerogenic effect of Breg or whether they enable B cell changes that promote a reduction in the overall alloimmune response, remains to be determined.

Interestingly, a comparison of tolerant liver transplant recipients to those receiving immunosuppression such as CNIs and azathioprine, failed to demonstrate the changes in B cell subsets and B cell-related genes that characterised spontaneous tolerance in kidney transplantation, but which have been attributed to immunosuppression [19]. Moreover, the IoT and ITN signatures demonstrate little overlap, and other immune compartments may warrant inclusion as demonstrated by a meta-analysis of tolerance studies which identified a gene signature involving proliferation of B and CD4⁺ T cells and the inhibition of CD14⁺ monocyte-related functions [20]. These observations highlight that whilst B cell signatures of tolerance identified in renal transplantation may not be wholly attributed to confounding immunosuppression, there are likely alternate tolerance mechanisms in different allograft settings.

Two studies have attempted to wean CNIs in patients predicted to be at a low risk of rejection on the basis of clinical characteristics (absence of DSA, stable graft function, biopsies without evidence of inflammation). Both were stopped prematurely due to high rates of rejection

and/or the formation of DSA following attempted drug withdrawal [21, 22], highlighting current challenges of immunosuppressive drug minimisation in kidney transplant recipients with stable function. B cell-predominant signatures may provide a standardised tool towards personalised medicine for the monitoring of tolerant or low-risk recipients. We are close approaching the time for immunosuppression-minimising clinical trials based on signatures that include not just clinical parameters, but also immunology-based criteria, alongside careful patient selection and stringent monitoring.

Predictors of Rejection

The discovery of Breg association with tolerance signatures has been complemented by the emergence of B cells as potential predictors of rejection in renal transplantation. Peripheral blood analysis in a tolerant kidney transplant recipients cohort revealed an increase in memory B cells with inhibitory characteristics when compared to recipients undergoing chronic rejection. B cells in the tolerant cohort exhibited a decreased FcγRIIA/FcγRIIB ratio and an increase in BANK1 expression, which serves as a preventative signal of hyperactive B cell responses [14]. Silva et al. found that operationally tolerant kidney transplant recipients had a diverse BCR repertoire similar to that of healthy controls, unlike recipients undergoing rejection, thus mirroring observations in mouse models [23-25]. Operationally tolerant patients retained the ability to activate CD40 and signal transducer and activator of transcription 3 (STAT3) signalling pathways in the CD24^{hi}CD38^{hi} B cell subset enriched for human Breg. This ability was decreased in CD24^{hi}CD38^{hi} B cells of patients undergoing chronic rejection or GVHD [23, 26].

When appraising such studies however, it is important to note that mechanisms that maintain tolerance are not necessarily the same as those that predict tolerance development. Moreover, the lack of such a signature does not necessarily predict rejection. Studies that follow patient outcomes over time may be more informative. Clatworthy et al. uncovered the involvement of human Bregs in the prevention of acute rejection in transplantation [27]. In this study, the investigators suspended a randomised controlled trial of anti-CD20 mAb rituximab induction therapy versus anti-CD25 mAb dactilizumab after recruitment of 13 human kidney transplant recipients: a high incidence of acute cellular rejection occurred in the rituximab group (83% vs. 35% in the dactilizumab group). The potential role of Bregs in the prevention of allograft rejection has since been reiterated in renal, liver and lung transplantation [28-32], with reduced frequencies of circulating IL-10⁺ Breg-enriched CD24^{hi}CD38^{hi} and/or CD24^{hi}CD27⁺ B cell subsets found in recipients who developed rejection. Similarly, graft versus host disease (GvHD) was associated with a decline in frequency and suppressive function of CD24^{hi}CD27⁺IL-10⁺ Breg in stem cell recipients [33].

Following on from this work, Cherukuri and colleagues found that kidney transplant recipients who underwent rejection demonstrated a decreased IL-10:TNFα ratio expressed by CD24^{hi}CD38^{hi} B cells, irrespective of steroid use [28]. CD24^{hi}CD38^{hi} B cells from rejecting recipients were functionally deficient *in vitro* when compared to those of control cohorts. Interestingly, the function and frequency of Treg were unchanged across all patient groups. At

the time of the late for-cause biopsy, the CD24^{hi}CD38^{hi} B cell IL-10 to TNF α ratio strongly predicted rejection, graft loss and the doubling of serum creatinine over a 3-year follow-up period, highlighting the clinical relevance of such findings. Using the definition of more naïve, CD24^{hi}CD38^{hi} B cells as ‘T1,’ and more mature CD24^{int}CD38^{int} B cells as ‘T2’ B cells, the authors identified that the T1/T2 ratio reflected the changes associated with IL-10:TNF α ratio [29]. Low T1/T2 ratios in stable kidney transplant recipients who were 2-years post-transplant, were independently associated with and strongly predicted graft outcomes over a 5-year follow-up period in an independent validation set, whereas clinical parameters including delayed graft function, eGFR and DSA levels were not predictive.

The interplay of different B cell effector and regulatory subsets in chronic rejection beyond DSA production, has been elegantly characterised in recent years. In an animal model of heart transplantation, AID/ μ S knockout mice in which B cells cannot undergo affinity maturation or secrete antibody, underwent chronic allograft vasculopathy (CAV) comparable to that of wild-type mice [34]. However, μ MT mice which were wholly deficient in both B cells and antibody, were protected from CAV. The use of chimeric mice, in which B cells were present but were unable to present antigen, demonstrated that both alloreactive T cell responses and CAV were significantly reduced in the absence of B cell-mediated antigen presentation. CAV was restored following adoptive transfer of polyclonal AID/ μ S KO B cells. These findings indicate that chronic rejection could occur in the complete absence of antibodies by mechanisms involving B cell-mediated antigen presentation. This may account for why some patients develop chronic rejection in the absence of detectable alloantibodies.

Work by Shiu and colleagues has further defined the complex interactions of B effector and regulatory subsets in the context of chronic rejection in human renal transplantation [35, 36]. The examination of peripheral blood from 65 patients with biopsy-proven AMR, non-immune related graft dysfunction, or stable graft function, revealed that approximately half of the biopsy-proven AMR samples did not demonstrate a CD4⁺ anti-donor IFN γ response unless either CD25⁺ Treg or CD19⁺ B cells were depleted. Notably, the depleted CD19⁺ B cells in these samples were polarised to IL-10 production following polyclonal stimulation with IgM and IgG. In addition, following depletion of Treg or CD19⁺ B cells, alloresponsiveness in patients with no histological signs of immune-mediated graft dysfunction was restored *in vitro*. Subsequent interrogation of the peripheral blood of renal transplant patients with AMR revealed the association of suppressed CD4⁺ anti-donor IFN γ production with a sustained improvement in eGFR following the optimisation of immunosuppression, rather than anti-CD19 mAb rituximab administration [37]. Those patients whose graft function improved upon immunosuppression optimisation demonstrated an increased proportion of T1 B cells in peripheral blood when compared to T2 B cells, and CD19⁺ B cell-dependent suppression of CD4⁺ anti-donor IFN γ production. Conversely, following rituximab administration, there was a reduction in the proportion of donor-reactive responses suppressed by CD19⁺ B cells, whilst the proportion of samples showing unregulated B cell-dependent IFN γ production increased. B cell-mediated suppression of IFN γ production, the associated increase in T1 B cells and slower declines in graft function was abolished by rituximab, which depleted transitional B cells for prolonged periods. Rituximab appeared to disturb the balance of the two opposing

roles of B cells, by selectively reducing the relative proportion of transitional regulatory B cells, while failing to sustainably deplete B effector cells that support anti-donor responses by antigen presentation.

Thus simple biomarkers based upon Breg and B cell effector composition and function, may be used in a carefully selected group of transplant recipients to monitor and pre-empt allograft dysfunction and rejection to improve both short and long-term outcomes.

Cellular Therapy

The use of Bregs as a cellular therapy will require the identification of a stable Breg population with unique phenotypic markers to allow their precise isolation. It remains unclear if the various methods of Breg *in vivo* or *in vitro* induction result in a distinct Breg lineage controlled by a specific set of transcription factors, or indeed whether different Breg populations are generated depending on the mode of stimulation and immunological microenvironment. Since IL-10⁺ B cells have been identified at all developmental stages of B cells, but only a fraction of each subset will express IL-10 upon appropriate stimulation, this suggests that a specific gene module within Breg precursors may exist [7]. The uncertainty that surrounds Breg identity and the lack of easily identifiable phenotypic markers beyond IL-10 may continue to hamper the progress of Breg cell therapy. Alternatively, protocols that could *ex vivo* expand B cells that are highly enriched for the expression of IL-10 and phenotypic markers associated with regulatory function, such as TIM-1, CD24 and CD38 [38], might suffice. To embark on such a strategy, the experimental approaches available that could be used clinically to generate human Bregs must be considered.

Adaptive signalling

As early as 2003, the Londei laboratory demonstrated the induction of mouse IL-10⁺ Bregs by stimulation of the CD154-CD40 pathway [39]. *In vitro* stimulation with an agonistic anti-CD40 monoclonal antibody (mAb) together with antigen (type II collagen) generated B cells capable of inhibiting Th1 differentiation, preventing and ameliorating collagen-induced arthritis (CIA) [39]. Short-term co-culture of agonistic CD40 mAb with mouse splenic B cells generated IL-10-producing Breg with a transitional 2 (T2) phenotype; this B cell subset could reverse autoimmunity *in vivo* upon injection and suppress Th1 responses in an IL-10-dependent manner [40]. In a study of mouse Breg expansion, the co-culture of mouse splenic B cells with NIH3T3 cells that expressed CD154 and BAFF, together with IL-4 and IL-21, resulted in greater than 4x10⁶ fold expansion of mouse B10 cells *ex vivo*, that were then able to inhibit autoimmune disease *in vivo*, in a model of experimental autoimmune encephalitis (EAE) [41]. Several studies have gone on to exploit this interaction to induce and expand human Bregs. Lemoine and Blair stimulated human B cells with membrane-bound CD154 to induce IL-10 production and Breg-mediated suppression *in vitro* [42, 43]. The involvement of the CD154-CD40 pathway highlights the importance of T cell-derived signals in IL-10⁺ Breg function. CD1d^{hi}CD5⁺ Breg express IL-10 and suppress inflammation in multiple immune-

related disorders upon *in vitro* 5-hour stimulation with LPS, phorbol 12-myristate 13-acetate (PMA), ionomycin and monensin, and are known as B10 cells [44, 45]. Using *CD40*^{-/-} and *MHC*^{-/-} mice in a model of experimental autoimmune encephalitis (EAE), it has been shown that the maturation of precursors to B10 cells ('B10pro') into functional IL-10-secreting Bregs was critically dependent on CD40 engagement and cognate interactions with T cells [41].

Antigen recognition is also important to Breg development and function. In MD4 mice, where the B cell receptor (BCR) has been fixed for an irrelevant antigen, Breg activation was impaired and Breg frequency reduced in mice that could not recover from EAE [45, 46]. Similar findings were observed in mice with decreased BCR signalling, such as in *CD19*^{-/-} mice in the context of inflammatory bowel disease (IBD) and contact hypersensitivity, whilst *CD19* overexpression expanded B10 cell numbers [44, 45, 47, 48]. Increasing CD40 signalling elevated B10 and B10pro cell numbers by at least 10-fold *in vivo* [49]. Moreover, the calcium sensors, stromal interaction molecule 1 (STIM-1) and STIM-2, which are involved in facilitating intracellular calcium influx following antigen recognition by the BCR, were required for B cell IL-10 production [50]. BCR signalling pathways may drive the capacity of B cells to express IL-10, thereby selecting for antigen-specific Breg *in vivo*. The requirement for Breg antigen specificity in transplantation was indicated when the adoptive transfer of mouse TIM-1⁺ Bregs from tolerant recipients of H2^d islet allografts prolonged survival of H2^d grafts but not those of 3rd party H2^k grafts [3, 51]. In a series of elegant experiments, tolerised C57BL/6 mice with ovalbumin (OVA)-specific BCR were used to demonstrate tolerance in the presence of OVA⁺ skin grafts, but not when the allograft lacked a recognizable antigen (BM12 skin grafts from MHC class II variant C57BL/6 mice). In contrast, wild-type C57BL/6 mice could accept BM12 skin grafts, which survived long-term. Antigen-specificity and recognition by the BCR therefore appears to be crucial for Breg function.

Innate Signalling

Whilst adaptive signals are important for Breg development and induction, innate signals also induce and expand IL-10⁺ Bregs. CpG oligonucleotides can stimulate mouse and human B10 cells to produce IL-10 [52, 53]. LPS can induce mouse B10pro cells to mature and become functional B10 cells [45]. Similarly, CD24^{hi}CD38^{hi} B cells can be expanded with IFN α and CpG *ex vivo* and retain IL-10 expression, CD24^{hi}CD38^{hi} phenotype, and *in vitro* suppressive function [54]. TLR signals, such as those that stimulate TLRs -2, -4 and -9, and apoptotic cells can expand B10 cell numbers *in vivo* [24, 55, 56]. MyD88 is a universal adaptor protein downstream of TLR. In the context of EAE, CD1d^{hi}CD5⁺ B cells from MyD88^{-/-} mice produced lower levels of IL-10 in response to LPS, PMA and ionomycin stimulation when compared to wild-type mice. MyD88-generated signals were critical for normal IL-10 producing, CD1d^{hi}CD5⁺ B10 cell development [45]. Moreover, a selective deficiency of MyD88 in B cells improved control of *Salmonella typhimurium* bacterial replication and prolonged survival of infected mice. B cells of MyD88-sufficient mice expressed IL-10 and could suppress neutrophil, NK cell and effector T cell activation, leaving mice more

susceptible to infection [57]. The release of IL-10 by TLR-stimulated B cells was increased upon CD40 engagement, suggesting that *in vivo*, optimal production of IL-10 occurred after TLR-activated B cells interacted with CD154-expressing CD4⁺ T cells [58, 59]. Activation of other molecules such as TIM-1, a phosphatidylserine receptor, resulted in the induction of IL-10⁺ Breg and transplant tolerance *in vivo*. B cells that expressed a mutant form of TIM-1 lacking the mucin domain, exhibited decreased phosphatidylserine binding and were unable to produce IL-10 in response to either apoptotic cells or ligation with anti-TIM-1 [60]. It has been hypothesised that both innate and adaptive Breg exist. Alternatively, the same Breg could respond to different stimuli to exert suppressive mechanisms including the secretion of IL-10 [61].

Cytokine-driven induction

Cytokines are involved in both adaptive and innate immune pathways and have proven critical to the induction and expansion of Breg subsets. The *in vitro* co-culture of mouse B cells with exogenous IL-21 and no other stimulation for 48-72 hours, is able to induce up to a x5.3 fold increase in mouse B10 frequencies [41]. Rosser et al. characterised Breg induction by pro-inflammatory cytokines IL-1 β and IL-6, as a method of auto-regulation in CIA [62]. In support of this view, Bregs increase in number during the inflammatory phase of several autoimmune disorders; in their absence, mice develop exacerbated disease [39, 53, 63, 64]. However, the immunosuppressive cytokine IL-35 can also induce IL-10⁺ and IL-35⁺ Breg, which profoundly suppresses inflammation in mouse models of autoimmunity [65, 66]. Mice in which only B cells did not express IL-35, demonstrated enhanced survival after *Salmonella* infection. IL-10 has been shown as critical to the generation of mouse Breg in early experiments [39], with more recent work also supporting an autocrine role for IL-10 in Breg function [67].

Clinical Protocols

The lessons learnt from experimental models and human studies could be used to generate a GMP-suitable IL-10⁺ Breg expansion or enrichment protocol (Fig.2). It is of course worth remembering that some of the experimental manipulations in animal models are potentially dependent on undefined and disease-specific factors which may be relevant only to that experimental context. Those mechanisms that appear to be consistent across multiple disease processes are more likely to result in replicable clinical protocols. These strategies may involve a combination of ligand-based stimuli including CD40- and TIM-1-ligation, and innate signals to stimulate the TLR. Soluble factors such as IL-21, IL-6, IL-10 and BAFF could be included as adjuncts to optimise IL-10⁺ Breg and improve efficiency. Following expansion, a standardised approach to detect IL-10-expression should be decided upon prior to *in vivo* administration: whether this is a short, 5 hour stimulation using LPS, PMA and ionomycin as advocated when identifying human B10 cells [53], or whether an additional preliminary step such as 48-72 hour stimulation with LPS, CpG or CD40 is required [43]. Given the phenotypic uncertainties surrounding Breg, it is difficult to draw the line between methods which can induce Breg that express IL-10, and those that can be subsequently used

to detect Breg by IL-10 expression. Furthermore, work by Cherukuri and colleagues has demonstrated that a distinguishing feature of human Breg was not only the capacity to express IL-10 upon stimulation, but also to express reduced levels of TNF α [28]. Any manufactured IL-10⁺ Breg therapy will need to be closely examined to ensure the reduction or absence of pro-inflammatory cytokine production.

Additional issues to consider include whether to use a human B cell population already enriched for IL-10⁺ Bregs, such as CD24^{hi}CD38^{hi} transitional, CD24^{hi}CD27⁺ memory or TIM-1⁺ B cells, or whether to stimulate whole CD19⁺ B cells as the initial substrate (Fig. 2). Whilst the proportional precursor frequencies of IL-10⁺ Breg are greater in circulating B cell subsets, they are rare in peripheral blood and so absolute starting numbers prior to expansion will be small [38, 43, 53, 68]. Since IL-10⁺ B cells can be generated from multiple B cell sub-populations by exploiting similar signalling pathways, it may be pragmatic for clinical use to initially isolate larger numbers of whole CD19⁺ B cells, and then subject the entire population to an optimised protocol for IL-10⁺ Breg induction and expansion.

Antigen specificity should also be considered. Polyclonal Treg cell therapy is being used in clinical trials with promising effect, but preclinical work has demonstrated the improved efficiency of alloantigen-specific Treg. In a humanised mouse model of alloimmune-mediated injury of human skin grafts, the infusion of alloantigen-specific Tregs resulted in a significant reduction in dermal tissue injury when compared to polyclonal Tregs [69, 70]. Antigen-specific B cells have a low precursor frequency and are difficult to isolate and expand in high yields [71]. The advent of chimeric antigen receptors (CAR) is revolutionising T cell therapy, and could facilitate development of an antigen-specific IL-10⁺ Breg therapy [72]. Constructs that include components of CD19, CD40, BCR and even the TLR could be combined for a potent Breg-CAR design (Fig. 2). Antigen-specific cell therapy has obvious advantages, including the reduction in off-target effects of immunosuppression such as opportunistic infection and cancer, which indeed are a source of serious morbidity in the transplant recipient. Increased potency will also encourage lower dose regimens, and the likelihood of an improved safety profile.

Amongst these exciting possibilities, a robust framework is needed to evaluate the functional stability and antibody-producing capacity of any such product prior to administration in the patient. As with mouse models of EAE, models of transplantation tolerance have highlighted the divergent roles for pro-inflammatory and regulatory B cells. For example, B cells can positively or negatively regulate allograft rejection depending on the nature of the allograft and the intensity of the rejection response [73]. Evidence supports the role of regulatory plasma cells [26, 66, 74], but given the detrimental consequences of ABMR, it will be vital to the safety and success of any Breg cell therapy to confidently interrogate its role in humoral immunity.

Pharmacological manipulation

The importance of Bregs in transplantation suggests that Breg promotion *in vivo* would benefit long-term allograft outcomes. Whilst we have explored the possibility of *ex vivo*-generated Breg cellular therapy infusion, we will now address those agents that may support *in vivo* expansion (Fig.3).

Existing therapies in transplantation

As discussed earlier, maintenance immunosuppressants such as prednisolone and azathioprine may have an effect on circulating IL-10⁺ Breg [16, 17]. Sirolimus may also support the *in vivo* and *in vitro* expansion of IL-10⁺ Bregs as well as FoxP3⁺ Tregs in liver transplantation [75].

The administration of several monoclonal antibody induction therapies has also been associated with Breg induction and/or expansion in renal transplantation. Alemtuzumab, a depleting anti-CD52 mAb, reduced the risk of biopsy-proven acute rejection in a broad range of human kidney transplant recipients, when compared to basiliximab induction [76]. Our group also observed a transient increase in CD24^{hi}CD38^{hi} Breg in peripheral blood during the first year post-transplant in alemtuzumab-treated human kidney transplant recipients [77]. Similar increases in transitional B cell frequency have been reported during B cell repopulation following a single dose of rituximab in the peri-transplant period in human kidney transplant recipients [78]. Intriguingly, a second dose of rituximab within 1–2 weeks of transplantation precipitated T-cell mediated rejection in kidney transplant recipients, and cardiac allograft vasculopathy in patients receiving a heart transplant [27, 79]. The dynamics of Breg repopulation following B cell depletion may be important in determining long-term allograft outcomes, in particular with regards to ABMR, and warrants further investigation.

A high serum concentration of the B cell survival factor, BAFF, in kidney transplant recipients is associated with the development of *de-novo* and increased pre-formed DSA levels, and increased ABMR rates [80, 81]. Administration of belimumab, an anti-BAFF mAb, to kidney transplant recipients, led to an increase in the B cell IL-10 to IL-6 ratio for the first 3 months post-transplant [82]. Despite a decrease in the frequency of transitional B cells, the belimumab group demonstrated a skewing of the IL-10 to IL-6 ratio in transitional and memory B-cell subsets, towards a more regulatory profile compared with controls. The belimumab group demonstrated lower frequencies of activated memory B cells, circulating plasmablasts and reduced formation of *de novo* IgG to new antigenic specificities post-transplant. The B cell compartment may have been diverted from a humoral response to a regulatory one. Longer-term outcomes from such trials will help to inform ABMR rates and the success of such an approach.

Treatment with belatacept, a CTLA-4-Ig fusion protein, is associated with lower BAFF levels and the persistent elevation in circulating transitional B cell frequencies in kidney transplant recipients [83]. Seven-year follow-up of kidney transplant recipients, comparing belatacept to cyclosporine-based immunosuppression, demonstrated higher patient and graft survival as well as increased mean eGFR in the belatacept group [84]. Ten-year follow-up analysis of transplant biopsies revealed increased frequencies of IL-10⁺ Breg, FoxP3⁺ Treg and

plasmacytoid dendritic regulatory cells in the allografts of belatacept group [85]. *In vitro* experiments have shown that belatacept induces STAT3 activation in stimulated B cells, thus providing a potential mechanistic explanation for the reported *in vivo* effects upon IL-10⁺ Breg [86]. The promotion of the regulatory immune compartment may contribute to better overall outcomes observed with belatacept, and illustrate the likely important interplay between different regulatory cell networks.

A recent Phase I clinical trial of modified immune cells (MIC) in renal transplantation exemplified such *in vivo* regulatory cell interactions [80]. Pre-clinical studies have demonstrated that monocytes, when treated with mitomycin C and matured into DCs, take on the morphology and phenotype of early immature DCs. This MIC population could strongly suppress T cell responses and induce donor-specific immunosuppression in experimental models [87, 88]. Kidney transplant recipients were infused with donor-derived MIC pre-transplant, and a control group retrospectively selected to match maintenance regimens. No DSAs or rejection episodes were noted in the MIC group. Allograft function was preserved and circulating Breg frequencies increased from day-30 post-transplant, despite steroid use. At the last follow-up time-point (median 810 days; range 720-1080 days), Breg frequencies were 24 times higher than levels in transplanted controls. Treg frequencies however, remained unchanged. Whilst the numbers of patients were small, the differences in Breg frequencies between the MIC and control groups were striking. Analysis of IGKV4-1 and IGKV1D-13, the ITN tolerance signature, revealed increased expression levels in the high dose MIC group when compared to lower dose groups. Importantly, the patients in the high-dose MIC group demonstrated absent donor-specific T cell responses at day 360 post-transplant, but preserved third-party responses, unlike control patients.

Longer-term outcomes and larger trials using these approaches will provide further insights as to whether such strategies can offer a Breg-targeted therapy that promotes B cell-mediated regulation and which reduces donor-specific humoral immunity.

Novel therapeutic targets

Whilst IL-10 expression is critical to both identification and function of most Breg subsets, no IL-10 therapy has passed clinical phase III studies [89]. A better understanding of Breg intracellular signaling pathways may lead to the development of novel agents that can target Breg function at a molecular level.

The cell surface molecule, TIM-1, plays an important role in IL-10⁺ Breg function and has been suggested as an almost universal marker of this subset [4]. IL-10⁺ Breg have been induced by TIM-1-ligand using the low affinity antibody, RMT10, and TIM-1⁺IL-10⁺ Breg could induce tolerance to islet allografts in an IL-10-dependent fashion. Mouse B cells expressing the TIM-1^{Δmucin} mutation were also unable to produce IL-10 in response to RMT10-ligation. Furthermore, TIM-1^{Δmucin} B cells displayed increased production of pro-inflammatory cytokines IL-1 and IL-6, promoted Th1 and Th17 responses, inhibited the generation of Treg and Tr1 cells and enhanced the severity of EAE [60, 90]. TIM-1 signalling

may play a pivotal role in Breg function, and so TIM-1 ligation may provide a therapeutic ‘B cell switch.’ Given that TIM-1 is also expressed as a stimulatory ligand by activated CD4⁺ T cells [91], a B-cell specific approach to TIM-1 ligation would probably be required to prevent off-target effects.

Many studies have demonstrated the importance of STAT3 phosphorylation in IL-10⁺ Breg function [26, 38, 41, 43, 65]. Selective targeting of this pathway may provide a coherent approach to promote Breg function. Hypoxia-inducible factor (HIF)-1 α , the oxygen-sensitive subunit of the HIF transcriptional complex, was recently shown to be important for IL-10 production in a STAT3-dependent manner by CD1d^{hi}CD5⁺ Breg in murine CIA and EAE models [92]. BCR-ligation increased the expression of HIF-1 α by B cells, which was suppressed in the presence of ERK or STAT3 inhibition. Crucially, pSTAT3 was found to bind the *Hif1 α* promoter, which was enhanced with BCR stimulation [92]. B cell-specific deletion of HIF-1 α reduced CD1d^{hi}CD5⁺ Breg proliferation, decreased IL-10 expression and lowered levels of glucose uptake and lactate secretion, revealing an important role for glycolysis in Breg expansion. Co-immunoprecipitation assays revealed two hypoxia responsive elements within the *il10* promoter to which HIF-1 α , HIF-1 β and pSTAT3 could bind; HIF-1 α and pSTAT3 could bind together upon BCR-stimulation. Thus BCR activation may induce IL-10 production through formation of HIF-1 α /pSTAT3 complexes at the *il10* promoter. Activating the HIF-1 α axis through pharmacological agents may provide tools to augment Breg function.

Work examining IL-10 signaling within Bregs has revealed roles for the c-Maf transcription factor and miRNA-146a [93, 94]. LPS stimulation of B220⁺ B cells results in increased c-Maf expression in mice, with c-Maf directly binding a Maf recognition element within the *il10* promoter. c-Maf knockdown results in decreased IL-10 production in collagen-induced arthritis (CIA) mice [94]. miRNA-146a deletion resulted in the development of autoimmune syndrome and a reduced frequency of CD1d^{hi}CD5⁺IL-10⁺ Breg in mice. Interestingly, *miRNA-146*^{-/-} B cells exhibited almost total abrogation of TIM-1 expression. Given the role of TIM-1 in IL-10 expression within Breg, miRNA-146 may influence Breg development and function through the regulation of TIM-1 expression [93].

Attention has also turned to negative regulators of IL-10 within Bregs, such as transcription factors including NFATc1 and histone deacetylases (HDAC). B cell-specific *Nfatc1* knockout resulted in amelioration of murine psoriasis in an IL-10-dependent manner. By binding to HDAC1 and the *Il10* gene, NFATc1 suppressed B cell IL-10 expression [95]. Shao et al. reported higher levels of HDAC11 and lower levels of IL-10 expression in B cells from allergic rhinitis patients when compared to healthy controls [96]. Introduction of HDAC11 shRNA significantly attenuated murine allergic rhinitis, and HDAC11-mediated suppression of IL-10 expression was elicited through HDAC-mediated deacetylation of c-Maf, which reduced c-Maf binding to the *il10* locus. Small molecule inhibitors of HDACs have received clinical approval for use in a range of diseases and so may present a rapidly translatable avenue to promote Breg function *in vivo*, in transplantation.

The use of mAbs directed against pro-inflammatory cytokines such as TNF α (anti-TNF α therapy) and IL-6 (anti-IL-6R mAb, tocilizumab) in the treatment of autoimmune disease has resulted in an increased frequency of human Breg in peripheral blood in association with improved clinical outcomes within 3-6 months of treatment [97, 98]. An observational study of patients with rheumatoid arthritis also reported that a high baseline level of IL-10⁺ Breg appeared to be predictive of a clinical response at 3 months upon anti-TNF α treatment [99]. These observations reinforce the importance of the expression of pro-inflammatory cytokines relative to that of immunosuppressive IL-10 in the context of Breg function. Work in transplantation has also demonstrated that whilst neutralisation of IL-10 inhibited CD24^{hi}CD38^{hi} Breg-mediated suppression of autologous Th1 cytokine expression, TNF- α blockade increased the suppressive capacity of both memory and naïve B-cell subsets [28]. The translation of such clinical observations from autoimmune disease may offer additional strategies to enhance endogenous Breg function and induction/expansion in transplantation.

Our rapidly expanding appreciation of Breg at transcriptional and post-transcriptional levels offers new approaches to target this subset's potent regulatory role. The introduction of novel small molecule agents and mAbs that can redirect B cell function away from humoral responses and towards regulatory roles may improve long-term clinical outcomes in transplantation.

Future Perspectives

Bregs present an exciting avenue by which to approach long term allograft tolerance, and to switch B cell responses from the humoral to the immunosuppressive. Evidence of the benefit of Bregs in transplantation has been steadily growing in animal models and human studies. The strategies discussed in this review provide a framework to continue to build on in order to eventually translate existing knowledge to clinical practice. Central to these efforts will be the identification of Breg markers for standardisation of generation protocols and monitoring of cellular therapy, as well as a greater appreciation of Breg humoral properties. Breg-specific signatures of tolerance and predictors of rejection could help to dictate immunosuppressant dose reduction, prevent immunological allograft damage, and minimise morbidity. Novel and existing reagents may also be prioritised to generate Bregs *in vivo*.

1. Geissler EK. The ONE Study compares cell therapy products in organ transplantation: introduction to a review series on suppressive monocyte-derived cells. *Transplant Res.* 2012; **1**: 11.
2. Sawitzki B, Harden PN, Reinke P, Moreau A, Hutchinson JA et al. The ONE Study: Evaluation of Regulatory Cell Therapy in Kidney Transplantation Using a Harmonized Trial Design. *Lancet.* 2020; **In Press**.
3. Lee KM, Kim JI, Stott R, et al. Anti-CD45RB/anti-TIM-1-induced tolerance requires regulatory B cells. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons.* 2012; **12**: 2072-8.
4. Ding Q, Yeung M, Camirand G, et al. Regulatory B cells are identified by expression of TIM-1 and can be induced through TIM-1 ligation to promote tolerance in mice. *The Journal of clinical investigation.* 2011; **121**: 3645-56.
5. Chesneau M, Michel L, Dugast E, et al. Tolerant Kidney Transplant Patients Produce B Cells with Regulatory Properties. *J Am Soc Nephrol.* 2015; **26**: 2588-98.
6. Asare A, Kanaparthi S, Lim N, et al. B Cell Receptor Genes Associated With Tolerance Identify a Cohort of Immunosuppressed Patients With Improved Renal Allograft Graft Function. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons.* 2017; **17**: 2627-39.
7. Mohib K, Cherukuri A, Rothstein DM. Regulatory B cells and transplantation: almost prime time? *Current opinion in organ transplantation.* 2018; **23**: 524-32.
8. Rosser EC, Mauri C. Regulatory B cells: origin, phenotype, and function. *Immunity.* 2015; **42**: 607-12.
9. Newell KA, Adams AB, Turka LA. Biomarkers of operational tolerance following kidney transplantation - The immune tolerance network studies of spontaneously tolerant kidney transplant recipients. *Human immunology.* 2018; **79**: 380-7.
10. Dudley JT, Tibshirani R, Deshpande T, Butte AJ. Disease signatures are robust across tissues and experiments. *Molecular systems biology.* 2009; **5**: 307.
11. Newell KA, Asare A, Kirk AD, et al. Identification of a B cell signature associated with renal transplant tolerance in humans. *The Journal of clinical investigation.* 2010; **120**: 1836-47.
12. Sagoo P, Perucha E, Sawitzki B, et al. Development of a cross-platform biomarker signature to detect renal transplant tolerance in humans. *The Journal of clinical investigation.* 2010; **120**: 1848-61.
13. Chesneau M, Pallier A, Braza F, et al. Unique B cell differentiation profile in tolerant kidney transplant patients. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons.* 2014; **14**: 144-55.
14. Pallier A, Hillion S, Danger R, et al. Patients with drug-free long-term graft function display increased numbers of peripheral B cells with a memory and inhibitory phenotype. *Kidney international.* 2010; **78**: 503-13.
15. Rebollo-Mesa I, Nova-Lamperti E, Mobillo P, et al. Biomarkers of Tolerance in Kidney Transplantation: Are We Predicting Tolerance or Response to Immunosuppressive Treatment? *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons.* 2016.

16. Bottomley MJ, Chen M, Fuggle S, Harden PN, Wood KJ. Application of Operational Tolerance Signatures Are Limited by Variability and Type of Immunosuppression in Renal Transplant Recipients: A Cross-Sectional Study. *Transplantation direct*. 2017; **3**: e125.
17. Rebollo-Mesa I, Nova-Lamperti E, Mobillo P, et al. Biomarkers of Tolerance in Kidney Transplantation: Are We Predicting Tolerance or Response to Immunosuppressive Treatment? *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons*. 2016; **16**: 3443-57.
18. Newell KA, Asare A, Sanz I, et al. Longitudinal studies of a B cell-derived signature of tolerance in renal transplant recipients. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons*. 2015; **15**: 2908-20.
19. Lozano JJ, Pallier A, Martinez-Llordella M, et al. Comparison of transcriptional and blood cell-phenotypic markers between operationally tolerant liver and kidney recipients. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons*. 2011; **11**: 1916-26.
20. Baron D, Ramstein G, Chesneau M, et al. A common gene signature across multiple studies relate biomarkers and functional regulation in tolerance to renal allograft. *Kidney international*. 2015; **87**: 984-95.
21. Hricik DE, Formica RN, Nickerson P, et al. Adverse Outcomes of Tacrolimus Withdrawal in Immune-Quiescent Kidney Transplant Recipients. *Journal of the American Society of Nephrology : JASN*. 2015; **26**: 3114-22.
22. Dugast E, Souillou JP, Foucher Y, et al. Failure of Calcineurin Inhibitor (Tacrolimus) Weaning Randomized Trial in Long-Term Stable Kidney Transplant Recipients. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons*. 2016; **16**: 3255-61.
23. Silva HM, Takenaka MC, Moraes-Vieira PM, et al. Preserving the B-cell compartment favors operational tolerance in human renal transplantation. *Molecular medicine*. 2012; **18**: 733-43.
24. Maseda D, Smith SH, DiLillo DJ, et al. Regulatory B10 cells differentiate into antibody-secreting cells after transient IL-10 production in vivo. *Journal of immunology*. 2012; **188**: 1036-48.
25. Kleffel S, Vergani A, Tezza S, et al. Interleukin-10+ regulatory B cells arise within antigen-experienced CD40+ B cells to maintain tolerance to islet autoantigens. *Diabetes*. 2015; **64**: 158-71.
26. de Masson A, Bouaziz JD, Le Buanec H, et al. CD24(hi)CD27(+) and plasmablast-like regulatory B cells in human chronic graft-versus-host disease. *Blood*. 2015; **125**: 1830-9.
27. Clatworthy MR, Watson CJ, Plotnek G, et al. B-cell-depleting induction therapy and acute cellular rejection. *N Engl J Med*. 2009; **360**: 2683-5.
28. Cherukuri A, Rothstein DM, Clark B, et al. Immunologic human renal allograft injury associates with an altered IL-10/TNF-alpha expression ratio in regulatory B cells. *Journal of the American Society of Nephrology : JASN*. 2014; **25**: 1575-85.
29. Cherukuri A, Salama AD, Carter CR, et al. Reduced human transitional B cell T1/T2 ratio is associated with subsequent deterioration in renal allograft function. *Kidney international*. 2017; **91**: 183-95.

30. Zhou H, Zhan F, Zhang H, et al. The proportion of CD19(+)CD24(hi)CD27(+) regulatory B cells predicts the occurrence of acute allograft rejection in liver transplantation. *Ann Transl Med.* 2019; **7**: 465.
31. Piloni D, Morosini M, Magni S, et al. Peripheral CD19+CD24(high)CD38(high) B-regulatory cells in lung transplant recipients. *Transpl Immunol.* 2019; **57**: 101245.
32. Svachova V, Sekerkova A, Hrubá P, et al. Dynamic changes of B-cell compartments in kidney transplantation: lack of transitional B cells is associated with allograft rejection. *Transplant international : official journal of the European Society for Organ Transplantation.* 2016; **29**: 540-8.
33. Sarvaria A, Basar R, Mehta RS, et al. IL-10+ regulatory B cells are enriched in cord blood and may protect against cGVHD after cord blood transplantation. *Blood.* 2016; **128**: 1346-61.
34. Zeng Q, Ng YH, Singh T, et al. B cells mediate chronic allograft rejection independently of antibody production. *The Journal of clinical investigation.* 2014; **124**: 1052-6.
35. Shiu KY, McLaughlin L, Rebollo-Mesa I, et al. Graft dysfunction in chronic antibody-mediated rejection correlates with B-cell-dependent indirect antidonor alloresponses and autocrine regulation of interferon-gamma production by Th1 cells. *Kidney international.* 2017; **91**: 477-92.
36. Shiu KY, McLaughlin L, Rebollo-Mesa I, et al. B-lymphocytes support and regulate indirect T-cell alloreactivity in individual patients with chronic antibody-mediated rejection. *Kidney international.* 2015; **88**: 560-8.
37. Shiu KY, Stringer D, McLaughlin L, et al. Effect of Optimized Immunosuppression (Including Rituximab) on Anti-Donor Alloresponses in Patients With Chronically Rejecting Renal Allografts. *Frontiers in immunology.* 2020; **11**: 79.
38. Aravena O, Ferrier A, Menon M, et al. TIM-1 defines a human regulatory B cell population that is altered in frequency and function in systemic sclerosis patients. *Arthritis research & therapy.* 2017; **19**: 8.
39. Mauri C, Gray D, Mushtaq N, Londei M. Prevention of arthritis by interleukin 10-producing B cells. *The Journal of experimental medicine.* 2003; **197**: 489-501.
40. Blair PA, Chavez-Rueda KA, Evans JG, et al. Selective targeting of B cells with agonistic anti-CD40 is an efficacious strategy for the generation of induced regulatory T2-like B cells and for the suppression of lupus in MRL/lpr mice. *Journal of immunology.* 2009; **182**: 3492-502.
41. Yoshizaki A, Miyagaki T, DiLillo DJ, et al. Regulatory B cells control T-cell autoimmunity through IL-21-dependent cognate interactions. *Nature.* 2012; **491**: 264-8.
42. Lemoine S, Morva A, Youinou P, Jamin C. Human T cells induce their own regulation through activation of B cells. *Journal of autoimmunity.* 2011; **36**: 228-38.
43. Blair PA, Norena LY, Flores-Borja F, et al. CD19(+)CD24(hi)CD38(hi) B cells exhibit regulatory capacity in healthy individuals but are functionally impaired in systemic Lupus Erythematosus patients. *Immunity.* 2010; **32**: 129-40.
44. Yanaba K, Bouaziz JD, Haas KM, Poe JC, Fujimoto M, Tedder TF. A regulatory B cell subset with a unique CD1dhiCD5+ phenotype controls T cell-dependent inflammatory responses. *Immunity.* 2008; **28**: 639-50.

45. Yanaba K, Bouaziz JD, Matsushita T, Tsubata T, Tedder TF. The development and function of regulatory B cells expressing IL-10 (B10 cells) requires antigen receptor diversity and TLR signals. *Journal of immunology*. 2009; **182**: 7459-72.
46. Fillatreau S, Sweenie CH, McGeachy MJ, Gray D, Anderton SM. B cells regulate autoimmunity by provision of IL-10. *Nature immunology*. 2002; **3**: 944-50.
47. Matsushita T, Yanaba K, Bouaziz JD, Fujimoto M, Tedder TF. Regulatory B cells inhibit EAE initiation in mice while other B cells promote disease progression. *The Journal of clinical investigation*. 2008; **118**: 3420-30.
48. Yanaba K, Yoshizaki A, Asano Y, Kadono T, Tedder TF, Sato S. IL-10-producing regulatory B10 cells inhibit intestinal injury in a mouse model. *The American journal of pathology*. 2011; **178**: 735-43.
49. Poe JC, Smith SH, Haas KM, et al. Amplified B lymphocyte CD40 signaling drives regulatory B10 cell expansion in mice. *PloS one*. 2011; **6**: e22464.
50. Matsumoto M, Fujii Y, Baba A, Hikida M, Kurosaki T, Baba Y. The calcium sensors STIM1 and STIM2 control B cell regulatory function through interleukin-10 production. *Immunity*. 2011; **34**: 703-14.
51. Kimura S, Rickert CG, Kojima L, et al. Regulatory B cells require antigen recognition for effective allograft tolerance induction. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons*. 2020; **20**: 977-87.
52. Lykken JM, Candando KM, Tedder TF. Regulatory B10 cell development and function. *International immunology*. 2015.
53. Iwata Y, Matsushita T, Horikawa M, et al. Characterization of a rare IL-10-competent B-cell subset in humans that parallels mouse regulatory B10 cells. *Blood*. 2011; **117**: 530-41.
54. Menon M, Blair PA, Isenberg DA, Mauri C. A Regulatory Feedback between Plasmacytoid Dendritic Cells and Regulatory B Cells Is Aberrant in Systemic Lupus Erythematosus. *Immunity*. 2016; **44**: 683-97.
55. Horikawa M, Weimer ET, DiLillo DJ, et al. Regulatory B cell (B10 Cell) expansion during *Listeria* infection governs innate and cellular immune responses in mice. *Journal of immunology*. 2013; **190**: 1158-68.
56. Miles K, Heaney J, Sibinska Z, et al. A tolerogenic role for Toll-like receptor 9 is revealed by B-cell interaction with DNA complexes expressed on apoptotic cells. *Proceedings of the National Academy of Sciences of the United States of America*. 2012; **109**: 887-92.
57. Neves P, Lampropoulou V, Calderon-Gomez E, et al. Signaling via the MyD88 adaptor protein in B cells suppresses protective immunity during *Salmonella typhimurium* infection. *Immunity*. 2010; **33**: 777-90.
58. Gray D, Gray M, Barr T. Innate responses of B cells. *European journal of immunology*. 2007; **37**: 3304-10.
59. Lampropoulou V, Hoehlig K, Roch T, et al. TLR-activated B cells suppress T cell-mediated autoimmunity. *Journal of immunology*. 2008; **180**: 4763-73.
60. Yeung MY, Ding Q, Brooks CR, et al. TIM-1 signaling is required for maintenance and induction of regulatory B cells. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons*. 2015; **15**: 942-53.
61. Tedder TF. B10 cells: a functionally defined regulatory B cell subset. *Journal of immunology*. 2015; **194**: 1395-401.

62. Rosser EC, Oleinika K, Tonon S, et al. Regulatory B cells are induced by gut microbiota-driven interleukin-1beta and interleukin-6 production. *Nature medicine*. 2014; **20**: 1334-9.
63. Carter NA, Rosser EC, Mauri C. Interleukin-10 produced by B cells is crucial for the suppression of Th17/Th1 responses, induction of T regulatory type 1 cells and reduction of collagen-induced arthritis. *Arthritis research & therapy*. 2012; **14**: R32.
64. Carter NA, Vasconcellos R, Rosser EC, et al. Mice lacking endogenous IL-10-producing regulatory B cells develop exacerbated disease and present with an increased frequency of Th1/Th17 but a decrease in regulatory T cells. *Journal of immunology*. 2011; **186**: 5569-79.
65. Wang RX, Yu CR, Dambuza IM, et al. Interleukin-35 induces regulatory B cells that suppress autoimmune disease. *Nature medicine*. 2014; **20**: 633-41.
66. Shen P, Roch T, Lampropoulou V, et al. IL-35-producing B cells are critical regulators of immunity during autoimmune and infectious diseases. *Nature*. 2014; **507**: 366-70.
67. Kim HS, Lee JH, Han HD, et al. Autocrine stimulation of IL-10 is critical to the enrichment of IL-10-producing CD40(hi)CD5(+) regulatory B cells in vitro and in vivo. *BMB reports*. 2015; **48**: 54-9.
68. van de Veen W, Stanic B, Yaman G, et al. IgG4 production is confined to human IL-10-producing regulatory B cells that suppress antigen-specific immune responses. *The Journal of allergy and clinical immunology*. 2013; **131**: 1204-12.
69. Sagoo P, Ali N, Garg G, Nestle FO, Lechler RI, Lombardi G. Human regulatory T cells with alloantigen specificity are more potent inhibitors of alloimmune skin graft damage than polyclonal regulatory T cells. *Science translational medicine*. 2011; **3**: 83ra42.
70. Landwehr-Kenzel S, Issa F, Luu SH, et al. Novel GMP-compatible protocol employing an allogeneic B cell bank for clonal expansion of allospecific natural regulatory T cells. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons*. 2014; **14**: 594-606.
71. Boonyaratanakornkit J, Taylor JJ. Techniques to Study Antigen-Specific B Cell Responses. *Frontiers in immunology*. 2019; **10**: 1694.
72. Zhang Q, Lu W, Liang CL, et al. Chimeric Antigen Receptor (CAR) Treg: A Promising Approach to Inducing Immunological Tolerance. *Frontiers in immunology*. 2018; **9**: 2359.
73. DiLillo DJ, Griffiths R, Seshan SV, et al. B lymphocytes differentially influence acute and chronic allograft rejection in mice. *Journal of immunology*. 2011; **186**: 2643-54.
74. Matsumoto M, Baba A, Yokota T, et al. Interleukin-10-producing plasmablasts exert regulatory function in autoimmune inflammation. *Immunity*. 2014; **41**: 1040-51.
75. Song J, Du G, Chen W, et al. The advantage of Sirolimus in amplifying regulatory B cells and regulatory T cells in liver transplant patients. *European journal of pharmacology*. 2020; **869**: 172872.
76. Group CSC, Haynes R, Harden P, et al. Alemtuzumab-based induction treatment versus basiliximab-based induction treatment in kidney transplantation (the 3C Study): a randomised trial. *Lancet*. 2014; **384**: 1684-90.

77. Heidt S, Hester J, Shankar S, Friend PJ, Wood KJ. B cell repopulation after alemtuzumab induction-transient increase in transitional B cells and long-term dominance of naive B cells. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons*. 2012; **12**: 1784-92.
78. Kamburova EG, Koenen HJ, van den Hoogen MW, Baas MC, Joosten I, Hilbrands LB. Longitudinal analysis of T and B cell phenotype and function in renal transplant recipients with or without rituximab induction therapy. *PloS one*. 2014; **9**: e112658.
79. Starling RC, Armstrong B, Bridges ND, et al. Accelerated Allograft Vasculopathy With Rituximab After Cardiac Transplantation. *Journal of the American College of Cardiology*. 2019; **74**: 36-51.
80. Morath C, Schmitt A, Kleist C, et al. Phase I trial of donor-derived modified immune cell infusion in kidney transplantation. *The Journal of clinical investigation*. 2020.
81. Snanoudj R, Candon S, Roelen DL, et al. Peripheral B-cell phenotype and BAFF levels are associated with HLA immunization in patients awaiting kidney transplantation. *Transplantation*. 2014; **97**: 917-24.
82. Banham GD, Flint SM, Torpey N, et al. Belimumab in kidney transplantation: an experimental medicine, randomised, placebo-controlled phase 2 trial. *Lancet*. 2018; **391**: 2619-30.
83. Leibler C, Matignon M, Pilon C, et al. Kidney transplant recipients treated with belatacept exhibit increased naive and transitional B cells. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons*. 2014; **14**: 1173-82.
84. Vincenti F, Rostaing L, Grinyo J, et al. Belatacept and Long-Term Outcomes in Kidney Transplantation. *The New England journal of medicine*. 2016; **374**: 333-43.
85. Furuzawa-Carballeda J, Uribe-Uribe NO, Arreola-Guerra JM, et al. Tissue talks: immunophenotype of cells infiltrating the graft explains histological findings and the benefits of belatacept at 10 years. *Clinical and experimental immunology*. 2019; **197**: 250-61.
86. Leibler C, Thiolat A, Henique C, et al. Control of Humoral Response in Renal Transplantation by Belatacept Depends on a Direct Effect on B Cells and Impaired T Follicular Helper-B Cell Crosstalk. *J Am Soc Nephrol*. 2018; **29**: 1049-62.
87. Radu CA, Kiefer J, Horn D, et al. Mitomycin-C-treated peripheral blood mononuclear cells (PBMCs) prolong allograft survival in composite tissue allotransplantation. *The Journal of surgical research*. 2012; **176**: e95-e101.
88. Dittmar L, Mohr E, Kleist C, et al. Immunosuppressive properties of mitomycin C-incubated human myeloid blood cells (MIC) in vitro. *Human immunology*. 2015; **76**: 480-7.
89. Saxena A, Khosraviani S, Noel S, Mohan D, Donner T, Hamad AR. Interleukin-10 paradox: A potent immunoregulatory cytokine that has been difficult to harness for immunotherapy. *Cytokine*. 2015; **74**: 27-34.
90. Xiao S, Brooks CR, Sobel RA, Kuchroo VK. Tim-1 is essential for induction and maintenance of IL-10 in regulatory B cells and their regulation of tissue inflammation. *Journal of immunology*. 2015; **194**: 1602-8.
91. Curtiss M, Colgan J. The role of the T-cell costimulatory molecule Tim-1 in the immune response. *Immunologic research*. 2007; **39**: 52-61.

92. Meng X, Grotsch B, Luo Y, et al. Hypoxia-inducible factor-1alpha is a critical transcription factor for IL-10-producing B cells in autoimmune disease. *Nature communications*. 2018; **9**: 251.
93. Amrouche L, You S, Sauvaget V, et al. MicroRNA-146a-deficient mice develop immune complex glomerulonephritis. *Scientific reports*. 2019; **9**: 15597.
94. Liu M, Zhao X, Ma Y, Zhou Y, Deng M, Ma Y. Transcription factor c-Maf is essential for IL-10 gene expression in B cells. *Scandinavian journal of immunology*. 2018; **88**: e12701.
95. Alrefai H, Muhammad K, Rudolf R, et al. NFATc1 supports imiquimod-induced skin inflammation by suppressing IL-10 synthesis in B cells. *Nat Commun*. 2016; **7**: 11724.
96. Shao JB, Luo XQ, Wu YJ, et al. Histone deacetylase 11 inhibits interleukin 10 in B cells of subjects with allergic rhinitis. *Int Forum Allergy Rhinol*. 2018; **8**: 1274-83.
97. Banko Z, Pozsgay J, Gati T, Rojkovich B, Ujfalussy I, Sarmay G. Regulatory B cells in rheumatoid arthritis: Alterations in patients receiving anti-TNF therapy. *Clinical immunology*. 2017; **184**: 63-9.
98. Snir A, Kessel A, Haj T, Rosner I, Slobodin G, Toubi E. Anti-IL-6 receptor antibody (tocilizumab): a B cell targeting therapy. *Clinical and experimental rheumatology*. 2011; **29**: 697-700.
99. Daïen CI, Gailhac S, Mura T, et al. Regulatory B10 cells are decreased in patients with rheumatoid arthritis and are inversely correlated with disease activity. *Arthritis & rheumatology*. 2014; **66**: 2037-46.
100. Evans JG, Chavez-Rueda KA, Eddaoudi A, et al. Novel suppressive function of transitional 2 B cells in experimental arthritis. *Journal of immunology*. 2007; **178**: 7868-78.
101. Amu S, Saunders SP, Kronenberg M, Mangan NE, Atzberger A, Fallon PG. Regulatory B cells prevent and reverse allergic airway inflammation via FoxP3-positive T regulatory cells in a murine model. *The Journal of allergy and clinical immunology*. 2010; **125**: 1114-24 e8.
102. Gray M, Miles K, Salter D, Gray D, Savill J. Apoptotic cells protect mice from autoimmune inflammation by the induction of regulatory B cells. *Proceedings of the National Academy of Sciences of the United States of America*. 2007; **104**: 14080-5.
103. Bankoti R, Gupta K, Levchenko A, Stager S. Marginal zone B cells regulate antigen-specific T cell responses during infection. *Journal of immunology*. 2012; **188**: 3961-71.
104. Wang WW, Yuan XL, Chen H, et al. CD19+CD24hiCD38hiBregs involved in downregulate helper T cells and upregulate regulatory T cells in gastric cancer. *Oncotarget*. 2015; **6**: 33486-99.
105. Flores-Borja F, Bosma A, Ng D, et al. CD19+CD24hiCD38hi B cells maintain regulatory T cells while limiting TH1 and TH17 differentiation. *Science translational medicine*. 2013; **5**: 173ra23.
106. Tebbe B, Wilde B, Ye Z, et al. Renal Transplant Recipients Treated with Calcineurin-Inhibitors Lack Circulating Immature Transitional CD19+CD24hiCD38hi Regulatory B-Lymphocytes. *PLoS One*. 2016; **11**: e0153170.
107. Nova-Lamperti E, Fanelli G, Becker PD, et al. IL-10-produced by human transitional B-cells down-regulates CD86 expression on B-cells leading to inhibition of CD4+T-cell responses. *Sci Rep*. 2016; **6**: 20044.

108. Latorre I, Esteve-Sole A, Redondo D, et al. Calcineurin and mTOR inhibitors have opposing effects on regulatory T cells while reducing regulatory B cell populations in kidney transplant recipients. *Transpl Immunol.* 2016; **35**: 1-6.
109. Hasan MM, Thompson-Snipes L, Klintmalm G, et al. CD24(hi)CD38(hi) and CD24(hi)CD27(+) Human Regulatory B Cells Display Common and Distinct Functional Characteristics. *J Immunol.* 2019; **203**: 2110-20.
110. Boonpiyathad T, Meyer N, Moniuszko M, et al. High-dose bee venom exposure induces similar tolerogenic B-cell responses in allergic patients and healthy beekeepers. *Allergy.* 2017; **72**: 407-15.
111. Kim HS, Kim AR, Kim DK, et al. Interleukin-10-producing CD5+ B cells inhibit mast cells during immunoglobulin E-mediated allergic responses. *Sci Signal.* 2015; **8**: ra28.
112. Lee JH, Noh J, Noh G, Choi WS, Cho S, Lee SS. Allergen-specific transforming growth factor-beta-producing CD19+CD5+ regulatory B-cell (Br3) responses in human late eczematous allergic reactions to cow's milk. *J Interferon Cytokine Res.* 2011; **31**: 441-9.
113. Natarajan P, Singh A, McNamara JT, et al. Regulatory B cells from hilar lymph nodes of tolerant mice in a murine model of allergic airway disease are CD5+, express TGF-beta, and co-localize with CD4+Foxp3+ T cells. *Mucosal immunology.* 2012; **5**: 691-701.
114. Kim AR, Kim HS, Kim DK, et al. Mesenteric IL-10-producing CD5+ regulatory B cells suppress cow's milk casein-induced allergic responses in mice. *Sci Rep.* 2016; **6**: 19685.
115. Yu CR, Choi JK, Uche AN, Egwuagu CE. Production of IL-35 by Bregs is mediated through binding of BATF-IRF-4-IRF-8 complex to il12a and ebi3 promoter elements. *J Leukoc Biol.* 2018; **104**: 1147-57.
116. Moore-Connors JM, Kim HS, Marshall JS, Stadnyk AW, Halperin SA, Wang J. CD43-, but not CD43+, IL-10-producing CD1dhiCD5+ B cells suppress type 1 immune responses during Chlamydia muridarum genital tract infection. *Mucosal Immunol.* 2015; **8**: 94-106.
117. Yang X, Yang J, Chu Y, et al. T follicular helper cells and regulatory B cells dynamics in systemic lupus erythematosus. *PLoS One.* 2014; **9**: e88441.
118. Zhang Y, Li J, Zhou N, et al. The Unknown Aspect of BAFF: Inducing IL-35 Production by a CD5(+)CD1d(hi)FcgammaRIIb(hi) Regulatory B-Cell Subset in Lupus. *J Invest Dermatol.* 2017; **137**: 2532-43.
119. Li S, Li X, Yang M, et al. Identification of the Subsets of IL-10-Producing Regulatory B Cells in the Course of Tolerance Induction and Maintenance in Islet Allograft Transplantation. *Transplant Proc.* 2018; **50**: 3900-5.
120. Qin Y, Zhang M, Jiang RM, et al. B10 cells play a role in the immune modulation of pro- and anti-inflammatory immune responses in mouse islet allograft rejection. *Cell Immunol.* 2016; **310**: 184-92.
121. Shen P, Roch T, Lampropoulou V, et al. IL-35-producing B cells are critical regulators of immunity during autoimmune and infectious diseases. *Nature.* 2014; **507**: 366-70.
122. Xiao S, Brooks CR, Zhu C, et al. Defect in regulatory B-cell function and development of systemic autoimmunity in T-cell Ig mucin 1 (Tim-1) mucin domain-mutant mice. *P Natl Acad Sci USA.* 2012; **109**: 12105-10.

123. Xiao S, Brooks CR, Sobel RA, Kuchroo VK. Tim-1 Is Essential for Induction and Maintenance of IL-10 in Regulatory B Cells and Their Regulation of Tissue Inflammation. *The Journal of Immunology*. 2015; **194**: 1602-8.
124. Kimura S, Rickert CG, Kojima L, et al. Regulatory B cells require antigen recognition for effective allograft tolerance induction. *Am J Transplant*. 2019.
125. Lee KM, Stott RT, Zhao G, et al. TGF-beta-producing regulatory B cells induce regulatory T cells and promote transplantation tolerance. *European journal of immunology*. 2014; **44**: 1728-36.
126. Kristensen B, Hegedus L, Lundy SK, Brimnes MK, Smith TJ, Nielsen CH. Characterization of Regulatory B Cells in Graves' Disease and Hashimoto's Thyroiditis. *PLoS One*. 2015; **10**: e0127949.
127. Griffin DO, Rothstein TL. Human "orchestrator" CD11b(+) B1 cells spontaneously secrete interleukin-10 and regulate T-cell activity. *Mol Med*. 2012; **18**: 1003-8.
128. Nakashima H, Hamaguchi Y, Watanabe R, et al. CD22 expression mediates the regulatory functions of peritoneal B-1a cells during the remission phase of contact hypersensitivity reactions. *J Immunol*. 2010; **184**: 4637-45.
129. Hsu LH, Li KP, Chu KH, Chiang BL. A B-1a cell subset induces Foxp3(-) T cells with regulatory activity through an IL-10-independent pathway. *Cell Mol Immunol*. 2015; **12**: 354-65.
130. Toubi E, Nussbaum S, Staun-Ram E, et al. Laquinimod modulates B cells and their regulatory effects on T cells in multiple sclerosis. *Journal of neuroimmunology*. 2012; **251**: 45-54.
131. Huber JE, Chang Y, Meinl I, Kumpfel T, Meinl E, Baumjohann D. Fingolimod Profoundly Reduces Frequencies and Alters Subset Composition of Circulating T Follicular Helper Cells in Multiple Sclerosis Patients. *Journal of immunology*. 2020; **204**: 1101-10.
132. Grutzke B, Hucke S, Gross CC, et al. Fingolimod treatment promotes regulatory phenotype and function of B cells. *Annals of clinical and translational neurology*. 2015; **2**: 119-30.
133. Salomon S, Guignant C, Morel P, et al. Th17 and CD24(hi)CD27(+) regulatory B lymphocytes are biomarkers of response to biologics in rheumatoid arthritis. *Arthritis research & therapy*. 2017; **19**: 33.

Table 1. IL-10⁺ Breg

Surface markers	Organism	Suppressive mechanism	Suppressive targets	Disease process	Activation protocol	Ref
CD21 ^{hi} CD23 ^{hi} CD24 ^{hi} IgM ^{hi} IgD ^{hi} CD1d ^{hi} (‘T2-MZP cells’)	Mouse	IL-10	T _H 17, T _H 1, Treg	Autoimmunity: CIA, lupus Allergy: Allergic airway inflammation	Collagen, PMA, iono; anti-CD40; <i>Schistosoma Mansoni</i> infection	[40, 100, 101]
CD21 ^{hi} CD23 ⁻ CD24 ^{hi} IgM ^{hi} IgD ^{lo} CD1d ^{hi} (‘MZ cells’)	Mouse	IL-10	CD4 ⁺ T cells, CD8 ⁺ T cells, Tr1	Autoimmunity: CIA Infection: <i>Leishmania Donavani</i>	AC; Anti-IgM & anti-CD21-coated beads; <i>Leishmania Donavani</i> infection	[102, 103]
CD24 ^{hi} CD38 ^{hi}	Human	IL-10, cell contact, TGF- β	T _H 1, Treg, T _H 17, pDC	Transplant: Lung, kidney, cGvHD Autoimmunity: SLE, RA Cancer: Gastric Healthy	PIB, CD40L; CpG, pDC; CpG, LPS, iono, PMA; CD154, PMA, iono; IL-10; Tumour environment; CpG, CD40L; CD40L, CpG, BCR ligation;	[23, 29, 31-33, 43, 54, 104-108]
CD24 ^{hi} CD27 ⁺	Human	IL-10, PD-L1, TGF- β , GrB, IL-6, TNF- α	Monocytes, T _H 1	Transplant: Liver, kidney, cGvHD Autoimmunity: SLE, RA, PSS, AVSD, MS,	CpG, PIB; CpG, CD40L; CpG, anti-BCR, CD40L; CpG, CD40L, PIB;	[26, 28, 30, 53, 109]
CD73 ⁻ CD25 ⁺ CD71 ⁺	Human	IL-10	CD4 ⁺ T cells	Allergy: Bee venom allergy Healthy	CpG; PMA, iono	[68, 110]
CD5 ⁺	Human Mouse	IL-10, TGF- β , IL-35	BMMCs, T _H 1, T _H 17, Treg	Autoimmunity: EAU Allergy: IgE-mediated allergic response, Late Eczematous Allergic Reactions to Cow's Milk, AAD, Casein-induced allergic response	IL-35, LPS; OVA; BSA, CD40L; Casein	[65, 111-114]
CD1d ^{hi} CD5 ⁺ (‘B10 cells’)	Human Mouse	IL-10, IL35	T _H 17, T _H 1, Tfh, Treg	Transplant: Islet Autoimmunity: EAU, EAE, SLE Infection: <i>Chlamydia muridarum</i> Allergy: CHS Healthy	Anti-CD45RB/Anti-TIM-1; BAFF; IL-21; LPS, PMA, iono; IRBP, anti-CD40 or PMA, iono; <i>Chlamydia muridarum</i> ; LPS, PIB, IL-21;	[41, 44, 115-120]
CD138 ⁺ CD138 ^{hi} TACI ⁺ CXCR4 ⁺ CD1d ^{int} Tim1 ^{int}	Human Mouse	GrB, cell contact, IL-35, IL-10	T _H 1, T _H 17, Macrophage, Neutrophil, NK cells	Transplant: Kidney Autoimmunity: EAE, EAU Infection: <i>Salmonella typhimurium</i>	CD40L, CpG; TLR-4, CD40L; PMA, iono; IL-2, IL-6, CpG, IFN- α or β ; <i>Salmonella typhimurium</i> ;	[5, 57, 74, 121]
TIM-1 ⁺	Human Mouse	IL-10, IL-4, TGF- β	T _H 1, T _H 17, Tr1, Treg	Transplant: Islet, skin, heart Autoimmunity: EAE, SSc, GD, HT Healthy	Anti-TIM-1 mAB (RMT1-10); AC, anti-IgM, anti-CD40, IL-21; CpG, PMA, iono, anti-IgG and IgM; CD45RB/Anti-TIM-1; PS; TG, CpG, PMA, iono	[4, 38, 60, 122-126]
CD11b ⁺	Human Mouse	IL-10; CD86	CD4 ⁺ T cells, Treg	Allergy: CHS Healthy	OVA; PMA, iono	[127-129]

Table 1. IL-10⁺ Breg Columns from left to right include the Breg phenotype, the organism in which they were identified, the proposed target of suppression, the disease model (where applicable) from which they were obtained, and Breg activation protocols. **Abbreviations:** AAD, allergic airway disease; AC, apoptotic cell; AVSD, Autoimmune vesiculobullous skin disease; BAFF, B cell activating factor; BCR, B cell receptor; BMMC, Bone marrow–derived mast cells; BSA, bovine serum albumin; cGVHD, chronic graft versus host disease; CHS, Contact hypersensitivity; CIA, Collagen-induced arthritis; CpG, CpG DNA; EAE, Experimental autoimmune encephalomyelitis; EAU, Experimental autoimmune uveitis; GD, Graves’ disease; GrB, granzyme B; HT, Hashimoto’s thyroiditis; IFN, Interferon; Ig, Immunoglobulin; iono, Ionomycin; IRBP, Interphotoreceptor retinoid-binding protein; LPS, Lipopolysaccharide; MS, Multiple sclerosis; MZ, Marginal zone; NK, natural killer; OVA, ovalbumin; pDC, plasmacytoid dendritic cell; PD-L1, Programmed death ligand 1; PIB, Phorbol 12-myristate 13-acetate, ionomycin and brefeldin A; PIM, phorbol 12-myristate 13-acetate, ionomycin, and monensin; PMA, Phorbol 12-myristate 13-acetate; PS, phosphatidyl serine; PSS, Primary Sjögren syndrome; RA, Rheumatoid arthritis; SLE, Systemic lupus erythematosus; SSc, systemic sclerosis; TG, thyroglobulin; TGF- β , Transforming growth factor beta; TIM-1, T-cell immunoglobulin and mucin domain 1; TLR, Toll like receptor; Tfh, T follicular helper cell; T2-MZP, Transitional 2-marginal zone precursor.

Figure.1. Breg-focused Immune Monitoring




Fig. 1. Breg-focused Immune Monitoring Breg signatures can be used in transplantation to improve graft and patient survival. Tolerance signatures have been identified and validated by consortia including the ITN [6, 18] and IoT [17]. Meta-analyses of tolerance studies also describe a B cell-specific signature [20]. Drug minimisation trials could be implemented in patients with predicted tolerance, alongside careful and long-term monitoring of graft function, DSA levels and protocol biopsies. Bregs have also been identified in rejection predictors [28, 29]. The tolerance signature genes and rejection tools in blue denote those associated with B cells and/or Bregs. Clinical utilisation of such tools may pre-empt rejection and facilitate accelerated anti-rejection treatment to improve graft survival and function within the context of dedicated trials.

Figure 2. Autologous Breg Cellular Therapy




Fig. 2. Autologous Breg Cellular Therapy Breg pre-clinical studies and clinical trials of other cell therapies in transplantation offer a template by which Breg cellular therapy could be introduced to the clinic. Autologous whole CD19⁺ B cells or B cell subsets enriched for IL-10⁺ Breg may be isolated from the potential organ recipient and expanded using known signalling pathways which promote Breg induction and/or generation *in vitro*. Antigen-specificity could be introduced by genetic engineering methods including lentiviral or CRISPR transfection. Breg- CAR constructs with intracellular components including those of the BCR, CD40, TIM-1 and TLR could generate antigen-specific, stable Breg cell therapy. Quality control measures should include *in vitro* evaluation of phenotype, IL-10 expression and capacity for STAT3 phosphorylation, prior to re-infusion into the recipient within a clinical trial.

Figure 3. Promotion of Bregs *in vivo*




Fig. 3. Promotion of Bregs *in vivo* Several induction agents and cell therapies have been demonstrated to support Breg survival and induction *in vivo* in transplantation. Multiple immune modulators in the context of autoimmune disease also promote Breg, offering additional therapeutic candidates [97, 98, 130-133]. Novel reagents in development may be able to soon offer specific Breg targeting. These reagents could be utilised to enhance Breg cell therapies both *in vivo* and during *in vitro* expansion protocols. Drug administration should be followed by careful monitoring of Breg-predominant tolerance signatures and predictors of rejection. Long-term graft outcomes and monitoring of humoral immunity will be critical to appreciating the benefits of such strategies.