

FROM BROWNIAN DYNAMICS TO POISSON-NERNST-PLANCK EQUATIONS: MULTI-RESOLUTION SIMULATIONS OF IONS*

JINYUAN ZHANG[†] AND RADEK ERBAN[†]

Abstract. The macroscopic description of Brownian dynamics (BD) particle-based models is often formulated as a system of partial differential equations (PDEs) that govern the evolution of particle concentrations in space and time. In certain applications, a BD-level of resolution is only necessary in specific regions of the computational domain, and the system can be efficiently simulated using multi-resolution (hybrid) approaches, which combine the detailed BD model in a part of the computational domain with coarser macroscopic equations in the remainder. When considering systems with charged particles (ions), their macroscopic PDEs can be expressed in the form of the Poisson-Nernst-Planck system. A multi-resolution approach, originally developed for electroneutral particles, is generalized to accommodate systems with ions. The strengths, limitations, and applicability of microscopic (BD), macroscopic (PDE), and multi-resolution simulation approaches are illustrated through a model consisting of a system of Na^+ and Cl^- ions, including both mobile and immobile charges, and periodic boundary conditions.

Key words. Brownian dynamics, Coulomb interactions, Poisson-Nernst-Planck system, multi-resolution simulations

MSC codes. 65C35,82C22,82C31,35J05,92C05

1. Introduction. At the molecular level, different classes of simulation methods can be applied to reaction-diffusion-advection processes in biological systems, including particle-based (individual-based) methods, which treat each particle separately, and coarse-grained methods, which describe particle concentrations [1]. Common coarse-grained approaches are written in terms of mean-field partial differential equations (PDEs), for example, the Poisson-Nernst-Planck (PNP) system, which is critical for understanding the dynamics of charged particles. The PNP system is particularly significant when examining the interactions between bulk and surface fixed charge [2, 3]. In biology, the PNP system has been used for analyzing ionic currents through protein channels embedded in membranes [4, 5, 6, 7].

There are two state variables in the PNP system, the charge density of different ionic species and the electrostatic potential. In Section 2, we will consider the system of two ionic species consisting of positive and negative ions in domain $\Omega \subset \mathbb{R}^3$. Denoting by $c^+ : \Omega \rightarrow [0, \infty)$ and $c^- : \Omega \rightarrow [0, \infty)$ the concentrations of the positive and negative ions, respectively, we obtain the Nernst-Planck equation by combining diffusive flux resulting from a concentration gradient with the flux arising from a potential gradient in the form

$$(1.1) \quad \frac{\partial c^\pm}{\partial t} = \nabla \cdot D^\pm \left(\nabla c^\pm + \frac{q^\pm}{k_b T} c^\pm \nabla \phi \right),$$

where ϕ is the electrostatic potential, k_b is the Boltzmann constant, T is the absolute temperature, and D^+ and q^+ (resp. D^- and q^-) is the diffusion constant and charge of the positive (resp. negative) ion. To get the PNP system, we couple equation (1.1)

*Submitted to the editors DATE.

Funding: This work was supported by the Engineering and Physical Sciences Research Council, grant number EP/V047469/1, awarded to Radek Erban. Jinyuan Zhang would like to thank Guangdong Postdoctoral Research Institute for studentship funding.

[†]Mathematical Institute, University of Oxford, Andrew Wiles Building, Radcliffe Observatory Quarter, Woodstock Road, Oxford OX2 6GG, United Kingdom

with the Poisson equation for the electrostatic potential

$$(1.2) \quad \nabla^2 \phi = -\frac{1}{\varepsilon_0 \varepsilon} [\varrho_p + q^+ c^+ + q^- c^-],$$

where ε is the dielectric permittivity constant of the medium, ε_0 is the dielectric permittivity constant of the vacuum and ϱ_p is the permanent charge density.

The PNP system (1.1)–(1.2) provides a mean-field description of a system of interacting charged particles. A more detailed description can be written using particle-based methods, such as molecular dynamics and Brownian dynamics (BD) simulations [1, 8]. Considering the system of N positive and N negative ions, and denoting their positions at time t by

$$(1.3) \quad \mathbf{X}_j^\pm(t) = [X_{j,1}^\pm(t), X_{j,2}^\pm(t), X_{j,3}^\pm(t)] \in \Omega, \quad \text{for } j = 1, 2, \dots, N,$$

where \mathbf{X}_j^+ (resp. \mathbf{X}_j^-) is the position of the j -th positive (resp. negative) ion, then the simplest BD model is given by the overdamped Langevin dynamics

$$(1.4) \quad d\mathbf{X}_j^\pm(t) = \sqrt{2D^\pm} d\mathbf{W}_j, \quad \text{for } j = 1, 2, \dots, N,$$

where \mathbf{W}_j is the vector of three independent Wiener processes. **The positions of particles calculated using equation (1.4) are then employed to describe their interactions, assuming that the particles interact if they are within a certain distance of each other** [1]. In computational studies, the BD equation (1.4) is commonly either discretized using fixed time step Δt [9, 10], or solved by using event-based algorithms, such as Green's function reaction dynamics [11], where the state of the system is updated by calculating the time at which the next interaction of particles occurs. The BD equation (1.4) provides a **coarse-grained description** of more detailed molecular dynamics, Langevin dynamics or stochastic coarse-grained models [12, 13], **while the BD modelling serves as the microscopic modelling in this manuscript.**

Although BD modelling based on equation (1.4) has previously been used for simulating ions in applications to calcium signalling [14, 15], the BD equation (1.4) must be generalized for use in other contexts. Considering applications to the dynamics of ions and ion channels [16, 17, 18, 19, 20], we generalize equation (1.4) by adding a drift term which includes a gradient of the (electric) potential. The resulting BD model is formulated in Section 2 as equation (2.3). This additional drift term depends not only on the position of the ion, but also on positions of other ions and on an external (electrical) field. In Section 3, we then show that our particle-based BD model converges in a suitable limit to the PNP system (1.1)–(1.2).

In Section 4, we present the details of computational implementations of both microscopic and macroscopic models. We discuss the strengths and limitations of each approach, applying them to an illustrative model comprising a system of Na^+ and Cl^- ions. In Section 5, we combine both approaches into a multi-resolution simulation framework, where BD simulations of individual ions and the mean-field PNP system (1.1)–(1.2) are used in different parts of the computational domain. **Such multi-resolution simulation frameworks have been developed in various contexts in the literature to simulate larger biological systems by using coarser (macroscopic) simulation approaches in parts of the computational domain [21, 22, 23, 24, 25]. Some of these studies couple BD based on equation (1.4) with coarser models given by reaction-diffusion PDEs [25, 23] or reaction-diffusion master equation [24]. In this case, the diffusion-limited interactions between particles are implemented whenever**

particles are at certain distances from each other, for example, particles react with a certain probability within the corresponding reaction radius [1]. While some other hybrid-continuum studies incorporate forces into the BD formulation as we will do in the next section, such generalizations have been restricted to one-dimensional examples [23, 26]. On the other hand, three-dimensional hybrid-continuum methods have been developed in other areas, for example, in rarefied gas dynamics, when continuum dynamics is given by the Navier-Stokes equations [27]. We conclude with the discussion of the applicability of the multi-resolution simulation techniques in Section 6.

2. Brownian dynamics. We model a system of N positive and N negative ions using BD in domain $\Omega \subset \mathbb{R}^3$. In our computational studies in Sections 4 and 5, the domain Ω is cuboid

$$(2.1) \quad \Omega = [0, L_1] \times [0, L_2] \times [0, L_3],$$

where L_1, L_2 and L_3 are positive parameters with units of length. The state of the system at time t is described by the $6N$ -dimensional vector

$$(2.2) \quad \bar{\mathbf{X}}(t) = [\mathbf{X}_1^+(t), \mathbf{X}_2^+(t), \dots, \mathbf{X}_N^+(t), \mathbf{X}_1^-(t), \mathbf{X}_2^-(t), \dots, \mathbf{X}_N^-(t)] \in \Omega^{2N},$$

where $\mathbf{X}_j^\pm(t)$ is given by (1.3) and denotes the position of the j -th positive ion (\mathbf{X}_j^+) or the j -th negative ion (\mathbf{X}_j^-) at time t for $j = 1, 2, \dots, N$. The time evolution of $\mathbf{X}_j^\pm(t)$ is given by the following Itô stochastic differential equation

$$(2.3) \quad d\mathbf{X}_j^\pm(t) = -\alpha^\pm \nabla_j^\pm U(\bar{\mathbf{X}}) dt + \sqrt{2D^\pm} d\mathbf{W}_j, \quad \text{for } j = 1, 2, \dots, N,$$

where α^+, α^-, D^+ and D^- are positive parameters, $U : \mathbb{R}^{6N} \rightarrow \mathbb{R}$ is the interaction potential and

$$\nabla_j^+ U = \left[\frac{\partial U}{\partial x_{j,1}^+}, \frac{\partial U}{\partial x_{j,2}^+}, \frac{\partial U}{\partial x_{j,3}^+} \right], \quad \nabla_j^- U = \left[\frac{\partial U}{\partial x_{j,1}^-}, \frac{\partial U}{\partial x_{j,2}^-}, \frac{\partial U}{\partial x_{j,3}^-} \right].$$

Our BD model (2.3) is a generalization of equation (1.4), where the interaction potential U is a function of $6N$ variables, because it depends not only on the position of the j -th positive or negative ion, but also on the positions of other ions in the system.

Our BD model (2.3) can be derived in the overdamped limit from the Langevin dynamics [1, 28]. In particular, the parameter α^\pm is the reciprocal of the corresponding friction coefficient, which has units of $[\text{mass}][\text{time}]^{-1}$. Since the interaction potential U has units of energy, we deduce that our BD equation (2.3) is dimensionally correct. However, to simplify our derivations of mean-field PNP equations, we first introduce the functional form of the interaction potential in Section 2.1 with a minimal number of parameters. In particular, our BD equation (2.3) can be considered non-dimensionalized. Our parameters are related to physical quantities in Section 2.2.

2.1. Functional form of the interaction potential. We assume that ions interact with fixed background potential and with each other. Moreover, we assume that our computational domain (2.1) is a small representative part of a larger domain that **contains ions**. Such a process can be modelled using suitable boundary conditions at the boundary $\partial\Omega$ of the cuboid. If the interactions were short-ranged, then ions leaving the domain Ω could be removed from the simulation, while additional ions could be introduced at the domain boundary $\partial\Omega$ according to a specified distribution [12, 29]. However, interactions between ions include long-range (Coulomb)

forces and the simulated ions (which are inside Ω) will also interact with ions which are outside of Ω and therefore not explicitly included in our simulation. To model such ions, we will implement standard periodic boundary conditions and assume that every ion interacts not only with the ions in Ω , but also with their periodic image copies and with the periodic image copies of the background potential. To simplify our presentation, we introduce the notation

$$(2.4) \quad \mathbf{L}_\ell = [\ell_1 L_1, \ell_2 L_2, \ell_3 L_3], \quad \text{for any integer valued vector } \ell = [\ell_1, \ell_2, \ell_3]$$

and we also denote summations over such integer valued vectors as:

$$(2.5) \quad \sum_{\ell} = \sum_{\ell_1=-\infty}^{\infty} \sum_{\ell_2=-\infty}^{\infty} \sum_{\ell_3=-\infty}^{\infty}$$

Denoting a point in the $6N$ -dimensional state space by

$$(2.6) \quad \bar{\mathbf{x}} = [\mathbf{x}_1^+, \mathbf{x}_2^+, \dots, \mathbf{x}_N^+, \mathbf{x}_1^-, \mathbf{x}_2^-, \dots, \mathbf{x}_N^-],$$

the interaction potential $U(\bar{\mathbf{x}})$ is given as the following sum of five terms

$$(2.7) \quad U(\bar{\mathbf{x}}) = \sum_{\ell} \sum_{i=1}^N \left(U_0^+(\mathbf{x}_i^+ - \mathbf{L}_\ell) + U_0^-(\mathbf{x}_i^- - \mathbf{L}_\ell) + \sum_{j=1}^{i-1} U_1(\|\mathbf{x}_i^+ - \mathbf{x}_j^+ - \mathbf{L}_\ell\|) \right. \\ \left. + \sum_{j=1}^N U_2(\|\mathbf{x}_i^- - \mathbf{x}_j^+ - \mathbf{L}_\ell\|) + \sum_{j=1}^{i-1} U_3(\|\mathbf{x}_i^- - \mathbf{x}_j^- - \mathbf{L}_\ell\|) \right),$$

where $U_0^\pm : \mathbb{R}^3 \rightarrow \mathbb{R}$ is a background potential which includes interactions with fixed charges in the environment, $\|\cdot\|$ denotes the standard Euclidean norm, potential $U_1 : [0, \infty) \rightarrow \mathbb{R}$ (resp. $U_3 : [0, \infty) \rightarrow \mathbb{R}$) describes distance-dependent interactions between two positive ions (resp. two negative ions) and $U_2 : [0, \infty) \rightarrow \mathbb{R}$ describes interactions between a positive and a negative ion. The distance-dependent potentials U_1 , U_2 and U_3 have the same functional form given as the sum of the Lennard-Jones and Coulomb potentials. We write them as

$$(2.8) \quad U_k(r) = \frac{A_k}{r^{12}} - \frac{B_k}{r^6} + \frac{C_k}{r},$$

where A_k , B_k and C_k , for $k = 1, 2, 3$, are parameters, which we relate to physical quantities in Section 2.2. We note that the sum in the third term of the interaction potential (2.7) does not include the case $i = j$, which is missing for $\ell = [0, 0, 0]$, because the i -th ion does not interact with itself, and for $\ell \neq [0, 0, 0]$ because the corresponding terms of the potential (2.7) would be constant and they would not change the drift term of our BD model (2.3), if we explicitly included them in (2.7). In our illustrative computational simulations in Section 4, we will implement the potential (2.7) using the Ewald summation, which is described in Section 4.2.

Considering r to be the distance between interacting particles with separation vector \mathbf{x} , we have $r = |\mathbf{x}|$ and equation (2.8) implies that $\nabla^2 U_k = \mathcal{H}_k$ where

$$(2.9) \quad \mathcal{H}_k(r) = \frac{132 A_k}{r^{14}} - \frac{30 B_k}{r^8} - \frac{C_k \delta(r)}{r^2} = \frac{132 A_k}{r^{14}} - \frac{30 B_k}{r^8} - 4\pi C_k \delta^3(\mathbf{x}),$$

where $\delta(\cdot)$ is the Dirac delta function and $\delta^3(\mathbf{x}) = \delta(x_1) \delta(x_2) \delta(x_3)$ is the three-dimensional Dirac delta function.

parameter	value	reference
A_1	$1.194 \times 10^{-8} \text{ nm}^{12} \text{ kcal mol}^{-1}$	[30]
B_1	$4.732 \times 10^{-5} \text{ nm}^6 \text{ kcal mol}^{-1}$	[30]
A_2	$5.186 \times 10^{-7} \text{ nm}^{12} \text{ kcal mol}^{-1}$	[30]
B_2	$4.171 \times 10^{-4} \text{ nm}^6 \text{ kcal mol}^{-1}$	[30]
A_3	$1.150 \times 10^{-5} \text{ nm}^{12} \text{ kcal mol}^{-1}$	[30]
B_3	$2.627 \times 10^{-3} \text{ nm}^6 \text{ kcal mol}^{-1}$	[30]
$C_1 = -C_2 = C_3$	$4.235 \times 10^{-1} \text{ nm kcal mol}^{-1}$	[33] and equation (2.12)
α^+	$3.241 \times 10^{11} \text{ kg}^{-1} \text{ s}$	[32] and equation (2.10)
α^-	$4.936 \times 10^{11} \text{ kg}^{-1} \text{ s}$	[32] and equation (2.10)
D^+	$1.334 \times 10^9 \text{ nm}^2 \text{ s}^{-1}$	[32]
D^-	$2.032 \times 10^9 \text{ nm}^2 \text{ s}^{-1}$	[32]

TABLE 1

Parameters of the potentials U_1 , U_2 and U_3 given by (2.8), and parameters α^\pm and D^\pm in the case of the positive ion being Na^+ and the negative ion being Cl^- .

2.2. Parameterization. While our minimal set of parameters A_k , B_k and C_k , for $k = 1, 2, 3$, in (2.8) can be considered dimensionless for the purposes of the theoretical derivations presented in Section 3, they can also be related to physical quantities and parameterized by the values in the literature. In Sections 4 and 5, we present illustrative simulations for the case when the positive ion is Na^+ and the negative ion is Cl^- . The parameters for these simulations are given in Table 1. The Lennard-Jones parameters A_j and B_j are calculated using the values in the CHARMM force field [30]. The parameters α^+ and α^- are reciprocals of the corresponding friction coefficient, which can be estimated by different methods [31]. We use the Einstein-Smoluchowski relation

$$(2.10) \quad \alpha^\pm = \frac{D^\pm}{k_b T}$$

where D^+ (resp. D^-) is the diffusion constant of the positive (resp. negative) ion, k_b is the Boltzmann constant and T is the absolute temperature. In Table 1, we use the values of D^+ and D^- at temperature $T = 25^\circ\text{C}$ for sodium and chloride ions [32], respectively, and we calculate α^\pm by (2.10). Since C_k corresponds to the Coulomb term, we have

$$(2.11) \quad C_1 = \frac{(q^+)^2}{4\pi\epsilon_0\epsilon}, \quad C_2 = \frac{q^+q^-}{4\pi\epsilon_0\epsilon} \quad \text{and} \quad C_3 = \frac{(q^-)^2}{4\pi\epsilon_0\epsilon},$$

where q^+ (resp. q^-) is the charge of the positive (resp. negative) ion, ϵ_0 is the dielectric permittivity constant of the vacuum and ϵ is the dielectric permittivity constant of the medium. Since we consider monovalent ions Na^+ and Cl^- , we have $q^+ = e$ and $q^- = -e$ in formulas (2.11), where $e = 1.602 \times 10^{-19} \text{ C}$ is the elementary charge. Consequently, formulas (2.11) imply

$$(2.12) \quad C_1 = -C_2 = C_3 = \frac{e^2}{4\pi\epsilon_0\epsilon},$$

which we use in our illustrative simulations together with ε being the permittivity of water at temperature $T = 25^\circ\text{C}$ given as $\varepsilon = 78.41$ in the literature [33].

3. Derivation of the PNP system from the BD model. In Section 5, we study a multi-resolution computational approach, which replaces BD simulations in a part of the computational domain by the PNP system (1.1)–(1.2). In this section, we illustrate that the PNP system (1.1)–(1.2) can, in fact, be derived under some assumptions from the microscopic BD model given by equation (2.3). The PNP system has been previously derived in the literature starting from a microscopic model written in terms of the Langevin dynamics [28]. Alternatively, the PNP system can also be derived from the alterations in the free energy functional, which encompasses both the electrostatic free energy and the ideal component of the chemical potential [34, 35]. We start with the derivation of the Poisson equation (1.2) in Section 3.1, followed by the derivation of the Nernst-Planck equations (1.1) in Section 3.2.

3.1. Derivation of the Poisson equation. The interaction potential U in our formulation (2.3) of the BD model is a function of $6N$ variables given by (2.7), while the potential ϕ in the Poisson equation (1.2) is a function of three spatial coordinates. To derive the PNP system, we first rewrite the BD model (2.3) using potentials $\Phi^\pm : \mathbb{R}^3 \rightarrow \mathbb{R}$ that depend on three spatial variables and are parametrized by current positions of ions $\bar{\mathbf{X}}(t)$ given by (2.2). We define

$$\begin{aligned}\Phi^+(\mathbf{x}) &= \sum_{\ell} \left(U_0^+(\mathbf{x} - \mathbf{L}_\ell) + \sum_{i=1}^N \left(U_1(\|\mathbf{x} - \mathbf{X}_i^+ - \mathbf{L}_\ell\|) + U_2(\|\mathbf{x} - \mathbf{X}_i^- - \mathbf{L}_\ell\|) \right) \right), \\ \Phi^-(\mathbf{x}) &= \sum_{\ell} \left(U_0^-(\mathbf{x} - \mathbf{L}_\ell) + \sum_{i=1}^N \left(U_2(\|\mathbf{x} - \mathbf{X}_i^+ - \mathbf{L}_\ell\|) + U_3(\|\mathbf{x} - \mathbf{X}_i^- - \mathbf{L}_\ell\|) \right) \right).\end{aligned}$$

We note that the potential $\Phi^\pm(\mathbf{x})$ is singular for $\mathbf{x} = \mathbf{X}_j^\pm$, $j = 1, 2, \dots, N$, so we also define potentials $\Phi_j^\pm : \mathbb{R}^3 \rightarrow \mathbb{R}$ that remove this singularity and all terms depending on \mathbf{X}_j^\pm by

$$(3.1) \quad \Phi_j^+(\mathbf{x}) = \Phi^+(\mathbf{x}) - \sum_{\ell} U_1(\|\mathbf{x} - \mathbf{X}_j^+ - \mathbf{L}_\ell\|),$$

$$(3.2) \quad \Phi_j^-(\mathbf{x}) = \Phi^-(\mathbf{x}) - \sum_{\ell} U_3(\|\mathbf{x} - \mathbf{X}_j^- - \mathbf{L}_\ell\|).$$

Then our BD model (2.3) can be rewritten as

$$(3.3) \quad d\mathbf{X}_j^\pm(t) = -\alpha^\pm \nabla (\Phi_j^\pm(\mathbf{X}_j^\pm)) dt + \sqrt{2D^\pm} d\mathbf{W}_j,$$

for $j = 1, 2, \dots, N$. Using our notation (2.9), we observe that the potential $\Phi^\pm(\mathbf{x})$ satisfies the Poisson equation in the form

$$\begin{aligned}\nabla^2 \Phi^+(\mathbf{x}) &= \sum_{\ell} \mathcal{H}_0^+(\mathbf{x} - \mathbf{L}_\ell) + \sum_{i=1}^N \left(\mathcal{H}_1(\|\mathbf{x} - \mathbf{X}_i^+ - \mathbf{L}_\ell\|) + \mathcal{H}_2(\|\mathbf{x} - \mathbf{X}_i^- - \mathbf{L}_\ell\|) \right), \\ \nabla^2 \Phi^-(\mathbf{x}) &= \sum_{\ell} \mathcal{H}_0^-(\mathbf{x} - \mathbf{L}_\ell) + \sum_{i=1}^N \left(\mathcal{H}_2(\|\mathbf{x} - \mathbf{X}_i^+ - \mathbf{L}_\ell\|) + \mathcal{H}_3(\|\mathbf{x} - \mathbf{X}_i^- - \mathbf{L}_\ell\|) \right),\end{aligned}$$

where we have defined $\mathcal{H}_0^\pm = \nabla^2 U_0^\pm$. In particular, we have formally obtained the Poisson equation, which includes the current positions of all ions. To get equation (1.2)

for the electrostatic potential, we focus on the terms on the right-hand side, which correspond to the electrostatic (Coulomb) forces. The Coulomb terms are included in equation (2.9) as terms corresponding to constant C_k . We also assume that there is an underlying permanent charge density ρ_p which helps us to express \mathcal{H}_0^\pm as

$$(3.4) \quad \mathcal{H}_0^\pm(\mathbf{x}) = \frac{q^\pm \rho_p(\mathbf{x})}{\varepsilon_0 \varepsilon} + \mathcal{H}_{0,\text{LJ}}^\pm(\mathbf{x}),$$

where the extra term $\mathcal{H}_{0,\text{LJ}}^\pm(\mathbf{x})$ corresponds to possible short-range interactions, which include some Lennard-Jones terms in our illustrative simulations in Section 4. The term $\mathcal{H}_{0,\text{LJ}}^\pm(\mathbf{x})$ does not appear in the Poisson equation (3.5) below, but it is included in the error term (3.14) in Section 3.2.

Using our notation (2.6) for points in the $6N$ -dimensional state space, we define the electrostatic potential as three-dimensional vector field $\hat{\phi}(\cdot; \bar{\mathbf{y}}) : \mathbb{R}^3 \rightarrow \mathbb{R}$ parameterized by the $6N$ -dimensional state vector $\bar{\mathbf{y}} \in \Omega^{2N}$. It is given as the solution of the Poisson equation

$$(3.5) \quad \nabla^2 \hat{\phi}(\mathbf{x}; \bar{\mathbf{y}}) = -\frac{1}{\varepsilon_0 \varepsilon} \sum_{\ell} \left[\rho_p(\mathbf{x} - \mathbf{L}_\ell) + \sum_{i=1}^N \left(q^+ \delta^3(\mathbf{x} - \mathbf{y}_i^+ - \mathbf{L}_\ell) + q^- \delta^3(\mathbf{x} - \mathbf{y}_i^- - \mathbf{L}_\ell) \right) \right],$$

which has the form of the Poisson equation (1.2) with concentrations replaced by the sums of the Dirac delta functions. Substituting the state space position $\bar{\mathbf{X}}(t)$ of ions at time t for $\bar{\mathbf{y}}$, we obtain that $\hat{\phi}(\cdot; \bar{\mathbf{X}}(t)) : \mathbb{R}^3 \rightarrow \mathbb{R}$ is a three-dimensional vector field which captures all Coulomb terms in our BD potential (3.1)–(3.2).

Denoting a point in the $6N$ -dimensional state space by (2.6) and the $6N$ -dimensional volume differential by

$$(3.6) \quad d\bar{\mathbf{x}} = d\mathbf{x}_1^+ d\mathbf{x}_2^+ \dots d\mathbf{x}_N^+ d\mathbf{x}_1^- d\mathbf{x}_2^- \dots d\mathbf{x}_N^-,$$

we can equivalently describe our BD model (3.3) by the time dependent probability density function $p : \Omega^{2N} \times [0, \infty) \rightarrow [0, \infty)$ which is defined so that

$$p(\bar{\mathbf{x}}, t) d\bar{\mathbf{x}} \quad \text{is the probability that} \quad \bar{\mathbf{X}}(t) \in [\bar{\mathbf{x}}, \bar{\mathbf{x}} + d\bar{\mathbf{x}}).$$

Using our notation (2.6) for points in the $6N$ -dimensional state space, we further simplify our presentation by denoting

$$(3.7) \quad \tilde{\mathbf{x}}_j^+ = (\mathbf{x}_1^+, \mathbf{x}_2^+, \dots, \mathbf{x}_{j-1}^+, \mathbf{x}_{j+1}^+, \dots, \mathbf{x}_N^+, \mathbf{x}_1^-, \mathbf{x}_2^-, \dots, \mathbf{x}_N^-),$$

$$(3.8) \quad \tilde{\mathbf{x}}_j^- = (\mathbf{x}_1^+, \mathbf{x}_2^+, \dots, \mathbf{x}_N^+, \mathbf{x}_1^-, \mathbf{x}_2^-, \dots, \mathbf{x}_{j-1}^-, \mathbf{x}_{j+1}^-, \dots, \mathbf{x}_N^-),$$

that is, $\tilde{\mathbf{x}}_j^+ \in \Omega^{2N-1}$ (resp. $\tilde{\mathbf{x}}_j^- \in \Omega^{2N-1}$) is a projection of the $6N$ -dimensional state space on the $(6N - 3)$ -dimensional subspace which excludes the position of the j -th positive (resp. negative) ion, for $j = 1, 2, \dots, N$. Using notation (3.7)–(3.8), the concentration $c^\pm(\mathbf{x}, t)$ is given as the sum of marginal densities of each ion being at the position \mathbf{x} at time t . Since ions are indistinguishable, we can write the concentration $c^\pm(\mathbf{x}, t)$ as N times the marginal density of the first ion in the list, **obtained by integrating over the positions of the remaining $2N - 1$ ions**, i.e.

$$(3.9) \quad c^\pm(\mathbf{x}, t) = N \int_{\Omega^{2N-1}} p(\mathbf{x}, \tilde{\mathbf{x}}_1^\pm, t) d\tilde{\mathbf{x}}_1^\pm.$$

Multiplying equation (3.5) by $p(\bar{\mathbf{y}}, t)$ and integrating over $\bar{\mathbf{y}} \in \Omega^{2N}$, we obtain

$$(3.10) \quad \nabla^2 \phi(\mathbf{x}) = -\frac{1}{\varepsilon_0 \varepsilon} \sum_{\boldsymbol{\ell}} \left[\rho_p(\mathbf{x} - \mathbf{L}_{\boldsymbol{\ell}}) + q^+ c^+(\mathbf{x} - \mathbf{L}_{\boldsymbol{\ell}}) + q^- c^-(\mathbf{x} - \mathbf{L}_{\boldsymbol{\ell}}) \right],$$

where $\phi(\mathbf{x})$ is an averaged electrostatic potential give by

$$(3.11) \quad \phi(\mathbf{x}) = \int_{\Omega^{2N}} \hat{\phi}(\mathbf{x}; \bar{\mathbf{y}}) p(\bar{\mathbf{y}}, t) d\bar{\mathbf{y}}.$$

Since the summation (2.5) over integer valued vectors (2.4) periodically covers \mathbb{R}^3 , we can equivalently rewrite equation (3.10) as equation

$$(3.12) \quad \nabla^2 \phi(\mathbf{x}) = -\frac{1}{\varepsilon_0 \varepsilon} \left[\rho_p(\mathbf{x}) + q^+ c^+(\mathbf{x}) + q^- c^-(\mathbf{x}) \right],$$

which is solved in cuboid domain (2.1) with periodic boundary conditions. In particular, we observe that the averaged electrostatic potential (3.11) satisfies the Poisson equation (1.2) in domain Ω .

3.2. Derivation of the Nernst-Planck equations. At time t , the electrostatic potential corresponding to both ions and permanent charge density is given as $\hat{\phi}(\mathbf{x}; \bar{\mathbf{X}}(t))$, where $\bar{\mathbf{X}}(t)$ is the state of the system at time t and $\hat{\phi}(\mathbf{x}; \bar{\mathbf{y}})$ is the solution to the Poisson equation (3.5). When using this potential in the evolution equation (3.3) for the j -th positive or negative ion, we need to remove the potential corresponding to this ion. Therefore, we also define potentials

$$(3.13) \quad \hat{\phi}_j^{\pm}(\mathbf{x}; \tilde{\mathbf{y}}_j^{\pm}) = \hat{\phi}(\mathbf{x}; \bar{\mathbf{y}}) - \sum_{\boldsymbol{\ell}} \frac{q^{\pm}}{4\pi\varepsilon_0 \varepsilon} \frac{1}{\|\mathbf{x} - \mathbf{y}_j^{\pm} - \mathbf{L}_{\boldsymbol{\ell}}\|},$$

where we use notation (3.7)–(3.8) to indicate that the left hand side is parameterized by the $(6N-3)$ -dimensional vector $\tilde{\mathbf{y}}_j^{\pm}$. The potential $\hat{\phi}_j^{\pm}(\mathbf{x}; \tilde{\mathbf{y}}_j^{\pm})$ does not capture all terms of the interaction potential $U(\bar{\mathbf{x}})$ given in (2.7). To rewrite the BD model (3.3) using the averaged electrostatic potential $\phi(\mathbf{x})$, we define the error term as the difference

$$(3.14) \quad E_j^{\pm}(\mathbf{x}; \tilde{\mathbf{X}}_j^{\pm}) = \Phi_j^{\pm}(\mathbf{x}) - q^{\pm} \hat{\phi}_j^{\pm}(\mathbf{x}; \tilde{\mathbf{X}}_j^{\pm}),$$

where $\Phi_j^{\pm}(\mathbf{x}) \equiv \Phi_j^{\pm}(\mathbf{x}; \tilde{\mathbf{X}}_j^{\pm})$ is given by (3.1)–(3.2). Substituting (3.14) into equation (3.3), we reformulate our BD model as

$$(3.15) \quad d\mathbf{X}_j^{\pm}(t) = -\alpha^{\pm} \nabla \left(q^{\pm} \hat{\phi}_j^{\pm}(\mathbf{X}_j^{\pm}; \tilde{\mathbf{X}}_j^{\pm}) + E_j^{\pm}(\mathbf{X}_j^{\pm}; \tilde{\mathbf{X}}_j^{\pm}) \right) dt + \sqrt{2D^{\pm}} d\mathbf{W}_j,$$

for $j = 1, 2, \dots, N$. The time evolution of $p(\bar{\mathbf{x}}, t)$ is given by the Fokker-Planck equation [36, 37]

$$(3.16) \quad \begin{aligned} \frac{\partial p}{\partial t}(\bar{\mathbf{x}}, t) &= \sum_{j=1}^N \nabla_j^+ \cdot \left(D^+ \nabla_j^+ p + p \alpha^+ \nabla_j^+ \left(q^+ \hat{\phi}_j^+(\mathbf{x}_j^+; \tilde{\mathbf{x}}_j^+) + E_j^+(\mathbf{x}_j^+; \tilde{\mathbf{x}}_j^+) \right) \right) \\ &+ \sum_{j=1}^N \nabla_j^- \cdot \left(D^- \nabla_j^- p + p \alpha^- \nabla_j^- \left(q^- \hat{\phi}_j^-(\mathbf{x}_j^-; \tilde{\mathbf{x}}_j^-) + E_j^-(\mathbf{x}_j^-; \tilde{\mathbf{x}}_j^-) \right) \right). \end{aligned}$$

Multiplying by N , integrating over $\tilde{\mathbf{x}}_1^\pm$, and using (3.9), (3.11) and (2.10), we obtain

$$(3.17) \quad \frac{\partial c^\pm}{\partial t} = \nabla \cdot D^\pm \left(\nabla c^\pm + \frac{q^\pm}{k_b T} c^\pm \nabla \phi + \mathbf{f}_a \right),$$

where the additional term \mathbf{f}_a is given by

$$(3.18) \quad \begin{aligned} \mathbf{f}_a(\mathbf{x}, t) &= \frac{N}{k_b T} \int_{\Omega^{2N-1}} p(\mathbf{x}, \tilde{\mathbf{x}}_1^\pm, t) \nabla_1^\pm E_1^\pm(\mathbf{x}; \tilde{\mathbf{x}}_1^\pm) d\tilde{\mathbf{x}}_1^\pm \\ &+ \frac{Nq^\pm}{k_b T} \int_{\Omega^{2N-1}} \int_{\Omega^{2N}} p(\mathbf{x}, \tilde{\mathbf{x}}_1^\pm, t) p(\bar{\mathbf{y}}, t) \left(\nabla_1^\pm \hat{\phi}_1^\pm(\mathbf{x}; \tilde{\mathbf{x}}_1^\pm) - \nabla \hat{\phi}(\mathbf{x}; \bar{\mathbf{y}}) \right) d\bar{\mathbf{y}} d\tilde{\mathbf{x}}_1^\pm. \end{aligned}$$

Consequently, we can formally recover the Nernst-Planck equations (1.1) by assuming that $\mathbf{f}_a \equiv \mathbf{0}$ in equation (3.17). This assumption neglects short-range Lennard-Jones interactions. In particular, we cannot expect that the derived PNP system (1.1)–(1.2) will provide good description of the macroscopic behaviour in the regions of space where the excluded volume interactions significantly influence the system dynamics. For example, the PNP system (1.1)–(1.2) is not applicable in the narrowest parts of ion channels, or close to the biological structures of interest [16, 18, 19]. In the next section, we will introduce a model system, which will include the regions of space where the excluded volume interactions cannot be neglected.

4. Comparison of BD with the solution obtained by the PNP system.

In this section, we will implement both the microscopic and macroscopic descriptions of an illustrative model comprising a system of Na^+ and Cl^- ions. The macroscopic description is given by the PNP equations (1.1)–(1.2) derived in Section 3. This is followed in Section 5 by developing a multi-resolution scheme that uses both BD and PNP in the same dynamical simulation.

We consider a simulation of N sodium and N chloride ions in domain Ω defined by (2.1). In particular, we use the parameters given for Na^+ and Cl^- in Table 1. The permanent charge density ϱ_p , which appears in (1.2) or (3.4), is given in our example as regular rectangular lattices of permanent positive (on plane $x_1 = 0$) and negative charges (on plane $x_1 = L_1/2$). Since we consider periodic boundary conditions, these layers are repeated in all planes $x_1 = jL_1$ and $x_1 = (j + 1/2)L_1$, where j is an integer. Our domain is schematically shown in Figure 1(a), where we highlight the layer of permanent charges in planes $x_1 = 0$, $x_1 = L_1/2$ and $x_1 = L_1$. More precisely, given an integer value N_p , we have N_p^2 elementary charges on the side $x_1 = 0$ of the cuboid (2.1) at points

$$(4.1) \quad \mathbf{z}_{m,n}^+ = \left[0, \frac{(m-1/2)L_2}{N_p}, \frac{(n-1/2)L_3}{N_p} \right] \quad \text{for } m, n = 1, 2, \dots, N_p,$$

and we also have N_p^2 negative elementary charges on the side $x_1 = L_1/2$ of the cuboid (2.1) at points

$$(4.2) \quad \mathbf{z}_{m,n}^- = [L_1/2, 0, 0] + \mathbf{z}_{m,n}^+ \quad \text{for } m, n = 1, 2, \dots, N_p.$$

Therefore the permanent charge density is given as

$$(4.3) \quad \varrho_p(\mathbf{x}) = \sum_{m=1}^{N_p} \sum_{n=1}^{N_p} e \delta^3(\mathbf{x} - \mathbf{z}_{m,n}^+) - e \delta^3(\mathbf{x} - \mathbf{z}_{m,n}^-).$$




FIG. 1. (a) Schematic of the computational domain Ω given by (2.1) with periodic lattices of point charges on planes $x_1 = 0$, $x_1 = L_1/2$ and $x_1 = L_1$. We use $N_p = 4$. (b) The steady state solutions $c^-(x_1)$ and $c^+(x_1)$ obtained by one-dimensional approximation (4.5) for three different choices of N and N_p , namely: (i) $N = 100$ and $N_p = 6$ (solid lines); (ii) $N = 100$ and $N_p = 4$ (dashed lines); and (iii) $N = 50$ and $N_p = 4$ (dot-dashed lines).

In our model, we identify the elementary charges on the side $x_1 = 0$ with Na^+ ions at fixed positions (4.1) and the elementary charges on the side $x_1 = L/2$ with Cl^- ions at fixed positions (4.2). In particular, ions are repelled from the fixed charges at short distances by the corresponding Lennard-Jones type interactions, giving the Laplacian of the background potential (3.4) in the form

$$\mathcal{H}_0^+(\mathbf{x}) = \frac{q^+ \rho_p(\mathbf{x})}{\varepsilon_0 \varepsilon} + \sum_{m,n=1}^{N_p} \left(\frac{132 A_1}{\|\mathbf{x} - \mathbf{z}_{m,n}^+\|^{14}} - \frac{30 B_1}{\|\mathbf{x} - \mathbf{z}_{m,n}^+\|^8} + \frac{132 A_2}{\|\mathbf{x} - \mathbf{z}_{m,n}^-\|^{14}} - \frac{30 B_2}{\|\mathbf{x} - \mathbf{z}_{m,n}^-\|^8} \right),$$

$$\mathcal{H}_0^-(\mathbf{x}) = \frac{q^- \rho_p(\mathbf{x})}{\varepsilon_0 \varepsilon} + \sum_{m,n=1}^{N_p} \left(\frac{132 A_2}{\|\mathbf{x} - \mathbf{z}_{m,n}^+\|^{14}} - \frac{30 B_2}{\|\mathbf{x} - \mathbf{z}_{m,n}^+\|^8} + \frac{132 A_3}{\|\mathbf{x} - \mathbf{z}_{m,n}^-\|^{14}} - \frac{30 B_3}{\|\mathbf{x} - \mathbf{z}_{m,n}^-\|^8} \right),$$

where parameters A_1 and B_1 (resp. A_2 and B_2 , or A_3 and B_3) correspond to Na^+ - Na^+ (resp. Na^+ - Cl^- or Cl^- - Cl^-) interactions and are given in Table 1. The cuboid domain (2.1) is chosen with equal parameters $L_1 = L_2 = L_3 = 10$ nm, i.e. Ω is a cube and we implement periodic boundary conditions as discussed in Section 2. Our model will be investigated using BD simulations in Section 4.2 and by numerically solving the PNP system (1.1)–(1.2) in Section 4.3. We begin by discussing the equilibrium properties of our model, when the PNP system reduces to solving the Poisson-Boltzmann equation.

4.1. One-dimensional approximation. In Figure 1(a), permanent charges are placed on a rectangular lattice with spacing L_2/N_p and L_3/N_p . To get some insight into the model behaviour, we first approximate the layers of sodium and chloride ions at positions (4.1) and (4.2) as uniformly charged planes at $x_1 = 0$ and $x_1 = L_1/2$ with surface charge density σ^+ and σ^- , respectively, where

$$\sigma^\pm = \pm \frac{N_p^2 e}{L_2 L_3}.$$

In particular, the permanent charge density is independent of x_2 and x_3 directions. Using this simplifying assumption, we can develop a one-dimensional approximation

of our system, because c^\pm and ϕ are constant in x_2 and x_3 directions. Considering the equilibrium properties of our system, we can substitute

$$(4.4) \quad c^\pm(\mathbf{x}) = c^\pm(x_1) \quad \text{and} \quad \phi(\mathbf{x}) = \phi(x_1)$$

into the Nernst-Planck equation (1.1) and pass $t \rightarrow \infty$ to get

$$(4.5) \quad c^\pm(x_1) = c_0 \exp\left(-\frac{q^\pm \phi(x_1)}{k_b T}\right),$$

where c_0 is a constant given by the normalization condition

$$(4.6) \quad N = \int_0^{L_1} \int_0^{L_2} \int_0^{L_3} c^\pm(\mathbf{x}) dx_3 dx_2 dx_1 = L_2 L_3 \int_0^{L_1} c^\pm(x_1) dx_1.$$

Substituting (4.4) and (4.5) into the Poisson equation (1.2), and using $q^+ = e$ and $q^- = -e$, we get

$$(4.7) \quad \phi''(x_1) = \frac{e}{\varepsilon_0 \varepsilon} \left[2c_0 \sinh\left(\frac{e\phi(x_1)}{k_b T}\right) + \frac{N_p^2}{L_2 L_3} \sum_{j=-\infty}^{\infty} \delta(x_1 - (j+1/2)L_1) - \delta(x_1 - jL_1) \right].$$

Since $\phi(x_1)$ is an L_1 -periodic function and symmetric around $x_1 = L_1/2$, we can find it by solving equation (4.7) in interval $[0, L_1/2]$. This is equivalent to solving

$$(4.8) \quad \phi''(x_1) = \frac{2e c_0}{\varepsilon_0 \varepsilon} \sinh\left(\frac{e\phi(x_1)}{k_b T}\right) \quad \text{in} \quad x \in [0, L_1/2],$$

with boundary conditions

$$(4.9) \quad \phi'(0) = -\frac{N_p^2 e}{2\varepsilon_0 \varepsilon L_2 L_3}, \quad \phi'(L_1/2) = -\frac{N_p^2 e}{2\varepsilon_0 \varepsilon L_2 L_3},$$

and the normalization condition (4.6). The solutions are presented in Figure 1(b), where we plot c^- and c^+ given by (4.5) as a function of x_1 for three different choices of N and N_p , namely: (i) $N = 100$ and $N_p = 6$; (ii) $N = 100$ and $N_p = 4$; and (iii) $N = 50$ and $N_p = 4$. In our calculations based on solving equation (4.8), we assume that $N/2$ of mobile ions are in interval $[0, L_1/2]$. To visualize the results in Figure 1(b), we extend the calculated solution of (4.8) from $[0, L_1/2]$ to the whole interval $[0, L_1]$ by symmetry.

4.2. BD implementation. The equilibrium solution of macroscopic equations given by the Poisson-Boltzmann system (4.5)–(4.7) in Figure 1(b) provides the coarsest description of our model system with Na^+ and Cl^- ions. In this section, we will present an implementation of the most detailed (microscopic, individual-based) model which is given as a BD simulation of the system of N sodium and N chloride ions in the cuboid domain (2.1) with permanent charges at locations (4.1)–(4.2) and with periodic boundary conditions. In particular, the permanent charges are periodically repeated in all planes $x_1 = jL_1/2$, where j is an integer. To simulate the evolution of the BD model, we will choose a small time step Δt and we discretize equation (3.3) by using the Euler-Maruyama method. We get

$$(4.10) \quad \mathbf{X}_j^\pm(t + \Delta t) = \mathbf{X}_j^\pm(t) - \alpha^\pm \nabla(\Phi_j^\pm(\mathbf{X}_j^\pm)) \Delta t + \sqrt{2D^\pm \Delta t} \boldsymbol{\xi}_j,$$

for $j = 1, 2, \dots, N$, where the coordinates of vector $\boldsymbol{\xi}_j$ are sampled from a normal distribution with zero mean and unit variance and $\Phi_j^\pm(\mathbf{x}) \equiv \Phi_j^\pm(\mathbf{x}; \tilde{\mathbf{X}}_j^\pm)$ is given by (3.1)–(3.2). In our BD simulations based on equation (4.10), we need to calculate the potential gradient at every time step. This includes sums (2.5) over all integer valued vectors (2.4). To evaluate these sums efficiently, we first decompose the potential into two terms (3.14) and rewrite the BD equation (3.3) as (3.15). The first term $E_j^\pm(\mathbf{X}_j^\pm; \tilde{\mathbf{X}}_j^\pm)$ only contains short-range interactions and we can calculate it by replacing the sum of infinitely many terms (2.5) with the most important term in this sum, which corresponds to the copy of the point closest to \mathbf{X}_j^\pm , i.e. we use the minimum-image convention to approximate all short-range interactions in $E_j^\pm(\mathbf{X}_j^\pm; \tilde{\mathbf{X}}_j^\pm)$. Moreover, we denote the Euclidean distance $\|\cdot\|$ modified by the minimum-image convention by

$$(4.11) \quad \|\mathbf{z}_1 - \mathbf{z}_2\|_{\min} = \min_{\boldsymbol{\ell}} \|\mathbf{z}_1 - \mathbf{z}_2 - \mathbf{L}\boldsymbol{\ell}\| \quad \text{for any } \mathbf{z}_1, \mathbf{z}_2 \in \mathbb{R}^3.$$

The second term in (3.14), $q^\pm \hat{\phi}_j^\pm(\mathbf{X}_j^\pm; \tilde{\mathbf{X}}_j^\pm)$, can be expressed using $\hat{\phi}(\mathbf{x}; \bar{\mathbf{y}})$ that solves the Poisson equation (3.5). We divide it into two terms corresponding to the mobile and fixed, ions, respectively. We obtain

$$\hat{\phi}(\mathbf{x}; \bar{\mathbf{y}}) = \hat{\phi}_m(\mathbf{x}; \bar{\mathbf{y}}) + \hat{\phi}_p(\mathbf{x}),$$

where $\hat{\phi}_m(\mathbf{x}; \bar{\mathbf{y}})$ and $\hat{\phi}_p(\mathbf{x})$ solve the Poisson equations in the following forms

$$(4.12) \quad \nabla^2 \hat{\phi}_m(\mathbf{x}; \bar{\mathbf{y}}) = -\frac{1}{\varepsilon_0 \varepsilon} \sum_{\boldsymbol{\ell}} \left[\sum_{i=1}^N \left(q^+ \delta^3(\mathbf{x} - \mathbf{y}_i^+ - \mathbf{L}\boldsymbol{\ell}) + q^- \delta^3(\mathbf{x} - \mathbf{y}_i^- - \mathbf{L}\boldsymbol{\ell}) \right) \right]$$

and

$$(4.13) \quad \nabla^2 \hat{\phi}_p(\mathbf{x}) = -\frac{1}{\varepsilon_0 \varepsilon} \sum_{\boldsymbol{\ell}} \rho_p(\mathbf{x} - \mathbf{L}\boldsymbol{\ell}).$$

To solve (4.12), we will use the Ewald summation [38, 39]. We get

$$(4.14) \quad \begin{aligned} \hat{\phi}_m(\mathbf{x}; \bar{\mathbf{y}}) &= \frac{1}{4\pi\varepsilon_0 \varepsilon} \sum_{i=1}^N \left(q^+ \frac{\operatorname{erfc}(\beta|\mathbf{x} - \mathbf{y}_i^+|_{\min})}{|\mathbf{x} - \mathbf{y}_i^+|_{\min}} + q^- \frac{\operatorname{erfc}(\beta|\mathbf{x} - \mathbf{y}_i^-|_{\min})}{|\mathbf{x} - \mathbf{y}_i^-|_{\min}} \right) \\ &+ \frac{1}{\varepsilon_0 \varepsilon L_1 L_2 L_3} \sum_{\boldsymbol{\ell} \neq [0,0,0]} \frac{1}{|\mathbf{k}_{\boldsymbol{\ell}}|^2} \exp\left(-\frac{|\mathbf{k}_{\boldsymbol{\ell}}|^2}{4\beta^2}\right) \\ &\quad \times \sum_{i=1}^N \left(q^+ \exp(i\mathbf{k}_{\boldsymbol{\ell}} \cdot (\mathbf{x} - \mathbf{y}_i^+)) + q^- \exp(i\mathbf{k}_{\boldsymbol{\ell}} \cdot (\mathbf{x} - \mathbf{y}_i^-)) \right) \end{aligned}$$

where $\beta > 0$ is a parameter, $|\cdot|_{\min}$ denotes the distance modified by the minimum image convention (4.11) and

$$(4.15) \quad \mathbf{k}_{\boldsymbol{\ell}} = \left[\frac{2\pi\ell_1}{L_1}, \frac{2\pi\ell_2}{L_2}, \frac{2\pi\ell_3}{L_3} \right] \quad \text{for any integer valued vector } \boldsymbol{\ell} = [\ell_1, \ell_2, \ell_3].$$

Since the permanent charge density is given by (4.3), the equation (4.13) has the same functional form as the equation (4.12) with permanent charges replacing the

mobile charges. In particular, the potential $\widehat{\phi}_p(\mathbf{x})$ can be calculated by the Ewald summation formula (4.14) with permanent charges replacing the mobile charges, with some computations performed only once at the beginning of the simulation. We define

$$(4.16) \quad \theta_p(\boldsymbol{\ell}) = \frac{\exp(-|\mathbf{k}_\ell|^2/(4\beta^2))}{|\mathbf{k}_\ell|^2 \varepsilon_0 \varepsilon L_1 L_2 L_3} \sum_{m=1}^{N_p} \sum_{n=1}^{N_p} q^+ \exp(i\mathbf{k}_\ell \cdot \mathbf{z}_{m,n}^+) + q^- \exp(i\mathbf{k}_\ell \cdot \mathbf{z}_{m,n}^-),$$

where $\mathbf{z}_{m,n}^+$ and $\mathbf{z}_{m,n}^-$ are positions of permanent positive and negative charges given by (4.1) and (4.2), respectively. Our BD algorithm uses equation (4.10) at every time step, which requires calculating the gradient of (4.14). Let r be the distance between two ions with separation vector \mathbf{r} , i.e. $r = |\mathbf{r}|$. Differentiating (2.8) and the first term (real part) of the potential (4.14), we define (the real part of) the force term between the ions as

$$(4.17) \quad \mathbf{F}_k(\mathbf{r}) = \left(\frac{12 A_k}{r^{14}} - \frac{6 B_k}{r^8} + \frac{2\beta C_k \exp(-\beta^2 r^2)}{\sqrt{\pi} r^2} + \frac{C_k \operatorname{erfc}(\beta r)}{r^3} \right) \mathbf{r},$$

where $k = 1$, $k = 2$, and $k = 3$, correspond to interactions between two Na^+ ions, one Na^+ and one Cl^- ion, and two Cl^- ions, respectively.

One iteration of the BD algorithm (*i.e.* the update of the system from time t to time $t + \Delta t$) is described in Table 2 as Algorithm [A1]–[A6]. We denote the drift term $-\alpha^\pm \nabla(\Phi_j^\pm(\mathbf{X}_j^\pm))$ in equation (4.10) as $\mathbf{a}_j^\pm(t)$, *i.e.* we rewrite the discretized equation (4.10) as

$$(4.18) \quad \mathbf{X}_j^\pm(t + \Delta t) = \mathbf{X}_j^\pm(t) + \mathbf{a}_j^\pm(t) \Delta t + \sqrt{2D^\pm \Delta t} \xi_j,$$

for $j = 1, 2, \dots, N$, where drift term $\mathbf{a}_j^\pm(t)$ is calculated in steps [A1]–[A5]. In step [A1], we initialize the drift term using the components of forces corresponding to the permanent charges, putting

$$(4.19) \quad \mathbf{a}_j^+(t) := \alpha^+ \sum_{m=1}^{N_p} \sum_{n=1}^{N_p} \mathbf{F}_1((\mathbf{X}_j^+ - \mathbf{z}_{m,n}^+)_{\min}) + \mathbf{F}_2((\mathbf{X}_j^+ - \mathbf{z}_{m,n}^-)_{\min})$$

$$(4.20) \quad \mathbf{a}_j^-(t) := \alpha^- \sum_{m=1}^{N_p} \sum_{n=1}^{N_p} \mathbf{F}_2((\mathbf{X}_j^- - \mathbf{z}_{m,n}^+)_{\min}) + \mathbf{F}_3((\mathbf{X}_j^- - \mathbf{z}_{m,n}^-)_{\min})$$

where we substitute separation vectors modified by the minimum image convention (4.11) in the formula (4.17), using notation $(\cdot)_{\min}$ and $|\cdot|_{\min}$ to denote the corresponding modified vectors and norms, respectively. In step [A2], we add force terms corresponding to interactions between mobile Na^+ ions. This is followed in steps [A3] and [A4] by adding force terms corresponding to Na^+ – Cl^- and Cl^- – Cl^- interactions, respectively, between mobile ions. In step [A5], we calculate sum

$$(4.21) \quad \theta(\boldsymbol{\ell}) = \theta_p(\boldsymbol{\ell}) + \frac{\exp(-|\mathbf{k}_\ell|^2/(4\beta^2))}{|\mathbf{k}_\ell|^2 \varepsilon_0 \varepsilon L_1 L_2 L_3} \sum_{i=1}^N q^+ \exp(i\mathbf{k}_\ell \cdot \mathbf{X}_i^+) + q^- \exp(i\mathbf{k}_\ell \cdot \mathbf{X}_i^-),$$

which is needed to evaluate the Fourier part of the potential (4.14). It includes the precomputed term $\theta_p(\boldsymbol{\ell})$, given by (4.16), corresponding to the permanent charges. The summation in step [A5] is over all $\boldsymbol{\ell} \in \mathcal{L}(\kappa)$, where $\kappa \in \mathbb{N}$ is a parameter to be specified, and set $\mathcal{L}(\kappa)$ is defined by

$$(4.22) \quad \mathcal{L}(\kappa) = \left\{ \boldsymbol{\ell} = [\ell_1, \ell_2, \ell_3] \mid |\ell_1| \leq \kappa, |\ell_2| \leq \kappa, |\ell_3| \leq \kappa \text{ and } \boldsymbol{\ell} \neq [0, 0, 0] \right\}.$$

- [A1] Initialize $\mathbf{a}_j^\pm(t)$, for $j = 1, 2, \dots, N$, using formulas (4.19)–(4.20).
- [A2] For each pair of Na^+ ions, labelled i and j (for $i \neq j$), calculate force $\mathbf{F}_1(\mathbf{r})$ between them by (4.17), where $\mathbf{r} = (\mathbf{X}_i^+ - \mathbf{X}_j^+)_{\min}$. Put
- $$\mathbf{a}_i^+(t) := \mathbf{a}_i^+(t) + \alpha^+ \mathbf{F}_1((\mathbf{X}_i^+ - \mathbf{X}_j^+)_{\min})$$
- $$\mathbf{a}_j^+(t) := \mathbf{a}_j^+(t) - \alpha^+ \mathbf{F}_1((\mathbf{X}_i^+ - \mathbf{X}_j^+)_{\min})$$
- [A3] For each Na^+ ion, labelled i , and each Cl^- ion, labelled j , calculate force $\mathbf{F}_2(\mathbf{r})$ between them by (4.17), where $\mathbf{r} = (\mathbf{X}_i^+ - \mathbf{X}_j^-)_{\min}$. Put
- $$\mathbf{a}_i^+(t) := \mathbf{a}_i^+(t) + \alpha^+ \mathbf{F}_2((\mathbf{X}_i^+ - \mathbf{X}_j^-)_{\min})$$
- $$\mathbf{a}_j^-(t) := \mathbf{a}_j^-(t) - \alpha^+ \mathbf{F}_2((\mathbf{X}_i^+ - \mathbf{X}_j^-)_{\min})$$
- [A4] For each pair of Cl^- ions, labelled i and j (for $i \neq j$), calculate force $\mathbf{F}_3(\mathbf{r})$ between them by (4.17), where $\mathbf{r} = (\mathbf{X}_i^- - \mathbf{X}_j^-)_{\min}$. Put
- $$\mathbf{a}_i^-(t) := \mathbf{a}_i^-(t) + \alpha^- \mathbf{F}_3((\mathbf{X}_i^- - \mathbf{X}_j^-)_{\min})$$
- $$\mathbf{a}_j^-(t) := \mathbf{a}_j^-(t) - \alpha^- \mathbf{F}_3((\mathbf{X}_i^- - \mathbf{X}_j^-)_{\min})$$
- [A5] Calculate sum (4.21) for all $\ell \in \mathcal{L}(\kappa)$, given by (4.22), and put
- $$\mathbf{a}_j^+(t) := \mathbf{a}_j^+(t) + \alpha^+ q^+ \sum_{\ell \in \mathcal{L}(\kappa)} \text{Im}(\theta(\ell) \exp(i \mathbf{k}_\ell \cdot \mathbf{X}_j^+)) \mathbf{k}_\ell$$
- $$\mathbf{a}_j^-(t) := \mathbf{a}_j^-(t) + \alpha^- q^- \sum_{\ell \in \mathcal{L}(\kappa)} \text{Im}(\theta(\ell) \exp(i \mathbf{k}_\ell \cdot \mathbf{X}_j^-)) \mathbf{k}_\ell$$
- where $\text{Im}(\cdot)$ denotes the imaginary part of a complex number.
- [A6] Generate $6N$ coordinates of vectors $\boldsymbol{\xi}_j$, for $j = 1, 2, \dots, N$, as normally distributed numbers with zero mean and unit variance. Calculate the positions of mobile ions at time $t + \Delta t$ by using equation (4.18).

TABLE 2

One iteration of the BD algorithm to calculate the positions of ions at time $t + \Delta t$ given positions of ions at time t .

Finally, after we evaluate the drift term in steps [A1]–[A5], we use equation (4.18) to calculate the positions of mobile ions at time $t + \Delta t$ in step [A6].

In Figure 2, we present illustrative results calculated by Algorithm [A1]–[A6]. We use $N = 100$ mobile Na^+ ions and $N = 100$ mobile Cl^- ions in domain Ω given by (2.1) with $L_1 = L_2 = L_3 = 10$ nm and the permanent charges at locations (4.1)–(4.2) with $N_p = 6$. To apply our BD Algorithm [A1]–[A6], we need to specify parameters $\beta > 0$ and $\kappa \in \mathbb{N}$, which are used to implement the periodic boundary conditions using the Ewald summation. In our illustrative simulations, we choose $\kappa = 3$ in (4.22), which means that we evaluate $|\mathcal{L}(\kappa)| = (2\kappa + 1)^3 - 1 = 342$ terms in the sum in step [A5]. This truncation of the infinite sum introduces an error in the evaluation of the Fourier part of the Ewald sum. The choice of the parameter $\beta > 0$ can then be optimized in the way that the error of the real part of the Ewald sum (which dominates for small values of β) is comparable to the error of the Fourier part (which dominates for large values of β). Using $\kappa = 3$, we find the optimal value of β to be $\beta = 0.05$, which is used to obtain the results presented in Figure 2, where we visualize the spatial histogram (density of ions) obtained by running the long-time simulation of Algorithm [A1]–[A6] over 10^7 time steps of length $\Delta t = 10^{-2}$ ps. To enable direct comparison, with our one-dimensional results in Figure 1(b), we plot the calculated equilibrium density over the first coordinate, x_1 , integrating over the




FIG. 2. The equilibrium distributions calculated using Algorithm [A1]–[A6] for $N = 100$, $N_p = 6$, $\kappa = 3$ and $\beta = 0.05$ are visualized as gray histograms for (a) mobile Na^+ ions; and (b) mobile Cl^- ions. The results are compared with density c^\pm calculated for the same parameter values in Figure 1(b) by using a one-dimensional approximation of the PNP system, given by (4.7) with boundary conditions (4.9) (green lines). The red lines are calculated by numerically solving the three-dimensional PNP system (1.1)–(1.2).

x_2 and x_3 coordinates. We present this one-dimensional (marginal) distribution of mobile Na^+ ions in Figure 2(a) as the green histogram. We compare it with the results (blue line) calculated by the approximate one-dimensional theory developed in Section 4.1. We observe similar qualitative behaviour, but the results quantitatively differ. We confirm this observation in Figure 2(b) where we present the BD results for mobile Cl^- ions as the green histogram. There are a number of reasons behind the observed difference. One of them is that our solution to the PNP system in Section 4.1 is itself a one-dimensional approximation. We will next compare it with numerically solving the PNP system (1.1)–(1.2) in the three-dimensional domain. This numerical approach will also be useful for developing the multi-resolution method in Section 5.

4.3. Numerical method for solving macroscopic PNP equations. To solve the macroscopic PNP equations (1.1)–(1.2) in the three-dimensional cuboid domain (2.1), we use a cuboid mesh with $n_1 \times n_2 \times n_3$ mesh points, where $n_i \in \mathbb{N}$ for $i = 1, 2, 3$ and define

$$(4.23) \quad \Delta x_1 = \frac{L_1}{n_1}, \quad \Delta x_2 = \frac{L_2}{n_2} \quad \text{and} \quad \Delta x_3 = \frac{L_3}{n_3}.$$

The mesh points are at locations

$$(4.24) \quad \mathbf{x}_{i,j,k} = [(i - 1/2) \Delta x_1, (j - 1/2) \Delta x_2, (k - 1/2) \Delta x_3]$$

for $i = 1, 2, \dots, n_1$, $j = 1, 2, \dots, n_2$, $k = 1, 2, \dots, n_3$, and we denote

$$(4.25) \quad c_{i,j,k}^\pm(t) = c^\pm(\mathbf{x}_{i,j,k}, t), \quad \phi_{i,j,k}(t) = \phi(\mathbf{x}_{i,j,k}, t).$$

Using finite differences to discretize the right-hand side of equation (1.1), we obtain

$$(4.26) \quad \frac{dc_{i,j,k}^\pm}{dt} = \mathcal{A}_{i,j,k}(c^\pm, \phi)$$

where $\mathcal{A}_{i,j,k}(c^\pm, \phi)$ is defined by

$$\begin{aligned} \mathcal{A}_{i,j,k}(c^\pm, \phi) = & D^\pm \left(\frac{c_{i+1,j,k}^\pm - 2c_{i,j,k}^\pm + c_{i-1,j,k}^\pm}{\Delta x_1^2} + \frac{c_{i,j+1,k}^\pm - 2c_{i,j,k}^\pm + c_{i,j-1,k}^\pm}{\Delta x_2^2} \right. \\ & \left. + \frac{c_{i,j,k+1}^\pm - 2c_{i,j,k}^\pm + c_{i,j,k-1}^\pm}{\Delta x_3^2} \right) \\ & + \frac{D^\pm q^\pm}{2k_b T} \left(\frac{(c_{i+1,j,k}^\pm + c_{i,j,k}^\pm)(\phi_{i+1,j,k} - \phi_{i,j,k}) + (c_{i-1,j,k}^\pm + c_{i,j,k}^\pm)(\phi_{i-1,j,k} - \phi_{i,j,k})}{\Delta x_1^2} \right. \\ & + \frac{(c_{i,j+1,k}^\pm + c_{i,j,k}^\pm)(\phi_{i,j+1,k} - \phi_{i,j,k}) + (c_{i,j-1,k}^\pm + c_{i,j,k}^\pm)(\phi_{i,j-1,k} - \phi_{i,j,k})}{\Delta x_2^2} \\ & \left. + \frac{(c_{i,j,k+1}^\pm + c_{i,j,k}^\pm)(\phi_{i,j,k+1} - \phi_{i,j,k}) + (c_{i,j,k-1}^\pm + c_{i,j,k}^\pm)(\phi_{i,j,k-1} - \phi_{i,j,k})}{\Delta x_3^2} \right), \end{aligned}$$

with the convention that indices i, j, k are periodic with periods n_1, n_2 and n_3 , respectively. For example, $c_{n_1+1,j,k}^+$ is identified with $c_{1,j,k}^+$ to implement the periodic boundary conditions. To solve the PNP equations (1.1)–(1.2) over the time interval $[t, t + \delta t]$, we need to implement a suitable numerical method to discretize the system of ODEs (4.26) in time. In the multi-resolution approach in Section 5, we will use an explicit numerical method to calculate concentrations $c_{i,j,k}^\pm(t + \delta t)$ at time $t + \delta t$ given the concentrations $c_{i,j,k}^\pm(t)$ and potential $\phi_{i,j,k}(t)$ at time t , with the simplest approach being the forward Euler scheme

$$(4.27) \quad c_{i,j,k}^+(t + \delta t) := c_{i,j,k}^+(t) + \delta t \mathcal{A}_{i,j,k}(c^+(t), \phi(t)).$$

$$(4.28) \quad c_{i,j,k}^-(t + \delta t) := c_{i,j,k}^-(t) + \delta t \mathcal{A}_{i,j,k}(c^-(t), \phi(t)).$$

This discretization can be also used for solving the macroscopic PNP equations (1.1)–(1.2), provided that δt is chosen sufficiently small. We note that we use different notation to denote the time step, Δt , in our BD model in Table 2, and the time step δt to solve the PNP equations (1.1)–(1.2) in (4.28), because time steps δt and Δt can be, in general, different.

We also need to calculate potential $\phi_{i,j,k}(t + \delta t)$ at time $t + \delta t$ by solving the Poisson equation (1.2). We write it as

$$(4.29) \quad \phi_{i,j,k}(t + \delta t) := \phi_{i,j,k}^m + \widehat{\phi}_p(\mathbf{x}_{i,j,k}),$$

where $\widehat{\phi}_p(\mathbf{x})$ is the solution of (4.13) giving the potential corresponding to the permanent charges. This potential is the same as in our BD simulations in Section 4.2 and is precomputed at the beginning of the simulation. Subtracting equations (1.2) and (4.13), we obtain $\phi_{i,j,k}^m$ in equation (4.29) as $\phi_{i,j,k}^m = \phi^m(\mathbf{x}_{i,j,k})$, where ϕ^m solves the Poisson equation

$$(4.30) \quad \nabla^2 \phi^m = -\frac{1}{\varepsilon_0 \varepsilon} [q^+ c^+ + q^- c^-].$$

To solve equation (4.30), we use the central difference method to discretize the Laplacian to obtain

$$(4.31) \quad \frac{\phi_{i+1,j,k}^m - 2\phi_{i,j,k}^m + \phi_{i-1,j,k}^m}{\Delta x_1^2} + \frac{\phi_{i,j+1,k}^m - 2\phi_{i,j,k}^m + \phi_{i,j-1,k}^m}{\Delta x_2^2} + \frac{\phi_{i,j,k+1}^m - 2\phi_{i,j,k}^m + \phi_{i,j,k-1}^m}{\Delta x_3^2} = -\frac{1}{\varepsilon_0 \varepsilon} \left[q^+ c_{i,j,k}^+ + q^- c_{i,j,k}^- \right]$$

and use the fast Fourier transform algorithm to calculate the discrete Fourier transform $f_{i',j',k'}$ of the right-hand side of equation (4.31), where $c_{i,j,k}^\pm$ are evaluated at time $t + \delta t$. We then calculate $\phi_{i,j,k}^m$, using the inverse fast Fourier transform algorithm, as the inverse discrete Fourier transform of $\lambda(i',j',k')f_{i',j',k'}$, where the factor $\lambda(i',j',k')$ is the discrete Fourier transform of the discretized Laplacian on the left hand side of equation (4.31), given by

$$(4.32) \quad \lambda(i',j',k') = \frac{1}{2} \left[\left(1 - \cos \frac{2\pi(i'-1)}{n_1} \right) \frac{n_1^2}{L_1^2} + \left(1 - \cos \frac{2\pi(j'-1)}{n_2} \right) \frac{n_2^2}{L_2^2} + \left(1 - \cos \frac{2\pi(k'-1)}{n_3} \right) \frac{n_3^2}{L_3^2} \right]^{-1}.$$

The potential $\phi_{i,j,k}(t + \delta t)$ is then calculated by adding the potential corresponding to the permanent charges using equation (4.29).

In Figure 2, we include the solutions of the PNP system (1.1)–(1.2) as the red lines. We use $n_1 = n_2 = n_3 = 96 = 3 \times 2^5$ mesh points in each direction. The small prime factors enable a relatively simple implementation of the Cooley and Tukey algorithm [40] for the discrete Fourier transform. In Figure 2, we present the calculated equilibrium densities as functions of the first coordinate, x_1 , integrating over the x_2 and x_3 coordinates. To highlight the discreteness of the calculated results, we plot this one-dimensional (marginal) distribution of mobile Na^+ ions in Figure 2(a) as the red dots (obtained at $n_1 = 96$ values) connected by the red line. The one-dimensional (marginal) distribution of mobile Cl^- ions is presented in Figure 2(b). The calculated values at mesh points in the vicinity of the layers of permanent charges (at $x_1 = 0$ and $x_1 = L/2$) are well above the BD results. This error cannot be improved by increasing the mesh sizes n_1 , n_2 and n_3 , because it is caused by the error term (3.18), which is missing in the PNP system. This error term includes the Lennard-Jones potential of permanent charges, which would prevent the density c^\pm accumulating in the mesh points close to the permanent charges, if it was included in macroscopic description. While this could decrease the observed difference between the PNP solution and BD simulations around $x_1 = 0$ and $x_1 = L/2$ in Figure 2, it is not straightforward to add the Lennard-Jones potential to the PDE simulations, because it introduces relatively steep potential gradients in a few mesh points next to the permanent charges, leading to numerical instabilities. An alternative approach is to solve the PDEs only in sub-regions not containing permanent charges. This leads to the multi-resolution algorithm developed in Section 5, where detailed BD simulations are used to capture the system dynamics in sub-regions close to the permanent charges.

5. Multi-resolution simulations. Considering reaction-diffusion processes of electroneutral particles, PDE-assisted Brownian Dynamics [25] combines BD simulations based on equation (1.4) with solving macroscopic reaction-diffusion PDEs in parts of the computational domain. In our model system, we would like to apply BD

close to the layers of permanent charges and use PNP equations in the rest of the computational domain. To achieve this, we need to extend the multi-resolutions approach to the models of charged particles, where the macroscopic system is given by the PNP equations (1.1)–(1.2). Following [25], we divide our computational domain (2.1) into two overlapping subdomains

$$(5.1) \quad \Omega_B = \left(\left[0, \omega_B \right] \cup \left[\frac{L_1}{2} - \omega_B, \frac{L_1}{2} + \omega_B \right] \cup \left[L_1 - \omega_B, L_1 \right] \right) \times [0, L_2] \times [0, L_3],$$

$$(5.2) \quad \Omega_P = \left(\left[\omega_P, \frac{L_1}{2} - \omega_P \right] \cup \left[\frac{L_1}{2} + \omega_P, L_1 - \omega_P \right] \right) \times [0, L_2] \times [0, L_3],$$

where ω_B and ω_P satisfy

$$(5.3) \quad 0 < \omega_P \leq \omega_B < \frac{L_1}{4}.$$

The subdomain Ω_B contains the layers of permanent charges, where we have observed the large difference between the results obtained by the PNP system (1.1)–(1.2) and BD simulations. In particular, we will use BD in domain Ω_B to follow trajectories of individual ions, while we use macroscopic concentration profiles c^\pm in domain Ω_P . In $\omega_P = \omega_B$, then the intersection of Ω_P and Ω_B only contains the two-dimensional boundaries of these regions, while we have an overlap region if $\omega_P < \omega_B$. We denote it by

$$(5.4) \quad O = \Omega_P \cap \Omega_B = O_x \times [0, L_2] \times [0, L_3],$$

where

$$O_x = \left[\omega_P, \omega_B \right] \cup \left[\frac{L_1}{2} - \omega_B, \frac{L_1}{2} - \omega_P \right] \cup \left[\frac{L_1}{2} + \omega_P, \frac{L_1}{2} + \omega_B \right] \cup \left[L_1 - \omega_B, L_1 - \omega_P \right].$$

We use grid sizes (4.23) to discretize the PNP equations at mesh points (4.24) and notation (4.25) to denote the values of concentrations c^\pm at the mesh points. The state of the multi-resolution system at time t is given by

- (s1) the values of concentrations $c_{i,j,k}^+(t)$ and $c_{i,j,k}^-(t)$ at mesh points $\mathbf{x}_{i,j,k} \in \Omega_P$,
- (s2) positions $\mathbf{X}_j^\pm(t) = [X_{j,1}^\pm(t), X_{j,2}^\pm(t), X_{j,3}^\pm(t)] \in \Omega_B$, for $j = 1, 2, \dots, N_B^\pm(t)$,
- (s3) potential $\phi_{i,j,k}(t)$ for $i = 1, 2, \dots, n_1$, $j = 1, 2, \dots, n_2$, $k = 1, 2, \dots, n_3$,

where $N_B^\pm(t)$ is the number of positive and negative ions described as individual particles by the BD approach. The number of simulated ions $N_B^\pm(t)$ depends on time t . Since we have, in general, $N_B^+(t) \neq N_B^-(t)$, the BD subsystem does not satisfy electroneutrality on its own, but the system will be electroneutral when both parts of the system (s1) and (s2) are considered together. In particular, we will not use the Ewald summation, but the potential ϕ at mesh points (4.24) will be calculated as the sum of two terms given by equation (4.29), where potential $\phi_{i,j,k}^m$ corresponds to the mobile charges expressed either as concentrations c^\pm of ions in Ω_P or as individual ions in Ω_B .

One iteration of the multi-resolution algorithm is given in Table 3 as Algorithm [M1]–[M8]. Given the state of the multi-resolution system (s1)–(s3) at time t , Algorithm [M1]–[M8] calculates the state of the system (s1)–(s3) at time $t + \Delta t$. In

step [M1], we evolve the concentrations c^\pm from time t to time $t + \Delta t$ by using (4.27)–(4.28). Since the state of the multi-resolution system (s1)–(s3) only defines c^\pm in subdomain Ω_P , we first extend it by zero to the rest of the simulation domain by putting

$$(5.5) \quad c_{i,j,k}^\pm(t) := 0 \quad \text{for mesh points } \mathbf{x}_{i,j,k} \in \Omega \setminus \Omega_P.$$

Then we can apply equations (4.27)–(4.28) to calculate the time evolution of concentrations c^\pm over one time step $[t, t + \Delta t]$. Since the calculated concentrations are further modified in step [M6] to take into account the transfer of ions between Ω_B and Ω_P , we will add an extra asterisk to c^\pm and denote the output of calculations in step [M1] as $c_{i,j,k}^{*,+}(t + \Delta t)$ and $c_{i,j,k}^{*,-}(t + \Delta t)$, rewriting equations (4.27)–(4.28) as

$$(5.6) \quad c_{i,j,k}^{*,+}(t + \delta t) := c_{i,j,k}^+(t) + \delta t \mathcal{A}_{i,j,k}(c^+(t), \phi(t)).$$

$$(5.7) \quad c_{i,j,k}^{*,-}(t + \delta t) := c_{i,j,k}^-(t) + \delta t \mathcal{A}_{i,j,k}(c^-(t), \phi(t)).$$

Equations (5.6)–(5.7) are formulated with time step δt , which we use in Section 4.3 to solve PNP equations. We can choose $\delta t \leq \Delta t$ such that $\Delta t/\delta t$ is an integer. If $\delta t < \Delta t$, then we apply equations (5.6)–(5.7) multiple times to calculate concentrations $c^{*,\pm}$ at time $t + \Delta t$ in step [M1].

In step [M2], we calculate drift terms $\mathbf{a}_j^\pm(t)$, for $j = 1, 2, \dots, N_B^\pm(t)$, which are used to update the positions of BD ions over one time step using equation (4.18). The calculation of drift terms follows steps [A1]–[A4], where N is substituted by $N_B^\pm(t)$ and we formally use $\beta = \infty$, *i.e.* we only consider the Lennard-Jones forces between all permanent and mobile ions in formulas used in [A1]–[A4] and add their contributions to drift terms $\mathbf{a}_j^\pm(t)$ in step [M2]. Then we also add all electric contributions in step [M2] by differentiating electric potential using the central difference scheme as

$$(5.8) \quad \nabla \phi(\mathbf{x}) = \left[\frac{\phi_{i+1,j,k} - \phi_{i-1,j,k}}{2\Delta x_1}, \frac{\phi_{i,j+1,k} - \phi_{i,j-1,k}}{2\Delta x_2}, \frac{\phi_{i,j,k+1} - \phi_{i,j,k-1}}{2\Delta x_3} \right],$$

where

$$(5.9) \quad i = \left\lceil \frac{x_1}{\Delta x_1} \right\rceil, \quad j = \left\lceil \frac{x_2}{\Delta x_2} \right\rceil, \quad k = \left\lceil \frac{x_3}{\Delta x_3} \right\rceil, \quad \text{for } \mathbf{x} = [x_1, x_2, x_3],$$

and $\lceil \cdot \rceil$ denotes the ceiling function, rounding a real number up to the nearest integer. Here, we again use the convention that indices i , j and k are periodic with periods n_1 , n_2 and n_3 , respectively. Step [M3] is the same as step [A6], where we perform the BD part of the simulation by substituting the calculated drift terms $\mathbf{a}_j^\pm(t)$, for $j = 1, 2, \dots, N_B^\pm(t)$, into equation (4.18).

While concentration c^\pm was equal to zero in $\Omega_B \setminus \Omega_P$ at time t by our definition (5.5), we will have a nonzero concentrations $c^{*,\pm}$ at time $t + \Delta t$. The mean number of ions ‘spilling over’ to $\Omega_B \setminus \Omega_P$ during the time interval $[t, t + \Delta t]$ is calculated in step [M4] by

$$(5.10) \quad \mu^\pm(t + \Delta t) = \int_{\Omega_B \setminus \Omega_P} c^{*,\pm}(\mathbf{x}, t + \Delta t) \, d\mathbf{x}.$$

If Δt is sufficiently small, then we have $\mu^\pm(t + \Delta t) \ll 1$. In particular, $\mu^\pm(t + \Delta t)$ can be interpreted as the probability of introducing new ion into $\Omega_B \setminus \Omega_P$. Its new

[M1]	Extend the values of concentrations $c_{i,j,k}^+(t)$ and $c_{i,j,k}^-(t)$ outside of the subdomain Ω_P by (5.5) and calculate the concentrations $c_{i,j,k}^{*,+}(t + \Delta t)$ and $c_{i,j,k}^{*,-}(t + \Delta t)$ by equations (5.6)–(5.7).
[M2]	Calculate drift terms $\mathbf{a}_j^\pm(t)$, for $j = 1, 2, \dots, N_B^\pm(t)$, using steps [A1]–[A4] where N is replaced by $N_B^\pm(t)$ and $\beta = \infty$. Put $\mathbf{a}_j^\pm(t) := \mathbf{a}_j^\pm(t) - \alpha^\pm q^\pm \bar{\nabla} \phi(\mathbf{X}_j^\pm), \quad \text{for } j = 1, 2, \dots, N_B^\pm(t),$ where $\bar{\nabla} \phi(\mathbf{X}_j^\pm)$ is a discretized gradient given by equation (5.8).
[M3]	Generate $6N_B(t)$ coordinates of vectors $\boldsymbol{\xi}_j$, for $j = 1, 2, \dots, N_B^\pm(t)$, as normally distributed numbers with zero mean and unit variance. Calculate the positions of mobile ions at time $t + \Delta t$ by using equation (4.18).
[M4]	Calculate $\mu^\pm(t + \Delta t)$ using (5.10). Put $\gamma^\pm = \gamma_{(i)}^\pm$ where $\gamma_{(i)}^\pm$ is given by (5.12).
[M5]	Generate two uniformly distributed random numbers r^\pm in $(0, 1)$. If $r^\pm < \mu^\pm(t + \Delta t)$, then create new ion in $\Omega_B \setminus \Omega_P$ according to the probability density $p_B^\pm(x, t + \Delta t)$ defined in (5.11) and set $\gamma^\pm = \gamma_{(ii)}^\pm$ where $\gamma_{(ii)}^\pm$ is given by (5.12).
[M6]	Terminate trajectories of BD ions which landed in $\Omega_P \setminus \Omega_B$. Calculate concentrations $c_{i,j,k}^+(t + \Delta t)$ and $c_{i,j,k}^-(t + \Delta t)$ in Ω_P by (5.14).
[M7]	Calculate the discrete Fourier transform $f_{i',j',k'}$ of $-\frac{1}{\varepsilon_0 \varepsilon} \left[q^+ c_{i,j,k}^+ + q^- c_{i,j,k}^- + q^+ \sum_{j=1}^{N_B^+} \frac{\delta(\mathbf{X}_j^+)}{\Delta x_1 \Delta x_2 \Delta x_3} + q^- \sum_{j=1}^{N_B^-} \frac{\delta(\mathbf{X}_j^-)}{\Delta x_1 \Delta x_2 \Delta x_3} \right]$ using the fast Fourier transform algorithm.
[M8]	Calculate $\phi_{i,j,k}^m$ as the inverse discrete Fourier transform of $\lambda(i', j', k') f_{i',j',k'}$, where $\lambda(i', j', k')$ is given by (4.32), using the inverse fast Fourier transform algorithm. Calculate $\phi_{i,j,k}(t + \Delta t)$ using equation (4.29).

TABLE 3
One iteration of the multi-resolution algorithm.

position is sampled in step [M5] according to the probability distribution

$$(5.11) \quad p_B^\pm(\mathbf{x}, t + \Delta t) = \frac{c^{*,\pm}(\mathbf{x}, t + \Delta t)}{\mu^\pm(t + \Delta t)}, \quad \text{for } \mathbf{x} \in \Omega_B \setminus \Omega_P.$$

To ensure the conservation of the total number of particles in the whole domain Ω , we also calculate scaling factors γ^\pm in steps [M4]–[M5], which are then used in step [M6] to rescale the concentration profiles $c^{*,\pm}$ in Ω_P . The scaling factors are given by

$$(5.12) \quad \gamma_{(i)}^\pm = \frac{N - N_B^\pm(t)}{N - N_B^\pm(t) - \mu^\pm(t + \Delta t)}, \quad \gamma_{(ii)}^\pm = \frac{N - N_B^\pm(t) - 1}{N - N_B^\pm(t) - \mu^\pm(t + \Delta t)},$$

where $N_B^+(t)$ (resp. $N_B^-(t)$) is the number of individually simulated positive (resp. negative) ions in the BD domain Ω_B at time t .

In step [M6], we identify the ions which left the subdomain Ω_B . They are removed from the simulation and added as Dirac delta function to obtain new concentration

profiles in Ω_P at time $t + \Delta t$. More precisely, we define the discretized version of the Dirac delta function centered at position \mathbf{x} by

$$(5.13) \quad \delta(\mathbf{x})_{i,j,k} = \begin{cases} 1 & \text{for } i, j \text{ and } k \text{ given by (5.9);} \\ 0 & \text{otherwise.} \end{cases}$$

Then the concentration update in step [M6] can be written as

$$(5.14) \quad c_{i,j,k}^{\pm}(t + \Delta t) = \gamma^{\pm} c_{i,j,k}^{*,\pm}(t + \Delta t) + \sum_{\ell \in \mathcal{J}^{\pm}} \frac{\delta(\mathbf{X}_{\ell}^{\pm}(t + \Delta t))_{i,j,k}}{\Delta x_1 \Delta x_2 \Delta x_3}$$

where the set \mathcal{J}^+ (resp. \mathcal{J}^-) contains indices of all terminated positive (resp. negative) ions. In particular, we transform the position of each terminated positive (resp. negative) ion into its nearest mesh point (i, j, k) by applying equation (5.9) and add $1/(\Delta x_1 \Delta x_2 \Delta x_3)$ to the corresponding concentration c^+ (resp. c^-) at this mesh point. The concentration update (5.14) preserves the number of particles, which we can formulate using our state variables (s1)–(s3) as

$$(5.15) \quad N = N_B^{\pm}(t) + \int_{\Omega_P} c^{\pm}(\mathbf{x}, t) d\mathbf{x} = N_B^{\pm}(t) + \sum_{i=1}^{n_1} \sum_{j=1}^{n_2} \sum_{k=1}^{n_3} c_{i,j,k}^{\pm}(t) \Delta x_1 \Delta x_2 \Delta x_3,$$

where we have used the extension (5.5) on the right-hand side, *i.e.* the summation on the right-hand side can be restricted to the mesh points, which are in Ω_P .

Finally, steps [M7]–[M8] calculate the potential $\phi_{i,j,k}(t + \Delta t)$. In step [M7], the right-hand side of the discretized Poisson equation (4.31) is modified to take into account all mobile ions at time $t + \Delta t$. In particular, potential $\phi_{i,j,k}^m$ (calculated in step [M8] by the inverse fast Fourier transform algorithm) will contain electric potential corresponding to all mobile ions, which are either represented as individual ions or as concentration fields. In step [M8], we add potential $\phi_p(\mathbf{x})$ corresponding to the permanent charges using equation (4.29) at time $t + \Delta t$. As it has been the case of our BD simulations in Section 4.2 or our PNP solutions, the permanent potential can be precomputed at the beginning of the simulation by solving equation (4.13).

In Figure 3, we present the equilibrium distributions of mobile Na^+ and Cl^+ ions calculated using Algorithm [M1]–[M8] for three sets of simulation parameters, systematically scaling the x_1 direction. All simulations are performed in three-dimensional domains, Ω given by (2.1), where we use $L_2 = L_3 = 10$ nm and $n_2 = n_3 = 96 = 3 \times 2^5$ mesh points in the x_2 and x_3 directions to discretize the Poisson equation, *i.e.* our spatial resolution for the calculation of electric potential is the same as we used in Section 4.3. The permanent charges are positioned at locations (4.1)–(4.2) with $N_p = 6$. The mesh sizes are given by (4.23) and we use three parameter sets constructed to preserve the average ion density $N/(L_1 L_2 L_3)$, the mesh size Δx_1 and the relative size of the overlap region. Specifically, in panels (a)–(b) we set $N = 100$, $L_1 = 10$ nm, $n_1 = 96$, $\omega_P = 5\Delta x_1$ and $\omega_B = 10\Delta x_1$; in (c)–(d) the system is doubled in length ($L_1 = 20$ nm) and particle number ($N = 200$), with $n_1 = 192$, $\omega_P = 10\Delta x_1$ and $\omega_B = 20\Delta x_1$; and in (e)–(f) the length and particle number are again doubled ($L_1 = 40$ nm, $N = 400$), with $n_1 = 384$, $\omega_P = 20\Delta x_1$ and $\omega_B = 40\Delta x_1$. This scaling strategy studies the effect of system size on the performance of Algorithm [M1]–[M8]. Since the multi-resolution simulation explicitly includes the Lennard-Jones potentials around permanent charges, the density c^{\pm} does not accumulate in the mesh points close to the permanent charges, which happens for the PNP solution in Figure 2.




FIG. 3. *The equilibrium distributions of mobile Na^+ ions (panels (a), (c) and (e)) and mobile Cl^- ions (panels (b), (d) and (f)) calculated using Algorithms [M1]–[M8] (histograms) and [A1]–[A6] (black dots). We present the results calculated for three sets of parameters:*

- (a)–(b) $N = 100$, $L_1 = 10$ nm, $n_1 = 96$, $\omega_P = 5 \Delta x_1$, and $\omega_B = 10 \Delta x_1$,
- (c)–(d) $N = 200$, $L_1 = 20$ nm, $n_1 = 192$, $\omega_P = 10 \Delta x_1$, and $\omega_B = 20 \Delta x_1$,
- (e)–(f) $N = 400$, $L_1 = 40$ nm, $n_1 = 384$, $\omega_P = 20 \Delta x_1$, and $\omega_B = 40 \Delta x_1$,

where $\Delta x_1 = L_1/n_1$ has the same value in all three simulations. Other parameters are also the same and equal to $N_p = 6$, $n_2 = n_3 = 96$, $L_2 = L_3 = 10$ nm, $\kappa = 3$ and $\beta = 0.05$.

In Figure 3, we present the calculated equilibrium densities as functions of the first coordinate, x_1 , integrating over the x_2 and x_3 coordinates. To highlight the multi-resolution nature of the simulation, the concentrations c^\pm in the region $\Omega_P \setminus \Omega_B$ are denoted using light blue (cyan) bars, the histograms in the region $\Omega_B \setminus \Omega_P$ are visualized using gray bars and the red bars show the average concentration in the overlap region (5.4), where some mobile ions are treated as concentration profiles and some mobile ions are described as individual particles (we add their contributions together to obtain the red bars). The simulations in panels (c)–(d) and (e)–(f) of Figure 3 use $n_1 = 2 \times 96 = 192$ and $n_1 = 4 \times 96 = 384$ mesh points in the x_1 direction, respectively, giving the same mesh size Δx_1 in each simulation. To enable comparison with the results shown in panels (a)–(b), we present the equilibrium densities over 96 mesh points in the x_1 direction for each simulation, adding the values over two or four neighbouring mesh points in panels (c)–(d) and (e)–(f), respectively. In Figure 3, we




FIG. 4. *The equilibrium distributions of mobile Na^+ ions (panel (a)) and mobile Cl^- ions (panel (b)) calculated using Algorithms [M1]–[M8] (histograms) and [A1]–[A6] (black dots) for parameters: $N = 100$, $L_1 = L_2 = L_3 = 10$ nm, $n_1 = n_2 = n_3 = 96$, $\omega_P = 5 L_1/n_1$, $\omega_B = 15 L_1/n_1$, $N_p = 6$, $\kappa = 3$ and $\beta = 0.05$.*

compare the results obtained by Algorithm [M1]–[M8] with the results obtained by Algorithm [A1]–[A6]. We also quantify the accuracy of the results by calculating the mean squared error

$$(5.16) \quad \text{Er}^\pm = \sum_{i=1}^{96} \left(c_{i,\text{MR}}^\pm - c_{i,\text{BD}}^\pm \right)^2$$

where $c_{i,\text{MR}}^\pm$, $i = 1, 2, \dots, 96$, are the values plotted in Figure 3 as histograms, showing the average concentrations of mobile ions in the region $[(i-1)L_1/96, iL_1/96] \times [0, L_2] \times [0, L_3]$ estimated by Algorithm [M1]–[M8], and $c_{i,\text{BD}}^\pm$ are average concentrations of mobile ions in the same regions estimated by Algorithm [A1]–[A6], shown as black dots in Figure 3. The mean square error Er^+ calculated by (5.16) is equal to $4.24 \times 10^{-5} \text{ nm}^{-6}$ for the simulation in Figure 3(a) with $N = 100$ of Na^+ mobile ions, while we have $\text{Er}^+ = 2.10 \times 10^{-5} \text{ nm}^{-6}$ for $N = 200$ in Figure 3(c), and $\text{Er}^+ = 1.12 \times 10^{-5} \text{ nm}^{-6}$ for $N = 400$ in Figure 3(e), illustrating that the error Er^+ decreases by a factor of two each time with double the domain size. The error Er^- also decreases as we increase the domain size. We have $\text{Er}^- = 4.64 \times 10^{-5} \text{ nm}^{-6}$ in Figure 3(b), $\text{Er}^- = 1.42 \times 10^{-5} \text{ nm}^{-6}$ in Figure 3(d), and $\text{Er}^- = 8.52 \times 10^{-6} \text{ nm}^{-6}$ in Figure 3(f).

The mean square errors defined by (5.16) also decrease as we increase the size of the overlap region. This is illustrated in Figure 4, where we consider the same domain size and boundary ω_P as in panels (a)–(b) of Figure 3, i.e. we have domain Ω given by (2.1) with $L_1 = L_2 = L_3 = 10$ nm and $\omega_P = 5 L_1/n_1 = 5\Delta x_1$. In Figure 4, we double the length of the overlap region by increasing its second boundary to $\omega_B = 15 L_1/n_1 = 15\Delta x_1$. The mean square errors defined by (5.16) are $\text{Er}^+ = 7.45 \times 10^{-6} \text{ nm}^{-6}$ in Figure 4(a) and $\text{Er}^- = 1.39 \times 10^{-5} \text{ nm}^{-6}$ in Figure 4(b). Comparing with panels (a)–(b) of Figure 3, we observe that errors Er^+ and Er^- decrease by factors 5.69 and 3.34, respectively.

6. Discussion. In this paper, we have extended the PDE-assisted BD approach for reaction-diffusion systems, developed in [25], to models of charged particles. One of the key changes for such systems is that the reaction-diffusion macroscopic PDE is, in general, replaced by a reaction-diffusion-advection PDE describing the time evolution of concentrations of biochemical species involved. This generalization of the method in [25] would be relatively straightforward if the BD particles passively responded to an underlying background potential. However, in applications to modelling ions and ion channels, BD particles carry charges which generate electrostatic

potential. In particular, the macroscopic equation can be written in the form of the PNP system (1.1)–(1.2), which we formally derived from the BD model (2.3) in Section 3. While the PNP system (1.1)–(1.2) is justified in a bulk solution [28], the error term (3.18) can have significant effects on the system dynamics close to the biological structures of interest. For example, considering applications to modelling ion channels, the PNP system (1.1)–(1.2) can be used in regions far away from an ion channel, but its applicability to the passage of ions through the ion channel is questionable, especially in its narrowest part as indicated by computational studies in the literature [16, 18, 19]. The behaviour of particles in such specific geometries is often not captured by the mean field limit, which employs the propagation of chaos assumption and appropriate scaling of interaction terms to derive macroscopic PDEs in the limit as $N \rightarrow \infty$ [41, 42].

In Section 4, we have presented an illustrative system with layers of permanent charges showing that the Lennard-Jones potential creates exclusion zones around permanent charges which are not captured by the PNP system (1.1)–(1.2). Such observations motivate the development of a multi-resolution approach in Section 5, which uses PDEs (1.1)–(1.2) in the bulk and more detailed BD simulations in the regions close to the permanent charges. While the short-range hard sphere correlations could be, in principle, included by modifying the PNP model [43], one of the advantages of hybrid (multi-resolution) simulation frameworks is that they enable to further increase the modeling accuracy in BD regions by incorporating additional modelling details, for example, by using a molecular dynamics description in small subsets of BD regions [44].

To design the multi-resolution scheme in Section 5, we have extended the PDE-assisted BD approach for reaction-diffusion systems, developed in [25], to models of charged particles. The resulting multi-resolution Algorithm [M1]–[M8] uses the discrete Fourier transform to solve the Poisson equation. In particular, we need to transform positions of individual ions to the corresponding mesh points by (5.13) and use the calculated mesh-based potential to extract the forces by (5.8). Equations (5.8) and (5.13) possess the necessary symmetry to ensure that each individual ion does not respond to the potential arising from its own charge. However, this remains a different approach compared to the BD method that employs the Ewald summation. In the case of reaction-diffusion systems [25], the multi-resolution algorithm used a BD approach without modifications, because there were no long-range forces to consider. If a modeller wants to use the Ewald summation for interactions between individually simulated particles in Ω_B , then they would need to ensure the electroneutrality in the BD subdomain, required by the Ewald summation. To further improve the accuracy, there is a potential to use some hybrid approaches like particle-particle-particle mesh algorithms [45, 46], or to use multi-grid approaches [47].

In our implementation of Algorithm [M1]–[M8] in Section 5, the multi-resolution method includes the overlap region (5.4), where both the PDE description and BD exist in parallel. In some multi-resolution approaches, it is possible to couple models with different resolutions using a simple interface, as it has been shown for coupling BD simulations with a coarser description given by a compartment-based model (reaction-diffusion master equation) in [21]. The advantage of a simple interface over an overlap region is that this can simplify some aspects of the software implementation of the resulting multi-resolution method [10]. It remains an open question for future investigation whether the BD modelling of charged particles can be coarse-grained as a compartment-based stochastic model which could potentially avoid using an overlap region in a multi-resolution scheme. However, if a BD model is coarse-

grained using mean-field PDEs (like the PNP system in this paper), then the use of an overlap region improves the accuracy, as it was previously shown for reaction-diffusion systems in the literature [25, 48]. The benefits of overlap (bridging, blending, transition) regions have been also demonstrated in other multi-resolution approaches, including coupling molecular dynamics and BD simulations [12, 8, 49], compartment-based models with macroscopic PDEs [50, 51, 52], to bridge particle-based BD models with macroscopic PDEs [53, 54], to coarse-grain the chemical master equation in the state space [60], and more broadly to connect atomistic and continuum modelling in material science [55, 56, 57, 58, 59, 61].

Considering our illustrative example in Section 5, the domain decomposition into the BD and PDE subdomains Ω_B and Ω_P remained fixed throughout the entire simulation. In some application areas, it is necessary to consider extending multi-resolution techniques to scenarios where the BD subdomain Ω_B changes over time [24], or the domain undergoes temporal growth [62, 63].

Data Availability. In compliance with EPSRC’s open access initiative, the data in this paper is available from: <https://doi.org/10.5281/zenodo.17979223>.

REFERENCES

- [1] R. Erban and S.J. Chapman, *Stochastic Modelling of Reaction-Diffusion Processes*, Cambridge Texts in Applied Mathematics, Cambridge University Press, 308 pages, 2020
- [2] M. Bazant, K. Thornton and A. Ajdari, “Diffuse-charge dynamics in electrochemical systems”, *Physical Review E—Statistical, Nonlinear, and Soft Matter Physics* **70**(2), 021506, 2004
- [3] V. Klika and E.A. Gaffney, “Upscaling the Poisson–Nernst–Planck equations for ion transport in weakly heterogeneous charged porous media”, *Applied Mathematics Letters* **137**, p. 108482, 2023
- [4] B. Eisenberg, “Ionic channels in biological membranes—electrostatic analysis of a natural nanotube”, *Contemporary Physics* **39**(6), pp. 447–466, 1998
- [5] M.G. Kurnikova, R.D. Coalson, P. Graf, and A. Nitzan, “A lattice relaxation algorithm for three-dimensional Poisson-Nernst-Planck theory with application to ion transport through the gramicidin A channel”, *Biophysical Journal* **76**(2), pp. 642–656, 1999
- [6] A.E. Cardenas, R.D. Coalson, and M.G. Kurnikova, “Three-dimensional Poisson-Nernst-Planck theory studies: Influence of membrane electrostatics on gramicidin A channel conductance”, *Biophysical Journal* **79**(1), pp. 80–93, 2000
- [7] U. Hollerbach, D.P. Chen, and R.S. Eisenberg, “Two-and three-dimensional Poisson–Nernst–Planck simulations of current flow through gramicidin A”, *Journal of Scientific Computing* **16**, pp. 373–409, 2001
- [8] R. Erban, “Coupling all-atom molecular dynamics simulations of ions in water with Brownian dynamics”, *Proceedings of the Royal Society A* **472**, 20150556, 2016
- [9] S. Andrews, N. Addy, R. Brent and A. Arkin, “Detailed simulations of cell biology with Smoldyn 2.1”, *PLoS Computational Biology* **6**(3), e1000705, 2010
- [10] M. Robinson, S. Andrews and R. Erban, “Multiscale reaction-diffusion simulations with Smoldyn”, *Bioinformatics* **31**(14), pp. 2406–2408, 2015
- [11] K. Takahashi, S. Tănase-Nicola and P. Ten Wolde, “Spatio-temporal correlations can drastically change the response of a MAPK pathway”, *Proceedings of the National Academy of Sciences* **107**(6), pp. 2473–2478, 2010
- [12] R. Erban, “From molecular dynamics to Brownian dynamics”, *Proceedings of the Royal Society A* **470**, 20140036, 2014
- [13] R. Erban, “Coarse-graining molecular dynamics: stochastic models with non-Gaussian force distributions”, *Journal of Mathematical Biology* **80**, pp. 457–479, 2020
- [14] M. Flegg, S. Rüdiger and R. Erban, “Diffusive spatio-temporal noise in a first-passage time model for intracellular calcium release”, *Journal of Chemical Physics* **138**(15), 154103, 2013
- [15] U. Dobramysl, S. Rüdiger and R. Erban, “Particle-based multiscale modeling of calcium

- puff dynamics”, *Multiscale Modelling and Simulation* **14**(3), pp. 997-1016, 2016
- [16] B. Corry, S. Kuyucak and S. Chung, “Tests of continuum theories as models of ion Channels. II. Poisson-Nernst-Planck theory versus Brownian dynamics”, *Biophysical Journal* **78**, pp. 2364-2381, 2000
- [17] W. Chen, R. Erban and S.J. Chapman, “From Brownian dynamics to Markov chain: an ion channel example”, *SIAM Journal on Applied Mathematics* **74**(1), pp. 208-235, 2014
- [18] C. Song and B. Corry, “Testing the applicability of Nernst-Planck theory in ion channels: comparisons with Brownian dynamics simulations”, *PLoS One* **6**(6), e21204, 2011
- [19] W. Im and B. Roux, “Ion permeation and selectivity of OmpF porin: a theoretical study based on molecular dynamics, Brownian dynamics, and continuum electrodiffusion theory”, *Journal of Molecular Biology* **322**, pp. 851-869, 2002
- [20] L. Li, J. Liu and G. Tang, “Some random batch particle methods for the Poisson-Nernst-Planck and Poisson-Boltzmann Equations”, *Communications in Computational Physics* **32**(1), pp. 41-82, 2022
- [21] M. Flegg, S.J. Chapman and R. Erban, “Two Regime Method for optimizing stochastic reaction-diffusion simulations”, *Journal of the Royal Society Interface* **9**(70), pp. 859-868, 2012
- [22] M. Flegg, S. Hellander and R. Erban, “Convergence of methods for coupling of microscopic and mesoscopic reaction-diffusion simulations”, *Journal of Computational Physics* **289**, pp. 1-17, 2015
- [23] P. Bauler, G. Huber and J.A. McCammon, “Hybrid finite element and Brownian dynamics method for diffusion-controlled reactions” *Journal of Chemical Physics* **136**(16), 164107, 2012
- [24] M. Robinson, M. Flegg and R. Erban, “Adaptive two-regime method: application to front propagation”, *The Journal of Chemical Physics* **140**(12), 2014
- [25] B. Franz, M. Flegg, S.J. Chapman and R. Erban, “Multiscale reaction-diffusion algorithms: PDE-assisted Brownian dynamics”, *SIAM Journal on Applied Mathematics* **73**(3), pp. 1224-1247, 2013
- [26] G. Huber, Y. Miao and S. Zhou, Shenggao, B. Li, Bo and J.A. McCammon, “Hybrid finite element and Brownian dynamics method for charged particles”, *Journal of Chemical Physics* **144**(16), 2016
- [27] Q. Sun, I. Boyd and G. Candler, “A hybrid continuum/particle approach for modeling subsonic, rarefied gas flows”, *Journal of Computational Physics* **194**, pp. 256-277, 2004
- [28] Z. Schuss, B. Nadler and R. Eisenberg, “Derivation of Poisson and Nernst-Planck equations in a bath and channel from a molecular model”, *Physical Review E* **64**(3), 036116, 2001
- [29] R. Erban and R. Van Gorder, “Langevin dynamics for a heavy particle immersed within a flow of light particles”, *Proceedings of the Royal Society A* **480**(2301), 20230851, 2024
- [30] H. Hwang, S. Austin, et al, “CHARMM at 45: Enhancements in accessibility, functionality, and speed”, *Journal of Physical Chemistry B* **128**(41), pp. 9976-10042, 2024
- [31] S. Koneshan, R. Lynden-Bell and J. Rasalah, “Friction coefficients of ions in aqueous solution at 25 °C”, *Jour. of the American Chemical Society* **120**, pp. 12041-12050, 1998
- [32] W. Haynes, et al. “CRC Handbook of Chemistry and Physics: A Ready-Reference Book of Chemical and Physical Data”, 95th edition, CRC Press, 2014
- [33] D. Fernández, et al. “Formulation for the static permittivity of water and steam at temperatures from 238 K to 873 K at pressures up to 1200 MPa, including derivatives and Debye-Hückel coefficients”, *Journal of Physical Chemistry* **26**(4), 1997
- [34] D. Gillespie, W. Nonner and R. Eisenberg, “Coupling Poisson-Nernst-Planck and density functional theory to calculate ion flux”, *Journal of Physics: Condensed Matter* **14**(46), 12129, 2002
- [35] B. Lu, M. Holst, J. McCammon and Y. Zhou, “Poisson-Nernst-Planck equations for simulating biomolecular diffusion-reaction processes I: Finite element solutions”, *Journal of Computational Physics* **229**(19), pp. 6979-6994, 2010
- [36] N. van Kampen, “Stochastic Processes in Physics and Chemistry”, 3rd edition, North-Holland, Amsterdam, 2007
- [37] X. Mao, “Stochastic Differential Equations and Applications”, 2nd edition, Horwood Publishing Limited, 2007
- [38] P. Ewald, “Die Berechnung optischer und elektrostatischer Gitterpotentiale”, *Annalen der Physik* **64**, pp. 253-287, 1921
- [39] D. Frenkel and B. Smit, “Understanding Molecular Simulation: From Algorithms to

- Applications”, Computational Science Series, Volume 1, Academic Press, Elsevier, 2002
- [40] J. Cooley and J. Tukey, “An algorithm for the machine calculation of complex Fourier series”, *Mathematics of computation* **19**(90), pp. 297–301, 1965
- [41] D. Lacker, “Hierarchies, entropy, and quantitative propagation of chaos for mean field diffusions”, *Probability and Mathematical Physics* **4**(2), pp. 377–432, 2023
- [42] R. Erban and J. Haškovec, “Impact of memory on clustering in spontaneous particle aggregation”, submitted, available as Arxiv.Org/abs/2510.15335, 2025
- [43] M. Ma, Z. Xu and L. Zhang, “Modified Poisson–Nernst–Planck Model with Coulomb and Hard-sphere Correlations”, *SIAM Journal on Applied Mathematics* **81**(4), pp. 1645–1667, 2021
- [44] R. Erban and Y. Togashi, “Asymmetric periodic boundary conditions for molecular dynamics and coarse-grained simulations of nucleic acids”, *Journal of Physical Chemistry B* **127**(38), 8257–8267, 2023
- [45] R. Hockney and J. Eastwood, *Computer simulation using particles*, IOP Publishing Ltd, 1988
- [46] P. Gibbon and G. Sutmann, “Long-range interactions in many-particle simulation”, *Quantum simulations of many-body systems: from theory to algorithm*. Eds. J. Groten-dorst, D. Marx and A. Muramatsu. NIC-series 10, pp. 467–506, 2002
- [47] R. Erban and S. Winkelmann, “Multi-grid reaction-diffusion master equation: applications to morphogen gradient modelling”, *Bulletin of Mathematical Biology* **87**(1), 2025
- [48] C. Smith and C. Yates, “The auxiliary region method: a hybrid method for coupling PDE- and Brownian-based dynamics for reaction–diffusion systems” *Royal Society Open Science* **5**(8), 180920, 2018
- [49] M. del Razo, M. Dibak, C. Schütte, Christof and F. Noé, “Multiscale molecular kinetics by coupling Markov state models and reaction-diffusion dynamics”, *Journal of Chemical Physics* **155**(12), 2021
- [50] H. Kang and R. Erban, “Multiscale stochastic reaction-diffusion algorithms combining Markov chain models with stochastic partial differential equations”, *Bulletin of Mathematical Biology* **81**(8), pp. 3185–3213, 2019
- [51] C. Yates and M. Flegg, “The pseudo-compartment method for coupling partial differential equation and compartment-based models of diffusion”, *Journal of The Royal Society Interface* **12**,(106), 20150141, 2015
- [52] C. Smith and C. Yates, “Spatially extended hybrid methods: a review”, *Journal of the Royal Society Interface* **15**(139) 20170931, 2018
- [53] C. Yates, A. George, A. Jordana, C. Smith, A. Duncan and K. Zygalakis, “The blending region hybrid framework for the simulation of stochastic reaction–diffusion processes”, *Journal of the Royal Society Interface* **17**(171), 20200563, 2020
- [54] M. del Razo and M. Kostré, “Open reaction-diffusion systems: bridging probabilistic theory and simulations across scales”, *Journal of Physics A: Mathematical and Theoretical* **58**(14), 145001, 2025
- [55] R. Miller and E. Tadmor, “A unified framework and performance benchmark of fourteen multiscale atomistic/continuum coupling methods”, *Modelling and Simulation in Materials Science and Engineering* **17**, 053001, 2009
- [56] M. Luskin and C. Ortner, “Atomistic-to-continuum coupling.” *Acta Numerica* **22**, pp. 397–508, 2013
- [57] B. Van Koten and M. Luskin, “Analysis of energy-based blended quasi-continuum approximations”, *SIAM Journal on Numerical Analysis* **49**(5), pp. 2182–2209, 2011
- [58] M. Parks, P. Bochev and R. Lehoucq, “Connecting atomistic-to-continuum coupling and domain decomposition”, *Multiscale Modelling and Simulation* **7**(1), pp. 362–380, 2008
- [59] S. Xiao and T. Belytschko, “A bridging domain method for coupling continua with molecular dynamics”, *Computer Methods in Applied Mechanics and Engineering* **193**, pp. 1645–1669, 2004
- [60] A. Duncan, R. Erban and K. Zygalakis, “Hybrid framework for the simulation of stochastic chemical kinetics”, *Journal of Computational Physics* **326**, pp. 398–419, 2016
- [61] E. Smith and P. Theodorakis, “Multiscale simulation of fluids: coupling molecular and continuum”, *Physical Chemistry Chemical Physics* **26**(2), 724–744, 2024
- [62] R. Baker, C. Yates and R. Erban, “From microscopic to macroscopic descriptions of cell migration on growing domains, *Bulletin of mathematical biology* **72**, pp. 719–762, 2010
- [63] C. Smith and C. Yates, “Incorporating domain growth into hybrid methods for reaction–diffusion systems”, *Journal of the Royal Society Interface* **18**(177), 20201047, 2021