

## The Physical Properties of Lyman Break Galaxies at $z > 5$ : Outflows and the “Pre-enrichment Problem”

M. D. Lehnert,<sup>1</sup> M. Bremer,<sup>2</sup> A. Verma,<sup>3</sup> L. Douglas,<sup>2</sup> and N. Förster Schreiber<sup>4</sup>

<sup>1</sup>*GEPI, Observatoire de Paris, France*

<sup>2</sup>*Physics Department, University of Bristol, UK*

<sup>3</sup>*Physics Department, University of Oxford, UK*

<sup>4</sup>*MPE, Germany*

**Abstract.** We discuss the properties of Lyman break galaxies at  $z > 5$  as determined from disparate fields covering approximately  $500 \text{ arcmin}^2$ . While the broad characteristics of the Lyman break galaxy population, such as luminosity functions and clustering amplitude, has been discussed extensively in the literature, we focus on the detailed physical properties of the sources in this large survey ( $\gtrsim 100$  with spectroscopic redshifts). Specifically, we discuss ensemble mass estimates, stellar mass surface densities, core phase space densities, star formation intensities, characteristics of their stellar populations, etc., as obtained from multi-wavelength data (rest-frame UV through optical) for a subsample of these galaxies. In particular, we focus on evidence that these galaxies drive vigorous outflows and speculate that this population may solve the so-called “pre-enrichment problem”. The general picture that emerges from these studies is that these galaxies, observed about 1 Gyr after the Big Bang, have properties consistent with being the progenitors of the densest stellar systems in the local Universe—the centers of old bulges and early-type galaxies.

### 1. High Redshift Galaxies and Cosmology

Galaxies at the highest redshifts,  $z > 5$ , are vital objects in observational cosmology. They formed during an era of one of the most dramatic phase changes in the history of the Universe—the epoch of re-ionization. Remarkably, these earliest galaxies were able to dictate the phase of the general medium in which all galaxies are imbedded. Because of their enhanced  $[\alpha/\text{Fe}]$  ratios, which are the product of core collapse supernova enrichment, and old ages, the densest (i.e., globular clusters) and most massive stellar systems (i.e., giant ellipticals) are thought to have formed at such high redshifts in spectacular bursts of star formation of relatively short duration ( $< 100 \text{ Myr}$  to  $1 \text{ Gyr}$ ). The short duration of the star formation but the large accumulated mass must mean that these galaxies had very high star formation intensities—the rate of star formation per unit surface area. It is well known that galaxies with high star formation intensities drive vigorous outflows of metals and energy. Such outflows from high redshift galaxies could have both cleared material from the galaxian surroundings allowing ionizing photons to escape and enrich the intergalactic medium (IGM). Through this mechanism, problems of how the IGM was re-ionized and enriched in metals even at the highest redshifts yet observed (this is the so-called

“pre-enrichment problem”) could be solved. But can we find direct evidence for this scenario in the physical properties of high redshift galaxies? This is what we will attempt to address here.




Figure 1. We compare the star formation rate intensities—the rate of star formation per unit projected area—of the  $z \approx 5$  LBGs (*diamonds*; Verma et al. 2007),  $z \approx 3$  LBGs (*squares*; Papovich, Dickinson, & Ferguson 2001), and a local sample of UV-selected galaxies (*squares*; Heckman et al. 2005). The horizontal line indicates the threshold above which local starburst exhibit large scale outflows (Heckman 2001; Lehnert & Heckman 1996). All of the galaxies studied here lie above this threshold and thus by analogy, must be driving outflows. We note that the evidence is overwhelming that virtually all  $z \approx 3$  LBGs are driving outflows (Adelberger et al. 2003).

## 2. Our Comprehensive Study of $z > 5$ Galaxies

At  $z \sim 5$ , there have been a number of significant spectroscopic surveys investigating the nature of Lyman break galaxies (LBGs, or “V- or R-band dropouts”), two of which have been led by some of the authors. There are now over 100 confirmed redshifts in total at  $z \gtrsim 5$  through the work of our ESO Large Programme (the ESO Distant Galaxy Survey—ERGS; L. Douglas et al., in preparation) and study of the field we call “BDFs” (e.g., Lehnert & Bremer 2003; M. D. Lehnert et al., in preparation). The ESO LP is based on the EDIScS fields, which are ten widely separated intermediate-redshift cluster fields, while the “BDFs” are four

adjacent fields observable from Paranal in winter. Our total sample constitutes the largest sample of  $z > 5$  LBGs (“ $R$ -band drop outs”) with spectroscopically confirmed redshifts. In all of these fields, we have deep  $R$ -,  $I$ -,  $z$ -band, and IRAC data (3.6, 4.5, 5.8, and  $8.0 \mu\text{m}$ ). Some of the fields also have observations in other optical/infrared bands, including MIPS  $24 \mu\text{m}$  data and *Hubble Space Telescope* ACS imaging. In addition, we have been studying a similar population of galaxies in the GOODS-S, taking advantage of the deep multi-wavelength data set (Verma et al. 2007).

### 3. What Have we Concluded?

From our analysis of some of these extensive data sets (Verma et al. 2007), we conclude:

- The LBGs at  $z \approx 5$  are young, with typical ages less than 100 Myr, compact, with typical half-light radii of about 1 kpc, and are rapidly forming stars, 10 to  $200 M_{\odot} \text{yr}^{-1}$ . These ages are very young suggesting that they have been intensely forming stars for only a few dynamical times (estimated from stellar mass estimates and sizes). They therefore may likely represent a population of “primordial galaxies”.
- These sources are generally so young that they likely did not substantially contribute to re-ionization. The estimates of their ages from the spectral energy distribution (SED) suggests that they formed at redshifts of about 6–7. Their contribution to re-ionization would likely be after the Universe was already substantially ionized.
- From SED fitting from the rest-frame UV through optical, the LBGs at  $z \approx 5$  are typically about a factor of 10 less massive than similarly selected galaxies at  $z \approx 3$  (a few  $10^9 M_{\odot}$  compared to a few  $10^{10} M_{\odot}$ ; Shapley et al. 2001). These galaxies represent about 1% of the local mass density (Verma et al. 2007).
- The LBGs at  $z \approx 5$  have high stellar mass surface densities,  $\mu_{\text{stars}} \approx 2\text{--}6 \times 10^8 M_{\odot} \text{kpc}^{-2}$ , and have very roughly approximated core phase densities of  $10^{-6} \text{pc}^3 \text{km}^3 \text{s}^{-3}$ . Both of these are similar to that of bulges and spheroids of  $M^*$  galaxies at low redshift.
- The LBGs at  $z \approx 5$  have high star formation intensities, well above those needed to drive winds at low redshift,  $\Sigma_{\text{SFR}} \gg 0.1 M_{\odot} \text{kpc}^{-2} \text{yr}^{-1}$  (Fig. 1; Heckman 2001; Lehnert & Heckman 1996).
- If the  $z \approx 5$  LBGs drive strong outflows then they may be able to solve the “pre-enrichment problem”—whereby the metallicity of the IGM does not appear to evolve strongly with redshift at least out to almost  $z = 6$  (Songaila 2001; Ryan-Weber et al. 2006). Quantitatively (crudely) estimating the likely metal contribution of these galaxies suggests that they are able to solve the pre-enrichment problem (see Fig. 2).

**Acknowledgments.** MDL wishes to thank Johan Knapen for organizing such a delightful meeting in honor of the contributions of John Beckman and for his extreme patience in waiting for this contribution.




Figure 2. The co-moving density of C IV absorbers does not appear to evolve strongly with redshift (Songaila 2001). This suggests that a large fraction of the metals in the IGM may have been in place within 1 Gyr after the Big Bang. Comparing the data of the contribution to the closure density due to C IV absorbers,  $\Omega_{\text{CIV}}$  (Songaila 2001; *squares*, and Pettini et al. 2003; Ryan-Weber et al. 2006; the highest redshift diamond, from Ryan-Weber et al., is a lower limit), we find that it is plausible that the  $z \approx 5$  LBGs could eject sufficient metals to explain the IGM absorption line observations. Our range of estimates was derived assuming that  $\dot{M}_{\text{outflow}} = \dot{M}_{\text{star formation}}$  (Lehnert & Heckman 1996), the closure density, that the ionization fraction of  $\text{C}^{3+} = 0.5$ , the metallicity of the outflow is  $0.2 Z_{\odot}$ ,  $\Omega_{\text{b}} h^2 = 0.023$ , and the range of star formation rates derived in Verma et al. (2007).

## References

- Adelberger, K. L., Steidel, C. C., Shapley, A. E., & Pettini, M. 2003, *ApJ*, 584, 45  
 Heckman, T. M., et al. 2005, *ApJ*, 619, 35  
 Heckman, T. M., 2001, in *ASP Conf. Ser.*, 240, Gas and Galaxy Evolution, ed. J. E. Hibbard, M. Rupen, & J. H. Gorkom (San Francisco: ASP), p. 345  
 Lehnert, M. D., & Bremer, M. 2003, *ApJ*, 593, 630  
 Lehnert, M. D., & Heckman, M. 1996, *ApJ*, 472, 546  
 Papovich, C., Dickinson, M., & Ferguson, H. C. 2001, *ApJ*, 559, 620  
 Pettini, M., Madau, P., Bolte, M., Prochaska, J. X., Ellison, S. L., & Fan, X. 2003, *ApJ*, 594, 695  
 Ryan-Weber, E. V., Pettini, M., & Madau, P. 2006, *MNRAS*, 371, L78  
 Shapley, A. E., Steidel, C. C., Adelberger, K. L., Dickinson, M., Giavalisco, M., & Pettini, M. 2001, *ApJ*, 562, 95  
 Songaila, A. 2001, *ApJ*, 561, L153  
 Verma, A., Lehnert, M. D., Förster Schreiber, N. M., Bremer, M. N., & Douglas, L. 2007, *MNRAS*, 377, 1024