

# Independent and Parallel Evolution of New Genes by Gene Duplication in Two Origins of C<sub>4</sub> Photosynthesis Provides New Insight into the Mechanism of Phloem Loading in C<sub>4</sub> Species

David M. Emms,<sup>1</sup> Sarah Covshoff,<sup>2</sup> Julian M. Hibberd,<sup>2</sup> and Steven Kelly<sup>\*,1</sup>

<sup>1</sup>Department of Plant Sciences, University of Oxford, Oxford, United Kingdom

<sup>2</sup>Department of Plant Sciences, University of Cambridge, Cambridge, United Kingdom

\*Corresponding author: E-mail: steven.kelly@plants.ox.ac.uk.

Associate editor: Michael Purugganan

## Abstract

C<sub>4</sub> photosynthesis is considered one of the most remarkable examples of evolutionary convergence in eukaryotes. However, it is unknown whether the evolution of C<sub>4</sub> photosynthesis required the evolution of new genes. Genome-wide gene-tree species-tree reconciliation of seven monocot species that span two origins of C<sub>4</sub> photosynthesis revealed that there was significant parallelism in the duplication and retention of genes coincident with the evolution of C<sub>4</sub> photosynthesis in these lineages. Specifically, 21 orthologous genes were duplicated and retained independently in parallel at both C<sub>4</sub> origins. Analysis of this gene cohort revealed that the set of parallel duplicated and retained genes is enriched for genes that are preferentially expressed in bundle sheath cells, the cell type in which photosynthesis was activated during C<sub>4</sub> evolution. Furthermore, functional analysis of the cohort of parallel duplicated genes identified SWEET-13 as a potential key transporter in the evolution of C<sub>4</sub> photosynthesis in grasses, and provides new insight into the mechanism of phloem loading in these C<sub>4</sub> species.

**Key words:** C<sub>4</sub> photosynthesis, gene duplication, gene families, parallel evolution.

## Introduction

The evolution of C<sub>4</sub> photosynthesis from ancestral C<sub>3</sub> photosynthesis involved multiple changes to biochemistry, subcellular compartmentalization, and leaf anatomy such that most modern-day C<sub>4</sub> species partition the entire photosynthetic process between two discrete cell types—bundle sheath (BS) and mesophyll (M) cells. This partitioning of the photosynthesis reactions between two cells facilitates an increase in CO<sub>2</sub> concentration around Rubisco leading to a reduction in photorespiration and a concomitant increase in photosynthetic efficiency (Hatch et al. 1967). Despite the numerous differences between C<sub>3</sub> and C<sub>4</sub> species, C<sub>4</sub> photosynthesis has evolved independently in over 60 different plant lineages (Sage et al. 2011) and thus represents one of the most widespread and successful examples of convergent evolution in eukaryotic biology.

Several complementary studies have identified patterns of gene expression associated with the evolution of C<sub>4</sub> photosynthesis. Comparisons of gene expression between closely related C<sub>3</sub> and C<sub>4</sub> species in the genera *Cleome* and *Flaveria* have identified ~3,500 gene expression changes associated with the evolutionary transition from C<sub>3</sub> to C<sub>4</sub> with smaller subsets of genes changing in parallel in multiple lineages (Gowik et al. 2011). Other studies identified genes with expression profiles that correlate with biochemical (Li et al. 2010; Majeran et al. 2010; Pick et al. 2011; Sage and Zhu 2011) or anatomical (Wang et al. 2013) features of C<sub>4</sub> species, or with biochemical differences between specialized C<sub>4</sub> cell types (Li et al. 2010; Chang et al. 2012; John et al. 2014). Further

studies have also identified parallel recruitment of transactors in two distantly related origins of C<sub>4</sub> (Aubry et al. 2014) and parallel recruitment of *cis*-elements to control C<sub>4</sub> gene expression (Brown et al. 2011). Although our understanding of gene expression changes associated with C<sub>4</sub> biology is rapidly increasing, almost nothing is known about the gene content differences that distinguish C<sub>4</sub> species from their C<sub>3</sub> relatives.

Individual gene and whole-genome duplications are a dominant feature of plant evolution and have been detected in all angiosperm lineages (Adams and Wendel 2005; Soltis et al. 2009). Although gene duplication is a stochastic process, retention of duplicate genes is affected by selection, and duplicates are frequently lost (Lynch and Conery 2000). In the case of whole-genome duplication events this loss process is known as fractionation (Thomas et al. 2006; Woodhouse et al. 2010). If retained, the duplicated genes may share the original gene function (subfunctionalization), or they may diverge and develop novel functions (neofunctionalization) (Ohno 1970; Lynch and Conery 2000; Moore and Purugganan 2005; Freeling et al. 2008). In maize, duplicated genes that share the same expression pattern evolve faster than those whose expression profiles diverge in time or space, and are more likely to be subject to positive selection (Hughes et al. 2014).

Although gene duplication has been studied in specific genes or at a genome scale in individual species such as *Arabidopsis thaliana*, *Glycine max*, and *Zea mays* (Blanc and Wolfe 2004; Duarte et al. 2006; Erdmann et al. 2010; Liu et al.

2011; Guo et al. 2013; Roulin et al. 2013; Hughes et al. 2014), it is unknown to what extent gene duplication events occur in parallel across species and to what extent these events underlie the emergence of complex convergent phenotypes. It is also unknown whether the evolution of  $C_4$  and the changes in photosynthetic competence of the bundle sheath and mesophyll cells, is associated with gene duplication. Analysis of gene expression data from different  $C_4$  species has led some authors to propose that neofunctionalization of genes following duplication has not played a major role in the evolution of  $C_4$  photosynthesis (Gowik and Westhoff 2011; Williams et al. 2012; van den Bergh et al. 2014). However, these analyses have primarily focused on the limited number of genes already known to be involved in  $C_4$  photosynthesis, and have not taken *ab initio* approaches to the discovery of duplicated genes associated with the evolution of  $C_4$  photosynthesis. Although gene duplication events associated with the evolution of  $C_4$  have yet to be identified, several lines of evidence show that gene duplications that predate the origin of  $C_4$  photosynthesis have played an important role in facilitating its emergence (Monson 1999, 2003; Sage 2004). Archetypal examples of this include specific members of the PEPC (phosphoenolpyruvate carboxylase), CA (carbonic anhydrase), and other gene families that are repeatedly co-opted for use in the  $C_4$  cycle in multiple independent  $C_4$  origins (Aldous et al. 2014; Christin et al. 2015).

Here we exploit recent advances in the ability to infer gene families and accurately localize gene duplication events (Boussau et al. 2013; Emms and Kelly 2015) to perform a genome-wide gene-tree species-tree reconciliation of all genes from seven monocot species that span two origins of  $C_4$  photosynthesis. We reveal that many  $C_4$  cycle genes duplicated in the common ancestor of the sampled *Poaceae* species, prior to the evolution of  $C_4$ . We show that subsequent co-option for  $C_4$  function of particular genes from these duplicate pairs was likely a result of expression divergence prior to the diversification of the *Poaceae*. We also provide an analysis of gene duplication associated with the evolution of  $C_4$  photosynthesis and reveal that there is significant independent, parallel duplication and subsequent retention of genes concurrent with the evolution of  $C_4$  photosynthesis. Furthermore, we show that genes that duplicated independently at both  $C_4$  origins are preferentially expressed in BS cells and comprise a suite of genes with novel functional significance for the evolution of  $C_4$  photosynthesis.

## New Approaches

Here we take a novel approach to elucidating the evolutionary changes that occurred in the transition from  $C_3$  to  $C_4$  photosynthesis in two independent lineages of  $C_4$  species. Through genome-wide gene-tree species-tree reconciliation, we reveal that hundreds of new genes evolved by gene duplication coincident with the evolution of  $C_4$  photosynthesis in both lineages. Furthermore, comparative analysis of these gene duplications reveals that there is commonality in the duplicated genes with a core set of 21 genes that duplicated and were retained independently in parallel in both lineages.


## Results

### Inference of Orthogroups and Identification of Gene Duplication Events


To provide the data for this analysis, a set of gene families (orthogroups) covering 42 sequenced plant genomes available in Phytozome V10 (Goodstein et al. 2012) was constructed (supplementary file S1 and fig. 1A, [Supplementary Material](#) online). Those orthogroups that contained sequences from monocot genomes were selected for further analysis (fig. 1). The set of monocots in this analysis comprise two independent origins of  $C_4$  photosynthesis (fig. 2A) (Aliscioni et al. 2012), an origin in the *Andropogoneae* of which *Z. mays* and *Sorghum bicolor* are descendants and an independent origin in the *Paniceae* of which *Setaria italica* and *Panicum virgatum* (DOE-JGI) are descendants. Although biochemical analyses suggested that *P. virgatum* and *Se. italica* were descendants of independent  $C_4$  origins (Hattersley and Watson 1992), the most recent phylogenetic analysis suggests that they share a single origin (Aliscioni et al. 2012). The independence of the  $C_4$  origins in this study is supported by the presence of a number of intervening clades containing  $C_3$  species including the *Arthropogoninae*, *Otachyriinae*, *Boivinellinae*, and *Neurachninae* (Aliscioni et al. 2012). The data set comprised 13,773 orthogroups and contained 248,266 genes (77.5% of all monocot genes) distributed across the 7 sampled monocot species. The phylogenetic position of individual gene duplication events contained within the 13,773 orthogroups were identified through construction of multiple sequence alignments, phylogenetic tree inference, and gene-tree species-tree reconciliation (fig. 1; see Methods). This resulted in the identification of 91,815 gene duplication events within the monocot phylogenetic tree. To identify retained gene duplication events the following two filtration criteria were applied: 1) A gene duplication event must have supporting evidence for its occurrence from at least two species and 2) all genes resulting from a gene duplication event must be retained in the genomes of all species that diverged after the duplication event. In total, 70,101 of the gene duplication events were on terminal branches of the species tree and so did not satisfy the first criterion. Of the remaining 21,714, a total of 3,782 gene duplications (distributed across the monocot phylogeny) satisfied the second criterion (fig. 2A), that is, that the genes descended from the gene duplication event were retained in all of the species that diverged after the gene duplication event. The complete set of phylogenetic trees for all these gene duplication events is provided in supplementary file S1, [Supplementary Material](#) online, and is available from the Dryad Digital Repository, <http://dx.doi.org/10.5061/dryad.17m42>. A Gene Ontology (GO) term enrichment analysis of the orthogroups in which these duplications occurred is provided in supplementary file S2, [Supplementary Material](#) online.

### Gene Duplication Events Occurred in Multiple Genes Associated with the $C_4$ Cycle

The filtered set of gene duplication events within the monocots contained six genes already known to be associated with



**FIG. 1.** Schematic overview of the gene duplication identification method used in this analysis. 1) Clustering genes into orthogroups using OrthoFinder. 2) Construction of multiple sequence alignments using MAFFT and inference of gene trees using Phyldog. 3) Reconciliation of gene trees with species tree using Phyldog. 4) Filtering of reconciled gene trees to identify gene duplication events that are retained in all species that evolved following the gene duplication event.



**FIG. 2.** (A) The number of fully retained gene duplication events that occurred along each nonterminal branch during the evolution of the monocot species in the study. The branches on which  $C_4$  evolved are indicated by green circles. (B) The  $C_4$  cycle associated genes that duplicated during this same period. Genes are depicted above the branch on which they duplicated. DIT2, dicarboxylate transporter 2; ME, NADP-dependent malic enzyme; PDK, pyruvate dehydrogenase kinase; TPI, triosephosphate isomerase; TPT, triosephosphate/phosphate translocator; PPDK-rp, pyruvate orthophosphate dikinase regulatory protein.


the  $C_4$  cycle (fig. 2B and supplementary fig. 1B, Supplementary Material online) (Langdale 2011). These comprise PPDK-rp, DIT2, NADP-ME, TPT, TPI, and a gene recently identified as upregulated in multiple  $C_4$  lineages, PDK (Brautigam et al. 2011, 2014) (fig. 3A–F). Of these, only pyruvate, orthophosphate dikinase regulatory protein (PPDK-rp) duplicated concurrently with an origin of  $C_4$  photosynthesis, in the ancestor of *So. bicolor* and *Z. mays* (figs. 2B and 3A). The remaining five duplicated in the ancestor of the *Poaceae* and three of these—NADP-ME2/3/4, PDK and DIT2—were duplicated in syntenic blocks associated with the  $\rho$  whole-genome duplication (Salse et al. 2008; Tang et al. 2010; Lee et al. 2013) while none showed evidence of being duplicated as part of the earlier  $\sigma$  or  $\tau$  whole-genome duplications that also occurred on this branch of the species tree (Tang et al. 2010; Jiao et al. 2014). Comparison of the expression profile of the PPDK-rp duplicates in *So. bicolor* and *Z. mays* with that of the single copy gene in *Se. italica* shows that subfunctionalization of this gene has occurred in the lineage leading to *So. bicolor* and *Z. mays* (fig. 3A). Here, the single PPDK-rp gene in *Se. italica* is expressed in both BS and M cells, while in *So. bicolor* and

*Z. mays* one of the duplicate pair is expressed in BS cells and the other is expressed in M cells (fig. 3A). All of the duplicate genes as well as the orthologous genes in *Se. italica*, *Oryza sativa*, and *Brachypodium distachyon* are predicted to encode chloroplast transit peptides. Although these duplicated genes now reside on different chromosomes in the fractionated *Z. mays* genome, they occur as a tandem pair in the genome of *So. bicolor* suggesting that they evolved by tandem duplication prior to the divergence of *So. bicolor* and *Z. mays*.

### Expression Divergence of Duplicate Genes Prior to $C_4$ Evolution Determined Recruitment to $C_4$ Function

A similar phenomenon is observed for the duplicates of glutamate/oxaloacetate malate exchange transporter, DIT2. Here two gene duplication events occurred, one prior to the divergence of the grasses and a second (not fully retained and hence not included in totals in fig. 2A) at the base of the *Panicoideae* (fig. 3B). In each species, one of the duplicated genes is expressed in M cells and another in BS cells (fig. 3B). The duplicate gene that is expressed in the BS in *So. bicolor* (Sobic.004G035500.1) and *Se. italica* (Si03501m) was lost from





**Fig. 3.** Expression profile of duplicate genes in bundle sheath and mesophyll cells of  $C_4$  species and in fully expanded foliar leaves of *Zea mays* and *Oryza sativa*. Only genes from species for which the corresponding data are available (BS/M and whole leaf expression) are shown in each tree. The complete trees including *Panicum virgatum* are included in supplementary file S1, **Supplementary Material** online, and are available from the Dryad Digital Repository, <http://dx.doi.org/10.5061/dryad.17m42>. Gene duplication events meeting our selection criteria are marked with a "D." Green and gray backgrounds distinguish genes descended from the two duplicates following the gene duplication event. Green bars indicate expression level in bundle sheath cells, blue bars indicate expression level in mesophyll cells. Gray bars indicate expression level in fully expanded leaves of *Z. mays* and *O. sativa*. TPM, transcripts per million. For explanation of gene abbreviations see legend to figure 2. PPKD-rp/2 duplicated in the ancestor of the *Andropogoneae*, the remainder duplicated in the ancestor of the *Poaceae*. (G) The number of duplication events in the ancestor of the *Poaceae* for which the most highly expressed of the duplicate genes in *Z. mays* is consistent (the same duplicate) or inconsistent (the other duplicate) with the most highly expressed of the duplicate genes in *O. sativa*. All duplication events for which one of the descendant genes in *O. sativa* and one of the descendant genes in *Z. mays* are both expressed at greater than 10 transcripts per million are included. For explanation of gene name abbreviations please see the legend to figure 2.

*Z. mays* and appears to be compensated for by BS-specific expression of the earlier duplicate (GRMZM2G086258) that originated at the base of the grasses (fig. 3B). The most parsimonious explanation for this phenomenon is that the gene used in  $C_4$  photosynthesis in *Z. mays* today was not the gene utilized for this function when  $C_4$  first evolved in this lineage. Furthermore, depending on the ancestral expression pattern of the single copy gene in the last common ancestor of the *Panicoideae*, the observed, divergent expression profile of the duplicate genes requires that either subfunctionalization or neofunctionalization of these duplicate genes occurred coincident with the evolution of  $C_4$  photosynthesis (fig. 3B).

For the remainder of the gene duplication events in  $C_4$  cycle genes that occurred prior to the divergence of the grasses (NADP-ME, TPT, TPI, and PDK), the same duplicate of the gene pair generated by the duplication event is expressed more highly in fully expanded leaves of both *O. sativa* (Covshoff et al. 2015) and *Z. mays* (Wang et al. 2013) (fig. 3C–F). These gene duplication events that occurred in

the ancestor of the grasses  $\geq 50$  Ma (Vicentini et al. 2008) predate the origin of  $C_4$  photosynthesis in the *Andropogoneae* and *Paniceae* by  $\geq 29$  My (Christin 2008). The fact that in each case the same duplicate gene was recruited for use in the  $C_4$  cycle in both independent origins raises the question as to whether in each case gene expression or biochemical changes modified the genes such that one duplicate was better suited for function in  $C_4$  photosynthesis. Comparison of the mRNA abundance of these duplicated genes in *Z. mays* and *O. sativa* revealed that in all cases the duplicated gene that was recruited for function in  $C_4$  photosynthesis was the ortholog of the one most highly expressed in mature rice leaves (fig. 3B–F) indicating that the duplicated genes had already undergone regulatory neofunctionalization prior to the divergence of the grass species in this study. To determine whether this pattern of duplicate gene recruitment was unique to  $C_4$  cycle genes, the mRNA abundance of all genes descended from gene duplication events that occurred prior to the divergence of the grasses were analyzed in mature leaves of *Z. mays* and

*O. sativa*. In the cases where both duplicate genes were expressed in fully expanded foliar leaves of both species, 74% ( $n = 893/1210$ ) of duplicate pairs exhibited a consistent expression pattern whereby the same gene of the duplicate pair was more highly expressed in the leaves of *Z. mays* and *O. sativa* (fig. 3G). Thus this recruitment phenomenon is not unique to  $C_4$  cycle genes, and it is more parsimonious to assume that expression differences, rather than biochemical differences, were the determining factors for recruitment to  $C_4$  function.

### Multiple Gene Duplication Events Occurred in the Same Gene Families at Both Origins of $C_4$

The retained gene duplication events that were localized to a branch that contained an origin of  $C_4$  were compared to determine whether there were any that occurred in parallel in both origins of  $C_4$ . Here a parallel gene duplication event is defined as a gene duplication that occurred independently to the same ortholog in both  $C_4$  lineages. The 535 gene duplications that occurred in the ancestor of the *Panicaceae* and the 248 in the ancestor of the *Andropogoneae* occurred in 381 and 225 orthogroups, respectively (supplementary file S1, [Supplementary Material](#) online), as multiple gene duplications occurred in some orthogroups. Of these gene duplications, 41 occurred independently to genes in the same orthogroup at both independent  $C_4$  origins (fig. 4A and supplementary file S3, [Supplementary Material](#) online). This overlap is significantly more than would be expected given the number of orthogroups and the number of gene duplication events on each branch (hypergeometric test  $P = 10^{-21}$ , expected number = 7). GO term analysis of this overlapping set of orthogroups revealed significant enrichment of multiple functional categories of genes. Of these, ~24% (10/42) had plausible association with  $C_4$  photosynthesis, including those categories concerning the biosynthesis of plant cell walls (Edwards and Voznesenskaya 2011), plasmodesmata (Botha 1992; Sowinski et al. 2007), and flavonols (Hannoufa et al. 1994), as well as those concerning leaf morphogenesis and development (Wang et al. 2013) (supplementary file S2, [Supplementary Material](#) online).

### Twenty-one Gene Duplication Events Occurred in the Same Genes at Both Origins of $C_4$


Of the 41 gene duplication events in the same orthogroup, 21 occurred in parallel whereby the same gene was duplicated at both  $C_4$  origins and was subsequently retained in all sampled descendant species (fig. 4A and supplementary file S3, [Supplementary Material](#) online). These 21 genes comprise two transcription factors, two transporters, two genes involved in hormone metabolism, two genes for redox homeostasis, six genes of unknown function, as well as one gene in each of the following categories: Cell wall biosynthesis, plastid import, flavonoid biosynthesis, lipid metabolism, plantacyanin, component of the ribosome, and signaling component (supplementary file S3, [Supplementary Material](#) online). The phylogenetic trees, BS and M cell expression profile, expression profile in fully expanded leaves of *Z. mays* and *O. sativa*, as well as the expression profile of the *Z. mays* genes

during leaf development are shown in supplementary figure 2A–U, [Supplementary Material](#) online. The complete set of phylogenetic trees for all of these gene duplication events is provided in supplementary file S1, [Supplementary Material](#) online, and is available from the Dryad Digital Repository, <http://dx.doi.org/10.5061/dryad.17m42>.


The evolution of  $C_4$  photosynthesis in these lineages involved the activation of photosynthesis in the BS. Given this, the BS and M cell expression of the genes in *So. bicolor*, *Se. italica* (John et al. 2014), and *Z. mays* (Chang et al. 2012) that were duplicated on branches that encompassed  $C_4$  was analyzed. Here, the proportion of duplicated genes that were preferentially expressed in BS or M cells was compared with the proportion of all genes preferentially expressed in these cell types in the same species. Analysis of these expression data revealed that there was no significant enrichment of BS or M preferentially expressed genes in the complete set of genes that duplicated at either  $C_4$  origin (fig. 4B). Neither was there a significant enrichment of BS or M preferentially expressed genes in the set of duplicate genes where the duplication event occurred in the same orthogroup (excluding those that occurred in the same gene; fig. 4C). However, there were significantly more BS preferentially expressed genes than expected in the set of genes that duplicated in parallel at both origins of  $C_4$  photosynthesis (fig. 4D). Of the parallel duplicated genes that show BS expression, ~45% of the descendant pairs are both preferentially expressed in the BS and ~50% show evidence of regulatory neofunctionalization where one of the duplicates is preferentially expressed in the BS and the other is either not preferentially expressed in either cell type or is preferentially expressed in M cells (supplementary fig. 2X, [Supplementary Material](#) online).

### Parallel Duplicated Genes Provide Insight into the Evolution of Phloem Loading in $C_4$

In  $C_4$  *Panacoid* grasses the plasmodesmata connections between BS and vascular parenchyma are significantly less numerous than those between M and BS (Botha 1992; Botha and Vanbel 1992; Sowinski et al. 2007). In addition, the plasmodesmata connections between the BS and the vascular parenchyma are significantly less numerous than those observed in related  $C_3$  species (Botha 1992; Botha and Vanbel 1992; Sowinski et al. 2007), and the mechanism by which sucrose is unloaded from BS cells into the vasculature is unknown. One of the two transporters that duplicated in parallel at both origins of  $C_4$  (fig. 5A) is the ortholog of the *O. sativa* SWEET-13 sucrose efflux transporter (Zhou et al. 2015). Although SWEET-13 in *O. sativa* is expressed to a low level in mature leaves (fig. 5B), the duplicate genes are highly expressed in whole leaves in *Z. mays* and are each among the most highly expressed genes in the BS (e.g., GRMZM2G173669, GRMZM2G179349, and GRMZM2G021706 are the 57th, 87th, and 166th most highly expressed genes in the BS respectively of 63,480 genes in *Z. mays*). It is possible, given the isolation protocol (Covshoff et al. 2013), that in *Se. italica* and *So. bicolor* some expression of SWEET-13 occurs in the vasculature rather than the BS. However, this is not the case for *Z. mays* where BS transcriptome data comes from mechanically isolated (Chang



**Fig. 4.** (A) Venn diagram showing the overlap between the orthogroups in which gene duplication events occurred. Number in parenthesis is the number of genes in which gene duplications happened to the same gene. (B) The expression profile in bundle sheath and mesophyll cells of all genes duplicated at the origins of  $C_4$  photosynthesis in this study. (C) The expression profile in bundle sheath and mesophyll cells of genes descended from paralogous duplication events (duplication in the same orthogroup in both origins). (D) The expression profile in bundle sheath and mesophyll cells of genes descended from orthologous duplication events (duplication in the same gene in both origins). Dark blue/green bars show the percentage of genes that are preferentially expressed in M/BS cells. Light blue/green bars show the percentage of all genes in each genome that are preferentially expressed in M/BS cells. \*\*\* $P < 0.001$ ; \* $P < 0.05$ .



**Fig. 5.** The SWEET-13 sucrose efflux transporter duplicated independently at both  $C_4$  origins. (A) The gene tree showing the independent, parallel gene duplication events. (B) The expression profile in bundle sheath (green bars) and mesophyll cells (blue bars) of each duplicate gene and a comparison of the whole leaf expression profile of each duplicate in *Zea mays* to the single copy ortholog in *Oryza sativa*. (C) The expression profile of each duplicate gene in *Z. mays* foliar (solid lines) and husk (dashed lines) leaf developmental gradient.

et al. 2012) and laser dissected BS cells (Li et al. 2010). Moreover, the genes are most highly expressed in fully expanded foliar leaves when compared with other developmental stages (fig. 5C).

### The Putative Functional Significance of Other Parallel Duplicated Genes

Of note is that 2 of the 21 parallel duplication events occurred in the same orthogroup of glutaredoxin genes (comprising four duplications in total). In total, 14 of the 17 genes descended from these duplication events are preferentially expressed in BS cells (supplementary fig. 2A and B,

Supplementary Material online). Moreover, all of the maize genes are expressed more highly in photosynthetically active tissue than in young developing tissue (supplementary fig. 2A and 2B, Supplementary Material online). Glutaredoxin genes are involved in antioxidant defense and stimulate photorespiratory glycine decarboxylase activity in plant mitochondria (Hoffmann et al. 2013). Thus given the expression of the duplicate genes in BS cells, it is possible that they function to protect the cell from oxidative damage as a result of activation of photosynthesis.

As described above, many of the parallel duplications occurred in genes of unknown function. Although some of



these duplicated genes have clues to function from the presence of conserved protein domains such as the parallel duplication that occurred in the ABC-type transporter (supplementary fig. 2S, [Supplementary Material](#) online), or the multiple parallel duplications that occurred in the glutaredoxin gene (above), many have no domains or other features that can provide clues to potential protein function. The most striking example of these duplications in unknown genes is that of the gene duplication events that happened in the orthogroup shown in supplementary figure 2L, [Supplementary Material](#) online. Here the majority of genes (14/20) are preferentially expressed in the BS cells of  $C_4$  species. Moreover, during *Z. mays* leaf development they are also more highly expressed in immature and fully expanded leaves than in earlier developmental stages. Although no clues to function can be obtained from the analysis of the protein sequence, it seems likely that the gene family provides some function to BS cells.

### The Transcription Factors That Duplicated in Parallel at Both Origins of $C_4$


In total, 40 transcription factors (*Andropogoneae* = 15, *Paniceae* = 25) duplicated at the 2 independent  $C_4$  origins in this study. Thus significant novel regulatory capacity evolved coincident with the evolution of  $C_4$  photosynthesis in these lineages. Of these 40 gene duplication events, 2 duplicated in parallel at both origins of  $C_4$  photosynthesis and the descendants of both are most highly expressed during early stages of leaf development (fig. 6C and F). One of the transcription factors that duplicated independently at both  $C_4$  origins is a SQUAMOSA promoter binding-like transcription factor (fig. 6A). Although neither the duplicate genes in the  $C_4$  species nor the related single copy gene in rice have been functionally characterized, other members of the gene family have been shown to modulate plant architecture and development (Jorgensen and Preston 2014). The duplicate genes are primarily expressed in early developmental stages of both foliar and husk leaves in *Z. mays* (fig. 6B and 6C). Analysis of the expression profile of the duplicate genes (GRMZM2G113779 and GRMZM2G067624) during leaf development in *Z. mays* shows that the duplicate gene pair has undergone regulatory neofunctionalization (fig. 6C). Moreover, one of the duplicates is highly expressed at the initial stages of foliar leaf development (fig. 6C) where Kranz anatomy is patterned (Wang et al. 2013). It is noteworthy that even though expression of these genes decreases as leaf tissue matures, in mature leaves the duplicate transcription factors still exhibit preferential expression in BS cells.

The other parallel duplicated transcription factor is an AP2/B3 transcription factor (fig. 6D and F). These duplicated transcription factor genes are also more highly expressed in developing leaf tissue as compared with fully expanded leaves, and comparison of the expression profile of the duplicate genes during foliar and husk leaf development in maize provides evidence of regulatory neofunctionalization in these leaf types (fig. 6F).

## Discussion

The evolution of  $C_4$  photosynthesis required numerous changes to plant anatomy and biochemistry such that many differences distinguish  $C_4$  species from their  $C_3$  relatives. Although a significant body of knowledge has amassed concerning genes that are differentially expressed between  $C_4$  and  $C_3$  species, it is unknown whether the evolution of  $C_4$  photosynthesis was concomitant with the evolution of new genes. Here we have shown that hundreds of new genes evolved by gene duplication coincident with the evolution of  $C_4$  photosynthesis and were subsequently retained in two independent lineages of  $C_4$  species, revealing a previously hidden complexity to  $C_4$  evolution. Although any of these gene duplication events may be important to the evolution of  $C_4$ , many are likely linked to other traits that evolved during the same period captured by the branch in the phylogenetic tree on which  $C_4$  evolved. Thus, to identify the genes most likely to be associated with  $C_4$  evolution, the sets of genes that duplicated at both origins and were retained in the descendant species were assessed for commonality. This revealed that many retained gene duplication events occurred within the same orthogroup (gene family) at both independent  $C_4$  origins. Furthermore, the analysis identified 21 genes that duplicated independently in parallel and were retained in the descendant species. The correspondence between the genes that duplicated in parallel at both  $C_4$  origins and their preferential expression in BS cell types implicates these genes in  $C_4$  function, and is consistent with the hypothesis that more biochemical changes happened to BS cells as compared with M cells during  $C_4$  evolution (Langdale 2011).

Of the genes that duplicated in parallel at both  $C_4$  origins, several have plausible functional significance for the evolution of  $C_4$ . Of particular note is the independent duplication of the SWEET-13 sucrose efflux transporter. In this case, although the orthologous single copy gene in *O. sativa* is expressed to a low level in mature leaves, the duplicate genes in the  $C_4$  lineages are each among the most highly expressed genes in the BS irrespective of whether BS was isolated as strands of vasculature (*Se. italic* [John et al. 2014], *So. bicolor* [presented here]) or as mechanically isolated cells (*Z. mays*; Chang et al. 2012) or laser dissected from leaf sections (*Z. mays*; Li et al. 2010). We propose that these duplicated transporters function to export sucrose out of the BS into the vasculature apoplast. This would be advantageous in  $C_4$  species where the number of plasmodesmata connections between BS and vascular parenchyma are reduced compared with those between the BS and M (Botha 1992; Botha and Vanbel 1992; Sowinski et al. 2007). In this model, the exported sucrose would be taken up by active proton-coupled sucrose transporter SUC2/SUT1 in the sieve elements and companion cells (fig. 7). Four lines of evidence support the role of these gene duplicates in  $C_4$  photosynthesis. First, the SWEET-13 gene duplicated in parallel in both independent origins of  $C_4$  photosynthesis in this analysis. Second, all gene duplicates are preferentially expressed in the BS of these  $C_4$  species. Third, expression of each of these duplicated genes is substantially higher in *Z. mays* than the related single copy ortholog in




**Fig. 6.** Two transcription factors duplicated independently at both C<sub>4</sub> origins. (A) SQUAMOSA promoter binding-like transcription factor, A–C; an AP2/B3 transcription factor, D–F. (A and D) The expression profile in bundle sheath (green bars) and mesophyll cells (blue bars) of each duplicate gene. (B and E) A comparison of the whole leaf expression profile of each duplicate in *Zea mays* to the single copy ortholog in *O. sativa*. (C and F) The expression profile of each duplicate gene in *Z. mays* foliar (solid lines) and husk (dashed) leaf developmental gradient.

*O. sativa* (C<sub>3</sub>). Finally, these duplicate genes are among the highest expressed genes in the BS.

Also of note among the retained, parallel duplicated genes are the duplicates of the SQUAMOSA promoter binding-like transcription factor. The parallel duplication of this gene, combined with the neofunctionalization of one duplicate in Kranz tissue, suggest that GRMZM2G067624 may function in early vascular patterning and contribute to the patterning of Kranz anatomy in these C<sub>4</sub> grasses.

It has previously been proposed that the evolution of C<sub>4</sub> photosynthesis did not rely on duplication and neofunctionalization of genes (Gowik and Westhoff 2011; Williams et al. 2012; van den Bergh et al. 2014;). However, this proposal was based on the analysis of the subset of genes already known to be involved in C<sub>4</sub> photosynthesis and thus there was potential for many such events to be overlooked. Here we show that there are hundreds of examples of gene duplication associated with the evolution of C<sub>4</sub> photosynthesis in two separate C<sub>4</sub> lineages. Moreover, we show that in at least one case a gene duplication event occurred in a gene known to be associated with C<sub>4</sub> photosynthesis. This occurred in the chloroplast targeted *PPDK-rp* gene in the ancestor of *So. bicolor* and *Z. mays*. Given the expression profile of the genes descended from this duplication event and the expression profile of the single copy ortholog in *Se. italica*, it appears that this duplicate pair has undergone subfunctionalization with one duplicate preferentially expressed in BS cells and the other expressed in M cells. Thus it is likely that gene



**Fig. 7.** Model for localization and function of duplicated SWEET-13 genes in C<sub>4</sub> photosynthesis. CC is companion cell, SE is sieve element, VP is vascular parenchyma, pyr is pyruvate, mal is malate, and suc is sucrose. Dashed arrows indicate multistep biochemical pathways. Solid arrows indicate movement of labeled compound.

duplication and subfunctionalization played a role in the evolution of C<sub>4</sub> photosynthesis in this lineage.

In agreement with previous investigations, this study revealed that several C<sub>4</sub> cycle genes duplicated prior to the evolution of C<sub>4</sub> photosynthesis. In all cases it appears that the same duplicate gene was co-opted for C<sub>4</sub> function in both independent C<sub>4</sub> lineages. Genome-wide analysis of the expression patterns of all duplicate genes that evolved on the same branch revealed that this concordance in co-option is most likely a function of expression divergence that



happened prior to the diversification of the *Poaceae*. That is, the duplicate copy that is most highly expressed in leaf tissue as a result of expression divergence prior to the evolution of  $C_4$  is the one most likely to be co-opted for  $C_4$  function. Given that 74% of duplicate genes display concordant patterns of expression dominance, it will be interesting to see whether this global trend in gene co-option holds true as more  $C_4$  cycle genes are discovered.

Taken together this study reveals that in addition to previously documented parallelism in gene expression changes between  $C_3$  and  $C_4$  species (Gowik et al. 2011; Aubry et al. 2014), and parallelism in recruitment of *cis*-elements (Brown et al. 2011), there is significant parallelism in the evolution of new genes by gene duplication coincident with the evolution of  $C_4$ . Furthermore, functional analysis of these newly evolved genes provides new insight into the biology of  $C_4$  species and suggests that duplication and neofunctionalization may have played a key role in the evolution of  $C_4$  photosynthesis.

## Materials and Methods

### Construction of Orthologous Gene Groups, Gene-Tree Species-Tree Reconciliation, and Filtration of Duplication Events

Protein primary transcripts were downloaded for the 41 species (supplementary file S1, [Supplementary Material](#) online) from Phytozome 10 (Goodstein et al. 2012) with genomes not under embargo in July 2014. The *Brassica rapa* database version of the *Brassica rapa* genome (Cheng et al. 2011) was used in place of the Phytozome version as there were restrictions on the use of the later. Orthologous gene groups (orthogroups) were inferred using OrthoFinder (Emms and Kelly 2015). Phyldog (Boussau et al. 2013) was used to simultaneously infer the gene trees for each orthogroup and reconcile them with the species phylogeny (supplementary fig. S1A, [Supplementary Material](#) online). The topology of the species phylogeny for the angiosperms was obtained from the Angiosperm Phylogeny (Stevens 2001). The topology of the tree for the remaining species was obtained from Phytozome. This topology was chosen as it represents the current consensus tree for the angiosperms and is supported by a number of recent phylogenies for plants (Stevens 2001; Bremer et al. 2009; Ruhfel et al. 2014). It is worth noting that the relationships between a small number of eudicot clades in the Phytozome tree disagree with this topology. However, these conflicting eudicot branches within the Phytozome tree are distant from the branches studied within the monocots, and thus use of the Phytozome tree would be unlikely to affect the results of the analysis. To estimate the molecular sequence branch lengths of this species phylogeny, a concatenated alignment of single copy gene orthogroups that each had representation of  $\geq 75\%$  of the species ( $n = 1,230$ ) was constructed and FastTree (Price et al. 2010) was used to estimate branch lengths given the constrained species topology.

To identify the phylogenetic position of gene duplication events, the multiple sequence alignments for the amino acid sequences of each orthogroup were inferred using mafft-linsi (Katoh and Standley 2013) and the alignments subject to

gene-tree to species-tree reconciliation using Phyldog and the known species tree described above. To identify retained gene duplication events the following two filtration criteria were applied: 1) A gene duplication event must have supporting evidence for its occurrence from at least two species and 2) all genes resulting from a gene duplication event must be retained in the genomes of all species that diverged after the duplication event. Thus all gene duplication events analyzed in this work are fully retained in all descendent species that evolved after the gene duplication event. The gene trees for all the gene duplication events are available from the Dryad Digital Repository, <http://dx.doi.org/10.5061/dryad.17m42>.

### Ontology Enrichment Analysis

GO (Ashburner et al. 2000) annotations for each orthogroup were taken from the annotations of the *A. thaliana* genes (Berardini et al. 2004) within the orthogroup. Enrichment analysis for each ontology term was performed using the hypergeometric test on the number of orthogroups with that term within a specific set (e.g., the set of orthogroups that expanded on a particular branch) given the number of orthogroups with that term in the population as a whole (e.g., the set of orthogroups containing a gene that was present on the particular branch). Correction for multiple comparisons was performed using the Benjamini and Hochberg procedure (Benjamini et al. 2001) with a false discovery rate cut-off of 0.05.

### Plant Growth and RNA Extraction

*Sorghum bicolor* was grown at the University of Cambridge at 28°C in a 16 h photoperiod under 400  $\mu\text{mol}/\text{m}^2/\text{s}$  photon flux density at 40% humidity as previously described (Covshoff et al. 2013). BS strands of *So. bicolor* were isolated by mechanical separation and M cell contents were collected by leaf rolling as previously described (Covshoff et al. 2013). In total, three biological replicates were collected for each cell type. Each biological replicate contained pooled RNA from the third leaf of 16 individual plants. Total RNA was extracted from the BS and M samples using the standard protocol of the mirVana micro RNA Isolation kit (Ambion).

### RNA Sequencing

RNA-seq libraries were prepared from 1  $\mu\text{g}$  of total RNA (TruSeq RNA sample preparation version 2 guide; Illumina). Six libraries (comprising the three biological replicates from each cell type) were sequenced with TruSeq version 3 chemistry using one lane of the HiSeq 2000 to generate approximately 161 million 100-bp paired-end reads. The full set of raw reads from each biological replicate was deposited to EBI array express under the accession number E-MTAB-4021.

### Gene Expression Analysis

Transcriptomes from bundle sheath and mesophyll cells for *S. italica* (John et al. 2014), *So. bicolor* (described above), and *Z. mays* (Chang et al. 2012) were used to investigate the expression of duplicated genes in these specialized  $C_4$  cell types. In each case, paired-end reads were subject to quality trimming and adaptor filtering using Trimmomatic (Bolger

et al. 2014) using the settings “LEADING:10 TRAILING:10 SLIDINGWINDOW:5:15 MINLEN:50.” The quality filtered paired-end reads were then mapped to the complete set of representative transcripts from Phytozome version 10 of the *Z. mays*, *So. bicolor*, and *Se. italica* using bowtie2 (Langmead and Salzberg 2012) and transcript abundances were estimated using Express (Roberts and Pachter 2013). All pairwise comparisons between developmental stages were made using DESeq (Anders and Huber 2010), using the default normalization method and identifying differentially expressed genes as those with a Benjamini–Hochberg corrected  $P \leq 0.05$  (Benjamini and Hochberg 1995).

## Supplementary Material

Supplementary figures S1 and S2 and files S1–S3 are available at *Molecular Biology and Evolution* online (<http://www.mbe.oxfordjournals.org/>).

## Acknowledgments

We thank the Department of Energy Joint Genome Institute for pre-publication access to the V1.0 *Panicum virgatum* genome. This project has received funding from the European Union’s Horizon 2020 research and innovation program under grant agreement no. 637765. S.K. is a Royal Society University Research Fellow. The work was also funded by the C<sub>4</sub> rice project grant to the University of Oxford from the Bill and Melinda Gates Foundation.

## References

- Adams KL, Wendel JF. 2005. Polyploidy and genome evolution in plants. *Curr Opin Plant Biol.* 8:135–141.
- Aldous SH, Weise SE, Sharkey TD, Waldera-Lupa DM, Stuhler K, Mallmann J, Groth G, Gowik U, Westhoff P, Arsova B. 2014. Evolution of the phosphoenolpyruvate carboxylase protein kinase family in C-3 and C-4 Flaveria spp. *Plant Physiol.* 165:1076–1091.
- Aliscioni S, Bell HL, Besnard G, Christin PA, Columbus JT, Duvall MR, Edwards EJ, Giussani L, Hasenstab-Lehman K, Hilu KW, et al. 2012. New grass phylogeny resolves deep evolutionary relationships and discovers C<sub>4</sub> origins. *New Phytol.* 193:304–312.
- Anders S, Huber W. 2010. Differential expression analysis for sequence count data. *Genome Biol.* 11:R106.
- Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, et al. 2000. Gene ontology: tool for the unification of biology. *Nat Genet.* 25:25–29.
- Aubry S, Kelly S, Kumpers BMC, Smith-Unna RD, Hibberd JM. 2014. Deep evolutionary comparison of gene expression identifies parallel recruitment of trans-factors in two independent origins of C-4 photosynthesis. *PLoS Genet.* 10:e1004365.
- Benjamini Y, Drai D, Elmer G, Kafkafi N, Golani I. 2001. Controlling the false discovery rate in behavior genetics research. *Behav Brain Res.* 125:279–284.
- Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate—a practical and powerful approach to multiple testing. *J R Stat Soc B* 57:289–300.
- Berardini TZ, Mundodi S, Reiser L, Huala E, Garcia-Hernandez M, Zhang PF, Mueller LA, Yoon J, Doyle A, Lander G, et al. 2004. Functional annotation of the *Arabidopsis* genome using controlled vocabularies. *Plant Physiol.* 135:745–755.
- Blanc G, Wolfe KH. 2004. Functional divergence of duplicated genes formed by polyploidy during *Arabidopsis* evolution. *Plant Cell* 16:1679–1691.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120.
- Botha CEJ. 1992. Plasmodesmatal distribution, structure and frequency in relation to assimilation in C3 and C4 grasses in southern Africa. *Planta* 187:348–358.
- Botha CEJ, Vanbel AJE. 1992. Quantification of symplastic continuity as visualized by plasmodesmograms—diagnostic-value for phloem-loading pathways. *Planta* 187:359–366.
- Boussau B, Szollosi GJ, Duret L, Gouy M, Tannier E, Daubin V. 2013. Genome-scale coestimation of species and gene trees. *Genome Res.* 23:323–330.
- Brautigam A, Kajala K, Wullenweber J, Sommer M, Gagneul D, Weber KL, Carr KM, Gowik U, Mass J, Lercher MJ, et al. 2011. An mRNA blueprint for C-4 photosynthesis derived from comparative transcriptomics of closely related C-3 and C-4 species. *Plant Physiol.* 155:142–156.
- Brautigam A, Schliesky S, Kulahoglu C, Osborne CP, Weber APM. 2014. Towards an integrative model of C-4 photosynthetic subtypes: insights from comparative transcriptome analysis of NAD-ME, NADP-ME, and PEP-CK C-4 species. *J Exp Bot.* 65:3579–3593.
- Bremer B, Bremer K, Chase MW, Fay MF, Reveal JL, Soltis DE, Soltis PS, Stevens PF, Anderberg AA, Moore MJ, et al. 2009. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG III. *Bot J Linn Soc.* 161:105–121.
- Brown NJ, Newell CA, Stanley S, Chen JE, Perrin AJ, Kajala K, Hibberd JM. 2011. Independent and parallel recruitment of preexisting mechanisms underlying C-4 photosynthesis. *Science* 331:1436–1439.
- Chang YM, Liu WY, Shih ACC, Shen MN, Lu CH, Lu MYJ, Yang HW, Wang TY, Chen SCC, Chen SM, et al. 2012. Characterizing regulatory and functional differentiation between maize mesophyll and bundle sheath cells by transcriptomic analysis. *Plant Physiol.* 160:165–177.
- Cheng F, Liu SY, Wu J, Fang L, Sun SL, Liu B, Li PX, Hua W, Wang XW. 2011. BRAD, the genetics and genomics database for Brassica plants. *BMC Plant Biol.* 11:136.
- Christin PA, Arakaki M, Osborne CP, Edwards EJ. 2015. Genetic enablers underlying the clustered evolutionary origins of C<sub>4</sub> photosynthesis in angiosperms. *Mol Biol Evol.* 32:846–858.
- Covshoff S, Furbank RT, Leegood RC, Hibberd JM. 2013. Leaf rolling allows quantification of mRNA abundance in mesophyll cells of sorghum. *J Exp Bot.* 64:807–813.
- Covshoff S, Hughes TE, Smith-Unna R, Kelly S, Szczewka M, Bailey KJ, Sage TL, Pachebat JA, Leegood RC, Hibberd JM. 2015. C<sub>4</sub> photosynthesis in the rice paddy: insights from the noxious weed *Echinochloa glabrescens*. *Plant Physiol.* 170:57–73.
- Christin PA, Besnard G, Samaritani E, Duvall MR, Hodkinson TR, Savolainen V, Salamin N. 2008. Oligocene CO<sub>2</sub> decline promoted C<sub>4</sub> photosynthesis in grasses. *Curr Biol.* 18:37–43.
- Duarte JM, Cui LY, Wall PK, Zhang Q, Zhang XH, Leebens-Mack J, Ma H, Altman N, dePamphilis CW. 2006. Expression pattern shifts following duplication indicative of subfunctionalization and neofunctionalization in regulatory genes of *Arabidopsis*. *Mol Biol Evol.* 23:469–478.
- Edwards GE, Voznesenskaya EV. 2011. C-4 photosynthesis: Kranz forms and single-cell C-4 in terrestrial plants. In: Raghavendra AP, Sage RF, editors. *C<sub>4</sub> photosynthesis and related CO<sub>2</sub> concentrating mechanisms*. Heidelberg (Germany): Springer. p. 29–61.
- Emms DM, Kelly S. 2015. OrthoFinder: solving fundamental biases in whole genome comparisons dramatically improves orthogroup inference accuracy. *Genome Biol.* 16:157.
- Erdmann R, Gramzow L, Melzer R, Theissen G, Becker A. 2010. GORDITA (AGL63) is a young paralog of the *Arabidopsis thaliana* B-sister MADS box gene ABS (TT16) that has undergone neofunctionalization. *Plant J.* 63:914–924.
- Freeling M, Lyons E, Pedersen B, Alam M, Ming R, Lisch D. 2008. Many or most genes in *Arabidopsis* transposed after the origin of the order Brassicales. *Genome Res.* 18:1924–1937.
- Goodstein DM, Shu SQ, Howson R, Neupane R, Hayes RD, Fazo J, Mitros T, Dirks W, Hellsten U, Putnam N, Rokhsar DS. 2012. Phytozome: a comparative platform for green plant genomics. *Nucleic Acids Res.* 40:D1178–D1186.

- Gowik U, Brautigam A, Weber KL, Weber APM, Westhoff P. 2011. Evolution of C-4 photosynthesis in the genus *Flaveria*: how many and which genes does it take to make C-4? *Plant Cell* 23:2087–2105.
- Gowik U, Westhoff P. 2011. The path from C-3 to C-4 photosynthesis. *Plant Physiol.* 155:56–63.
- Guo H, Lee TH, Wang XY, Paterson AH. 2013. Function relaxation followed by diversifying selection after whole-genome duplication in flowering plants. *Plant Physiol.* 162:769–778.
- Hannoufa A, Brown RH, Ibrahim RK. 1994. Variations in flavonoid sulfate patterns in relation to photosynthetic types of 5 flaveria species. *Phytochemistry* 36:353–356.
- Hatch MD, Slack CR, Johnson HS. 1967. Further studies on a new pathway of photosynthetic carbon dioxide fixation in sugar-cane and its occurrence in other plant species. *Biochem J.* 102:417.
- Hattersley PW, Watson L. 1992. Diversification of photosynthesis. In: Chapman GP, editor. *Grass evolution and domestication*. Cambridge: Cambridge University Press. p. 38–116.
- Hoffmann C, Plochanski B, Haferkamp I, Lerach M, Ewald R, Bauwe H, Riemer J, Herrmann JM, Neuhaus HE. 2013. From endoplasmic reticulum to mitochondria: absence of the *Arabidopsis* ATP antiporter endoplasmic reticulum adenylate transporter1 perturbs photorespiration. *Plant Cell* 25:2647–2660.
- Hughes TE, Langdale JA, Kelly S. 2014. The impact of widespread regulatory neofunctionalization on homeolog gene evolution following whole-genome duplication in maize. *Genome Res.* 24:1348–1355.
- Jiao YN, Li JP, Tang HB, Paterson AH. 2014. Integrated syntenic and phylogenomic analyses reveal an ancient genome duplication in monocots. *Plant Cell* 26:2792–2802.
- John CR, Smith-Unna RD, Woodfield H, Covshoff S, Hibberd JM. 2014. Evolutionary convergence of cell-specific gene expression in independent lineages of C-4 grasses. *Plant Physiol.* 165:62–75.
- Jorgensen SA, Preston JC. 2014. Differential SPL gene expression patterns reveal candidate genes underlying flowering time and architectural differences in *Mimulus* and *Arabidopsis*. *Mol Phylogenet Evol.* 73:129–139.
- Katoh K, Standley DM. 2013. MAFFT: Multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol.* 30:772–780.
- Langdale JA. 2011. C-4 cycles: past, present, and future research on C-4 photosynthesis. *Plant Cell* 23:3879–3892.
- Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods.* 9:357–359.
- Lee TH, Tang HB, Wang XY, Paterson AH. 2013. PGDD: a database of gene and genome duplication in plants. *Nucleic Acids Res.* 41:D1152–D1158.
- Li PH, Ponnala L, Gandotra N, Wang L, Si YQ, Tausta SL, Kebrom TH, Provart N, Patel R, Myers CR, et al. 2010. The developmental dynamics of the maize leaf transcriptome. *Nat Genet.* 42:1060–1067.
- Liu SL, Baute GJ, Adams KL. 2011. Organ and cell type-specific complementary expression patterns and regulatory neofunctionalization between duplicated genes in *Arabidopsis thaliana*. *Genome Biol Evol.* 3:1419–1436.
- Lynch M, Conery JS. 2000. The evolutionary fate and consequences of duplicate genes. *Science* 290:1151–1155.
- Majeran W, Friso G, Ponnala L, Connolly B, Huang MS, Reidel E, Zhang CK, Asakura Y, Bhuiyan NH, Sun Q, et al. 2010. Structural and metabolic transitions of C-4 leaf development and differentiation defined by microscopy and quantitative proteomics in maize. *Plant Cell* 22:3509–3542.
- Monson RK. 1999. The origins of C4 genes and evolutionary pattern in the C4 metabolic phenotype. In: Sage RF, Monson RK, editors. *C4 plant biology*. San Diego (CA): Academic Press. p. 377–410.
- Monson RK. 2003. Gene duplication, neofunctionalization, and the evolution of C-4 photosynthesis. *Int J Plant Sci.* 164:S43–S54.
- Moore RC, Purugganan MD. 2005. The evolutionary dynamics of plant duplicate genes. *Curr Opin Plant Biol.* 8:122–128.
- Ohno S. 1970. *Evolution by gene duplication*. London: Allen & Unwin.
- Panicum virgatum v1.0, DOE-JGI. Available from: <http://phytozome.jgi.doe.gov/>.
- Pick TR, Brautigam A, Schluter U, Denton AK, Colmsee C, Scholz U, Fahnstich H, Pieruschka R, Rascher U, Sonnewald U, et al. 2011. Systems analysis of a maize leaf developmental gradient redefines the current C-4 model and provides candidates for regulation. *Plant Cell* 23:4208–4220.
- Price MN, Dehal PS, Arkin AP. 2010. FastTree 2—approximately maximum-likelihood trees for large alignments. *PLoS One* 5:E9490.
- Roberts A, Pachter L. 2013. Streaming fragment assignment for real-time analysis of sequencing experiments. *Nat Methods.* 10:71–73.
- Roulin A, Auer PL, Libault M, Schlueter J, Farmer A, May G, Stacey G, Doerge RW, Jackson SA. 2013. The fate of duplicated genes in a polyploid plant genome. *Plant J.* 73:143–153.
- Ruhfel BR, Gitzendanner MA, Soltis PS, Soltis DE, Burleigh JG. 2014. From algae to angiosperms inferring the phylogeny of green plants (Viridiplantae) from 360 plastid genomes. *BMC Evol Biol.* 14:23.
- Sage RF. 2004. The evolution of C-4 photosynthesis. *New Phytol.* 161:341–370.
- Sage RF, Christin PA, Edwards EJ. 2011. The C-4 plant lineages of planet Earth. *J Exp Bot.* 62:3155–3169.
- Sage RF, Zhu XG. 2011. Exploiting the engine of C-4 photosynthesis. *J Exp Bot.* 62:2989–3000.
- Salse J, Bolot S, Throude M, Jouffe V, Piegu B, Quraishi UM, Calcagno T, Cooke R, Delseny M, Feuillet C. 2008. Identification and characterization of shared duplications between rice and wheat provide new insight into grass genome evolution. *Plant Cell* 20:11–24.
- Soltis DE, Albert VA, Leebens-Mack J, Bell CD, Paterson AH, Zheng CF, Sankoff D, dePamphilis CW, Wall PK, Soltis PS. 2009. Polyploidy and angiosperm diversification. *Am J Bot.* 96:336–348.
- Sowinski P, Bilska A, Baranska K, Fronk J, Kobus P. 2007. Plasmodesmata density in vascular bundles in leaves of C4 grasses grown at different light conditions in respect to photosynthesis and photosynthate export efficiency. *Environ Exp Bot* 61:74–84.
- Stevens P. 2001. Angiosperm phylogeny website.
- Tang HB, Bowers JE, Wang XY, Paterson AH. 2010. Angiosperm genome comparisons reveal early polyploidy in the monocot lineage. *Proc Natl Acad Sci U S A.* 107:472–477.
- Thomas BC, Pedersen B, Freeling M. 2006. Following tetraploidy in an *Arabidopsis* ancestor, genes were removed preferentially from one homeolog leaving clusters enriched in dose-sensitive genes. *Genome Res.* 16:934–946.
- van den Bergh E, K  lahoglu C, Br  utigam A, Hibberd JM, Weber AP, Zhu XG, Schranz ME. 2014. Gene and genome duplications and the origin of C 4 photosynthesis: Birth of a trait in the Cleomaceae. *Curr Plant Biol.* 1:2–9.
- Vicentini A, Barber JC, Aliscioni SS, Giussani LM, Kellogg EA. 2008. The age of the grasses and clusters of origins of C(4) photosynthesis. *Glob Change Biol.* 14:2963–2977.
- Wang P, Kelly S, Fouracre JP, Langdale JA. 2013. Genome-wide transcript analysis of early maize leaf development reveals gene cohorts associated with the differentiation of C-4 Kranz anatomy. *Plant J.* 75:656–670.
- Williams BP, Aubry S, Hibberd JM. 2012. Molecular evolution of genes recruited into C-4 photosynthesis. *Trends Plant Sci.* 17:213–220.
- Woodhouse MR, Schnable JC, Pedersen BS, Lyons E, Lisch D, Subramaniam S, Freeling M. 2010. Following tetraploidy in maize, a short deletion mechanism removed genes preferentially from one of the two homeologs. *PLoS Biol.* 8:e1000409.
- Zhou JH, Peng Z, Long JY, Sossso D, Liu B, Eom JS, Huang S, Liu SZ, Cruz CV, Frommer WB, et al. 2015. Gene targeting by the TAL effector PthXo2 reveals cryptic resistance gene for bacterial blight of rice. *Plant J.* 82:632–643.