

Current Challenges in the Management of Sepsis in ICUs in Resource-poor Settings and Suggestions for the Future

Marcus J. Schultz^{1,2}, Martin W. Dunser³, Arjen M. Dondorp^{1,2}, Neill K.J. Adhikari⁴, Shivakumar Iyer⁵, Arthur Kwizera⁶, Yoel Lubell², Alfred Papali⁷, Luigi Pisani^{1,2}, Beth D. Riviello⁸, Derek C. Angus⁹, Luciano C. Azevedo¹⁰, Timothy Baker¹¹, Janet V. Diaz¹², Emir Festic¹³, Rashan Haniffa¹, Randeep Jawa¹⁴, Shevin T. Jacob¹⁵, Niranjan Kisson¹⁶, Rakesh Lodha¹⁷, Ignacio Martin-Loeches¹⁸, Ganbold Lundeg¹⁹, David Misango²⁰, Mervyn Mer²¹, Sanjib Mohanty²², Srinivas Murthy¹⁶, Ndidiamaka Musa²³, Jane Nakibuuka⁶, Ary Serpa Neto^{2,24}, Mai Nguyen Thi Hoang²⁵, Binh Nguyen Thien²⁶, Rajyabardhan Pattnaik²², Jason Phua²⁷, Jacobus Preller²⁸, Pedro Povoas²⁹, Suchitra Ranjit³⁰, Daniel Talmor⁸, Jonarthan Thevanayagam³¹, C. Louise Thwaites³² for the Global Intensive Care Working Group of the European Society of Intensive Care Medicine

Author affiliations:

- ¹Mahidol University, Bangkok, Thailand
²Academic Medical Center & University of Amsterdam, Amsterdam, The Netherlands
³University College of London Hospital, London, UK
⁴Sunnybrook Health Sciences Centre & University of Toronto, Toronto, ON, Canada
⁵Bharati Vidyapeeth Deemed University Medical College, Pune, India
⁶Mulago National Referral Hospital, Kampala, Uganda
⁷University of Maryland School of Medicine, Baltimore, MD
⁸Beth Israel Deaconess Medical Center and Harvard Medical School, Boston, MA
⁹University of Pittsburgh, Pittsburgh, PA
¹⁰Hospital Sirio-Libanes, São Paulo, Brazil
¹¹Karolinska Institute, Stockholm, Sweden
¹²California Pacific Medical Center, San Francisco, CA
¹³Mayo Clinic, Jacksonville, FL, USA
¹⁴Stony Brook University Medical Center, Stony Brook, NY, USA
¹⁵University of Washington, Seattle, WA
¹⁶British Columbia Children's Hospital, Vancouver, Canada
¹⁷All India Institute of Medical Science, Delhi, India
¹⁸St. James's University Hospital, Dublin, Ireland
¹⁹Mongolian National University of Medical Sciences, Ulaanbaatar, Mongolia
²⁰Aga Khan University Hospital, Nairobi, Kenya
²¹Johannesburg Hospital and University of the Witwatersrand, Johannesburg, South Africa
²²Ispat General Hospital, Rourkela, Sundargarh, Odisha, India

- ²³Seattle Children's Hospital and University of Washington, WA
²⁴Medical Intensive Care Unit, Hospital Israelita Albert Einstein, Sao Paulo, Brazil
²⁵Oxford University Clinical Research Unit, Hospital for Tropical Diseases, Ho Chi Minh City, District 5, Vietnam
²⁶Trung Vuong Hospital, Ho Chi Minh City, Vietnam
²⁷National University Hospital, Singapore
²⁸Addenbrooke's Hospital, Cambridge University Hospitals NHS Foundation Trust, Cambridge, UK
²⁹Nova Medical School, CEDOC, New University of Lisbon & Hospital de Sao Francisco Xavier, Centro Hospitalar de Lisboa Ocidental, Lisbon, Portugal
³⁰Appolo Hospitals, Chennai, India
³¹Mzuzu Central Hospital, Mzuzu, Malawi
³²Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine, University of Oxford, UK

Correspondence:

Prof. Marcus J. Schultz
Department of Intensive Care
Academisch Medical Center
Meibergdreef 9
1105 AZ Amsterdam
The Netherlands
Email: marcus.j.schultz@gmail.com

Take home message: ICUs in low- and middle-income countries face many challenges that could affect patient outcome. Addressing both disease-specific and setting-specific factors is important to improve performance in these settings. Detailed evaluation at both at a macro- and micro-economy level is highly needed.

Tweet: Care for the critically ill could improve ICUs in low- and middle-income countries

ABSTRACT

Sepsis is a major reason for intensive care unit (ICU) admission, also in resource-poor settings. ICUs in low- and middle-income countries (LMICs) face many challenges that could affect patient outcome. The aim of this review is to describe differences between resource-poor and resource-rich settings regarding the epidemiology, pathophysiology, economics and research aspects of sepsis. We restricted this manuscript to the ICU setting even knowing that many sepsis patients in LMICs are treated outside an ICU. Although many bacterial pathogens causing sepsis in LMICs are similar to those in high-income countries, resistance patterns to antimicrobial drugs can be very different; in addition, causes of sepsis in LMICs often include tropical diseases in which direct damaging effects of pathogens and their products can sometimes be more important than the response of the host. There are substantial and persisting differences in ICU capacities around the world; not surprisingly the lowest capacities are found in LMICs, but with important heterogeneity within individual LMICs. Although many aspects of sepsis management developed in rich countries are applicable in LMICs, implementation requires strong consideration of cost implications and the important differences in resources. Addressing both disease-specific and setting-specific factors is important to improve performance of ICUs in LMICs. Although critical care for severe sepsis is likely cost-effective in LMIC setting, more detailed evaluation at both at a macro- and micro-economy level is necessary. Sepsis management in resource-limited settings is a largely unexplored frontier with important opportunities for research, training, and other initiatives for improvement.

INTRODUCTION

In many low- and middle-income countries (LMICs), with improved public health services like sanitation and immunization, the relative importance of curative care to improve health becomes more important. This includes care for severe sepsis, a major reason for intensive care unit (ICU) admission in LMIC. These patients will currently often be treated in general wards, but basic intensive care facilities are becoming increasingly available. The scope of the current review is limited to the ICU setting. ICUs in resource-restricted settings have to function with important limitations, including both infrastructure and materials, as well as human resources. It is important to address economic aspects around the provision of relatively expensive intensive care in low-income countries. Most LMIC have tropical or subtropical climates, and causes of sepsis will often be different from high-income countries (HIC), where most sepsis treatment guidelines have been developed. Because of the different setting and the different causes of sepsis, existing guidelines will need prudent interpretation. There is a broad research agenda around this, which is currently still hardly addressed. Finally, expansion of setting-adapted training will be important to improve ICU performance in LMIC.

In this review, written by a group of physicians from resource-poor and -rich ICUs in LMICs and high-income countries, respectively, who were involved in the development of a series of sets of recommendations for sepsis management in resource-poor settings as recently published in this journal [1-4], the estimated burden, pathogens and pathophysiology of sepsis are compared between resource-poor and resource-rich settings. The availability of critical care and guidelines, and costs of critical care in LMICs are compared to that in high-income countries. Suggestions for future directions are provided.

BURDEN AND CAUSES OF SEPSIS AND ITS MANAGEMENT

Disease burden

While detailed information has been reported on the epidemiology and outcome of sepsis in high-income countries [5, 6], only scant systematically collected epidemiological data exist from LMICs [7, 8], despite that these countries carry about 80% of the global mortality caused by infections [9]. At present, the epidemiology of sepsis in LMICs can only be loosely estimated from the epidemiology of acute infectious diseases with a potential to cause sepsis captured in the 'Global Burden of Disease' database [10-12]. This database reported important regional differences in the epidemiology of such acute infectious diseases (Figure 1). For instance, while the majority of acute infections in resource-limited settings are acquired in the community [9], the incidence of nosocomial infections such as catheter-related bloodstream infections or ventilator-associated pneumonia are several fold higher in LMICs than in high-income countries like the United States [13]. Similarly, only few data on sepsis mortality in LMICs have so far been published. These suggest that sepsis-related mortality greatly varies among regions and countries according to their income level. Sepsis attributable case fatality in high-income countries has been decreasing over the last decades to 30 to 40% [5, 6], whereas case fatality rates of up to 80% continue to be reported from resource-poor regions of the world [14-17].

Causative pathogens and pathogenesis

Although many bacterial pathogens causing sepsis in LMICs are similar to those in high-income countries, resistance patterns to antimicrobial drugs can be very different. It has been suggested that the high prevalence of multi-drug resistant bacteria, including methicillin-resistant *Staphylococcus aureus*, extended spectrum

beta-lactamases producing bacteria, carbapenamase producing enterobacteriaceae and mycobacterium tuberculosis, contribute to the excess deaths observed in LMICs caused by invasive infections with these bacteria, particularly among infants [13, 18]. Five countries with the highest burden of under 5 deaths (China, Nigeria Pakistan, India and the DRC) have the highest neonatal deaths from antimicrobial resistance [19]. The problems with antimicrobial resistance and its implications for the treatment of sepsis in LMIC have been described in an earlier publication in this journal [4].

Whereas the majority of severe sepsis in high-income countries is caused by bacterial infections, in LMICs, many of which are located in the tropics, causes of severe sepsis also include acute non-bacterial diseases, including protozoal diseases such as malaria, and viral diseases such as measles, dengue, or viral hemorrhagic fevers. The international literature mainly focuses on sepsis caused by invasive bacterial infections and the associated systemic inflammatory response [20, 21]. Therefore, non-bacterial causes are understudied and the acquired knowledge on the pathophysiology and treatment of sepsis may not be generalizable to these other causes of sepsis [22].

Previous sepsis definitions put a large emphasis on the 'dysregulated response of the host's immune system' as the key element of the pathogenesis of sepsis [23, 24]. Although this will be generally correct, this ignores potential direct damaging effects of certain pathogens or pathogen products, which can in particular play a role in tropical diseases. For instance, in severe falciparum malaria, a blocked microcirculation resulting from the sequestered infected red blood cell biomass is a direct cause of vital organ failure [25]; and in dengue shock syndrome virus proteins are thought to damage the glycocalyx lining the endothelium directly [26, 27].

Intervening in these pathophysiological pathways will obviously require therapeutic approaches different from those in bacterial sepsis.

Poor availability of critical care

There is persisting substantial heterogeneity in ICU capacity around the world. In high-income countries in Europe and Northern America, capacity is between 5 and 30 beds per 100,000 inhabitants. The scarce data available show capacity is much lower, albeit quite variable in LMICs [28]. For instance, in Asia reported ICU capacity is only 0.3 beds per 100,000 inhabitants in Bangladesh, 2.4 per 100,000 inhabitants in Malaysia; 2.5 per 100,000 inhabitants in Sri Lanka, and 3.9/100,000 inhabitants in China [29], but in contrast 11.7 adult and pediatric ICU beds per 100,000 inhabitants in Mongolia [30]. Studies from several countries in Sub-Saharan Africa reported as few as 0.1 to 0.2 ICU beds per 100,000 inhabitants [31, 32]. With the exception of Mongolia [30] and Latin-American countries [33], almost no data on the availability of dedicated pediatric ICU capacities have been published for resource-limited settings [34].

We tend to look at LMICs as if they are ‘uniform’, but this is a too simplistic if not naive approach. Within and between LMIC, there is important heterogeneity in availability of intensive care, resourcing of ICUs, quality of services and case-mix [35]. The rapidly expanding urban population in many LMIC will provide a challenge for current urban ICU capacity because of the associated increase in case load [36].

Another problem in reporting ICU capacity is the lack of a commonly agreed definition of an ICU or an ICU bed [37, 38]. The spectrum of how ICUs are staffed and equipped differs vastly between countries and regions. Table 1 summarizes published evidence and personal experience of the authors in an attempt to categorize different ICU structures worldwide. As surveys from various countries

suggest, the availability of ICU-related material resources directly correlates with the countries' income level and health care spending [7, 31, 32, 39-43]. The shortage of medical professionals specifically trained in the care of acutely and critically ill patients is another widespread and serious challenge for ICU services in many LMICs [31, 32, 39-42]. A notable exception to this is well-staffed and equipped ICUs in private health care facilities. Such services can typically be found in some high- and middle-income countries, but these usually remain only accessible for those who can afford this.

The increasing but still low ICU capacity in poorer regions [44] implies that access to ICU services for critically ill patients is usually severely limited. This results in frequent triage decisions [32], which likely increases preventable disease mortality [45]. Even though costs of care in ICUs of resource-limited settings are only a fraction of those in ICUs of high-income countries [46], expenses for ICU care are usually to a large extent covered by the patient's family and relatives in LMICs. Unwanted consequences can be denial or refusal of ICU admission of poor patients, but also the premature withdrawal of life-saving interventions [47, 48]. In other instances, costs of care for a critically ill patient, who may eventually die, can exceed the limited budget of many families leaving them with high debts or even causing private bankruptcy.

Incomplete and unadjusted guidelines

The principle of 'evidence-based medicine' is equally important in resource-limited as in resource-rich settings. Development of local evidence is important, since case-mix and causes of sepsis, but also available infrastructure and facilities are essentially different from that in Western countries. Resource-limited ICUs are frequently limited in availability of equipment, laboratory support and skilled physician

and nursing staff. As a result, recommendations on sepsis management in resource poor settings, such as those developed by the '*Global Intensive Care Working Group*' of the European Society of Intensive Care Medicine (ESICM) [49] differ in several aspects from the Surviving Sepsis Campaign–guidelines, which were developed in high–income settings [50]. Obvious examples include targeting strict blood glucose levels with insulin, which can be safe with frequent and reliable blood glucose monitoring, while it is a dangerous strategy when the effects of insulin titrations are determined infrequently or not at all. Other modifications in recommendations for sepsis management could result from the differences in availability of and indications for fluids in sepsis patients between resource–rich and resource–poor settings, but also the cause of sepsis (see also textbox 'Availability of and indications for fluids in sepsis patients differ between HICs and LMICs'). Recently, our group of physicians from resource–poor and –rich countries took the initiative to develop sets of recommendations for management of patients with sepsis in resource–poor ICUs [1-4].

Despite the importance of building 'local' evidence, many aspects of existing guidelines and basic principles of good critical care are universal. Access to adequate information has improved massively in this cyber age, and several courses (such as the '*BASIC for Developing Healthcare Systems*' courses developed by the Chinese University of Hong Kong, Hong Kong, China, and *Médecins Sans Frontières*) are available. Theoretical knowledge of specialist doctors in developing countries is often impressive, but translating this into practical implementation can lag behind, which is an important scope for training. Training should not only be for ICU physicians, also for nurses, and other clinical personnel. It will be important that education on sepsis management does not focus only on ICU physicians, but include

medical schools, nursing schools and training of other health workers, as many sepsis patients in LMICs receive treatment outside an ICU. 'Train-the-trainer' models and fostering local champions for positive change are important to sustain improvements of care.

COSTS OF CARE IN SEPSIS

Expensive but likely cost-effective critical care

The vast resources consumed by ICUs, up to a staggering 1% of total gross domestic product in the United States [51], demands their subjection to explicit economic cost-benefit considerations [52]. The literature from high-income settings has indeed expanded over the past decade, demonstrating that many critical care interventions offer significant health returns with costs per Quality-adjusted Life Year (QALY) gained well below a threshold of US \$50,000, indicative of a cost-effective intervention in this context.

For countries in which more basic services are still relatively undeveloped it might be questioned whether investment in ICUs should take priority over the strengthening of lower tiers of the healthcare system [53]. Irrespective of these considerations, the reality is that in many growing economies ICUs are increasing in number [44], with virtually no evidence regarding the cost-effectiveness of their services.

Costs of critical care amongst regions

One approach is to consider the evidence from high-income countries, with a much lower willingness to pay the abovementioned threshold. Applying a threshold of US \$4,000 per QALY gained [54] suggests that numerous ICU interventions and those for sepsis in particular [55-61], are likely to be cost-effective in ICUs in LMICs (Table 2). For several reasons these cost-effectiveness ratios might be conservative in the context of LMIC. First, the cost of ICUs in these countries is much lower than in high-income countries. The two largest standardized multi-country reviews of critical care costs in high-income settings estimated the cost per ICU-day ranging from US \$850 to US \$3,400 in 2015 [62, 63], with labor being the dominant cost-driver (61 and

67%, respectively). The proportional difference in labor costs between high-income countries and LMICs will be similar to or higher than that of their respective gross domestic product per capita [64]. This alone implies that the critical care costs in LMICs will be less than a third that in high-income countries. Other costs, such as laboratory services (10% of total critical care cost [63]), also are likely to be far lower for similar reasons, as well as drug costs, and cost-saving practices such as recycling of consumables [65]. In one of the few costing analyses of an ICU from a LMIC, the total cost per ICU admission day in India was estimated at just over US \$200 [66], between approximately 5 to 20% that for high-income countries.

However, also the absolute financial outlay per intervention multiplied by the number of interventions is important to consider, in particular in LMICs. Even in a wealthy country with a high threshold, its tolerance depends exquisitely on the number of times it must spend the money, especially if up-front costs are high. In other words, an intervention might appear cost-effective, but if it would have to be applied to a massive portion of the public, the costs would still be perceived as financially impossible. This also requires there is an effective system of triage in place to see that these interventions are offered to patients who are most likely to benefit, at times a challenge in LMICs.

Finally, we are uncertain whether applying a threshold of US \$4,000 per QALY gained is one that could be promoted or generalized, as there is a large intra- and inter-region variation in resources. Country, and ideally region-specific currency conversions/purchase power parity comparisons should be made.

Benefit of critical care amongst regions

The mean age of adult patients admitted to ICUs in high-income countries is consistently higher than those reported in LMIC. For example, in European ICUs the

mean age of adult admissions is typically 55 to 66 years, much higher than the median age of 34 in a Rwandan ICU [67], the mean age of 32 in one of the few ICUs in Uganda [1], and the median age of 37, 51 and 49 in ICUs in Bangladesh, Nepal and India [68]. Although not specified for admission diagnosis, it likely also reflects the younger age of sepsis patients. This can be taken as an argument to invest in better intensive care to save these young lives from an in principle treatable disease. Also, in LMICs where social security networks are usually lacking, the loss of an individual's economic activity has far more extensive economic consequences for the families involved.

By generalizing the evidence from high-income countries and considering the lower costs and higher potential gains there are strong indications that a wide range of critical care services are likely to be cost-effective in LMICs. Unfortunately, there is a dearth of evidence directly from these settings to confirm this. The only economic evaluation of an ICU in Bosnia-Herzegovina, concluded that critical care was highly cost-effective in their setting [46]. A study in Brazil showed that the implementation of a sepsis management protocol was associated with an absolute reduction of 18% in mortality and with cost-savings [55].

Impact of certain interventions amongst regions

Despite these suggestions that a broad range of ICU services could be cost-effective in LMICs there are also reason for caution. First, some interventions are likely to have a much lower impact in poorly functioning environments than they might have in resource-rich settings. For example, invasive monitoring in an ICU with poor basic infection control could result in more harm than benefit. For this reason it is imperative that basic standards of care are in place prior to introduction of costly interventions whose effect might otherwise be compromised. Training programs to

improve the general quality of care that require no costly interventions have been shown to have beneficial effects [68]. Second, while hypothetically cost–effectiveness should correspond with affordability, in LMICs where health systems are often fragmented and divided between the private and public sector and an abundance of vertically funded health programs, this is often not the case. It is imperative therefore that only interventions with modest budgetary impacts are shortlisted for consideration and from those that have the greatest beneficial effects, with the lowest incremental costs are selected for implementation. Some such low–cost approaches have been identified and should be prioritized for evaluation. Better surveillance systems for local etiologies of sepsis and antimicrobial susceptibility patterns has been recognized as a key requirement for improving the management of sepsis and critical care [4], and a modeling based economic evaluation supported the notion that this is likely to be highly cost–effective [69].

Likewise, the development and adaptation of LMIC–specific risk prediction models have been shown to potentially outperform models widely used in high–income settings like the frequently used Acute Physiology and Chronic Health Evaluation score, and the Simplified Acute Physiology Score [67], which might not even be feasible in many ICUs in LMICs. The same is true for scoring systems in children and scoring systems that are context relevant should also be explored because the PRISM may under predict mortality [70].

Critical care in general and a subset of specific interventions can be an efficient use of scarce resources in LMICs. But despite the aforementioned lower costs, an ICU admission in e.g., India would most likely represent a catastrophic expenditure consuming over half the mean annual household income [44, 66] (see also: textbox 2 – Critical Care Expenditure in Indian Hospitals). As such facilities

expand in LMICs and inevitably consume increasing resources, it is imperative for health authorities to ensure these offer affordable and cost-effective services, while monitoring and continuing to seek opportunities to improve the quality of critical care and their efficiency at low cost. This was exemplified by the United Kingdom in the late 1990s, where a country-wide initiative to transform and modernize critical care was associated with a 10% reduction in mortality at lower cost increases than would otherwise be expected [71].

SEPSIS RESEARCH IN RESOURCE-LIMITED SETTINGS

A yet largely untouched research agenda

With only 1.7% of all biomedical research publications originating in low-income countries [72], and a likely even greater disparity in critical care [44], both the needs and the opportunities for critical care research are vast. Recent rigorous attempts to quantify the global burdens of sepsis, infection and respiratory failure confirm a profound lack of epidemiologic critical illness data from low-income countries, with the only reliable data limited to single-center descriptions [67, 73-76]. The ‘*International Severe Acute Respiratory and Emerging Infection Consortium*’ (ISARIC), the ‘*Global Intensive Care Working Group*’ of the ESICM, and the Mahidol Oxford Research Unit in Bangkok, Thailand, are three examples of groups working to create the infrastructure for global epidemiologic data on critical illness, for use both as baseline data at regular intervals but also in preparation for disease outbreaks [77].

Research on infectious diseases that lead to critical illness are perhaps the most successful areas of investigation thus far, with impressive scientific advancements in diseases such as malaria, tuberculosis, and melioidosis [78-80]. Defining and testing quality metrics is an area prioritized in both critical care and global public health [81], but quality improvement in resource-limited ICUs remains largely unexplored [44, 82]. The latter requires some ability to benchmark intensive care units with severity of illness scores, also a nascent area of study [67]. Research into medical education models like the ‘*Human Resources for Health*’ program in Rwanda [83], decision-support tools like the ‘*Checklist for Early Recognition and Treatment of Acute Illness*’ (CERTAIN) [84] and ‘*Vital Signs Directed Therapy*’ [85]

could help in better defining how to optimize knowledge acquisition and application for providers in resource-limited settings.

Challenges with research in resource-restricted settings

The realities of restriction in resources interact to create complex challenges to producing quality critical care research. Many short-term multi-site epidemiologic studies depend on individual sites to participate without funding for the perceived minimal data collection burden. However, staff at resource-constrained sites often cannot spare even the short-term investment required to participate and 'standard' clinical data such as arterial blood gas analyses and chest radiographs are often not available in these sites [75, 77]. Likewise, multi-center clinical trials are significantly more expensive to perform in resource-limited settings since baseline clinical or analytic infrastructure of any sort cannot be assumed. Efforts to build resource capacity in research and to ensure fair authorship opportunities also mean that research has two resource-intensive and sometimes competing goals: that of developing local researchers without prior training and that of producing high-quality research. An unintended consequence of (appropriate) increasing involvement by local institutional review boards is that lack of staffing and experience may lead to unnecessary delays in research approvals. These resource barriers to collecting epidemiologic data are exacerbated by the fact that 'critical illness' is not a laboratory-defined condition, but rather one that is often defined by the expensive resources used to treat it [28, 86].

Ethical considerations are complex as well. All participants in resource-limited settings must be considered potentially 'vulnerable' populations due to extreme need and lack of health or research literacy. Deciding where equipoise exists for interventions proven to work in high-resource settings in settings of low resources is

difficult. Asking the question, 'how do we do this better given limited resources' uncomfortably raises the question 'how do we increase resources and to what extent are we obligated to do so?' Ethical considerations also include relationships between local and foreign researchers, the latter who often bring financial and experiential resources leading to unequal power dynamics and potential for abuse [87]. Allowing lower standards for publication for research originating in low-resource settings has been considered in order to decrease some of the publication bias toward resource-rich settings; however, it is not at all clear that publishing poor-quality research that could impact clinical care is ethical or advisable.

SUGGESTIONS FOR THE FUTURE

Better definition of sepsis

The latest sepsis definition [88], which refers to sepsis as life-threatening organ dysfunction caused by a dysregulated host response to infection, better reflects the fact that sepsis can complicate any serious acute infection. As pointed out earlier, direct damaging effects of the pathogen itself, or its products can sometimes be the main process. Consequently, research on sepsis treatment should not focus only on immunomodulating strategies in bacterial sepsis, but also on faster illumination of the pathogen and its products, which could be in particular relevant in certain non-bacterial causes of sepsis like falciparum malaria, cryptococcal and various viral infections.

Another unmet need is to validate the various definitions for sepsis in more varied populations, like children in LMICs, adults in LMICs, and areas where the causative pathogens of sepsis differ from Western countries where the present definitions have been developed.

Better research infrastructures and planning

Experience, resources, and human power are critical for building evidence, which are not routinely available in resource-poor settings. The global critical care community should help building local clinical research capacities and contribute to obtaining adequate funding. Engagement of a variety of stakeholders, including local intensive care societies, healthcare authorities (e.g., ministries of health) and universities will be crucial in this respect [89]. There are large funding bodies, such as the Wellcome Trust that support several efforts, but additional funds are clearly needed. Teaming up of researchers from established groups in high-income countries with local groups in resource-limited settings has proven a good model, provided equal and

reciprocal relationships are guaranteed. Formulating essential topics for research will be important and could benefit from increased networking between critical care physicians from developing countries. This would also foster research networks needed to perform adequately powered clinical trials. Requirements for quality clinical research are not necessarily available in many developing countries, with often little research infrastructure or human resources for research available. Hospitals and their doctors can be overburdened by their service delivery tasks, leaving little room for research. Offering a career path to clinical research physicians could free up manpower for this important aspect of improving evidence based critical care. Priority settings and governance regarding financial resources for research and implementation projects will become extremely important. Also, local institutions, including ethical review boards, should help creating an enabling environment for research benefiting the local population, and not promote instituting unnecessary barriers, which is now at times the case.

Obtaining relevant evidence, adapting guidelines

Obtaining local intelligence on the most important causes of sepsis and the resistance patterns of the infecting microorganisms is crucial to guide local empirical antimicrobial treatment. Since microbiological capacity is often lacking in hospitals in LMICs, research collaborations could help obtaining this evidence from strategically located sentinel study sites.

We need additional recommendations for those interventions not yet covered in the published guidelines. Multi-center trials assessing clinical efficacy and effectiveness of new interventions and interventions known to be effective in resource-rich settings are sorely needed, given differences in epidemiologic and treatment contexts [44]. The Fluid Expansion As Supportive Therapy (FEAST) trial on

fluid resuscitation in febrile African children with shock demonstrates that these trials are feasible and may yield unexpected results [90]. Networks of critical care researchers are increasingly established in more resource-limited areas of the world, which could lead to additional studies of this kind [44]. Beyond these broad areas of research, specific subtypes deserve special attention in resource-limited settings. Social determinants of disease and barriers to care are of particular relevance to all research in resource-limited settings including intensive care [81]. While cost-effectiveness analyses are arguably important in all environments without infinite resources, their need is more acute in places with fewer resources to allocate [91]. Locally adapted prediction models can help ensure that interventions are targeted in a cost-effective manner.

Another important area for critical care research is on developing templates for expanding urban ICU capacity in fast growing cities [36]. Setting-specific guidance, instead of country-specific guidance how to build and equip an ICU probably is an additional and substantial area of research.

Finally, research on the process whereby scientific knowledge is translated into improved quality of care, ‘global health delivery science’, gets at the immediate challenge in patient care in resource-poor settings, the fact that so much of what is known is not implemented effectively [81].

Opportunities

While the challenges are daunting, the opportunities are similarly impressive.

Outbreaks like Ebola have increased the drive for intensive care research in resource-limited settings both by demonstrating how critical illness in these settings impacts people in resource-rich settings, and by highlighting the need for improved critical care capacity in all areas of the world [92]. Interest in and funding for global

health has increased steadily over the last few decades, such that career path and funding opportunities for high-income country researchers are better than ever before. Researchers from resource-limited settings now also have opportunities for high-quality training and mentorship through programs, for example through *Wellcome Trust* or United Kingdom '*Medical Research Council*' schemes and the American Thoracic Society's '*Methods in Epidemiologic, Clinical, and Operations Research*' (MECOR) program [93].

An example of building a research structure is one from India, where the Indian Society of Critical Care Medicine initiated a 'cloud based database' called 'Customized, Health in Intensive care, Trainable Research & Analysis (CHITRA) (<http://www.isccm.org/chitra.aspx>).

Finally, LMIC and high-income countries could mutually benefit from their research agendas; HIC also gain from more LMIC research. For example, LMIC considerations about direct pathogen effects could, in turn, generate better critical thinking about sepsis, from which HIC and LMIC gain. Similarly, technologic innovation under resource constraints can yield gains for all. Critical care could be ripe for this at times called 'frugal innovation' or 'reverse innovation' concept [94]. Also, in our personal experience [68], nurses from HIC that have worked in LMIC, and even for a short time, are more cost-conscious than their colleagues.

CONCLUSIONS

Strategies to improve the quality of sepsis management in resource-poor settings require consideration of disease-specific and setting-specific factors and meticulous evaluation of the best way to adapt and deploy quality improvement initiatives.

Critical care, including sepsis management, is expensive but likely cost-effective in LMICs, but we need to better understand what the true financial impact of critical care is, both at a macro- and micro-economy level. Sepsis management in resource-limited settings is a largely unexplored frontier with a clear mandate and exciting opportunities for impact.

REFERENCES

1. Kwizera A, Festic E, Dunser MW, (2016) What's new in sepsis recognition in resource-limited settings? *Intensive Care Med*
2. Musa N, Murthy S, Kissoon N, (2016) Pediatric sepsis and septic shock management in resource-limited settings. *Intensive Care Med*
3. Serpa Neto A, Schultz MJ, Festic E, (2016) Ventilatory support of patients with sepsis or septic shock in resource-limited settings. *Intensive Care Med* 42: 100-103
4. Thwaites CL, Lundeg G, Dondorp AM, sepsis in resource-limited settings-expert consensus recommendations group of the European Society of Intensive Care M, the Mahidol-Oxford Research Unit in Bangkok T, (2016) Recommendations for infection management in patients with sepsis and septic shock in resource-limited settings. *Intensive Care Med*
5. Martin GS, Mannino DM, Eaton S, Moss M, (2003) The epidemiology of sepsis in the United States from 1979 through 2000. *N Engl J Med* 348: 1546-1554
6. Vincent JL, Sakr Y, Sprung CL, Ranieri VM, Reinhart K, Gerlach H, Moreno R, Carlet J, Le Gall JR, Payen D, Sepsis Occurrence in Acutely Ill Patients I, (2006) Sepsis in European intensive care units: results of the SOAP study. *Crit Care Med* 34: 344-353
7. Phua J, Koh Y, Du B, Tang YQ, Divatia JV, Tan CC, Gomersall CD, Faruq MO, Shrestha BR, Gia Binh N, Arabi YM, Salahuddin N, Wahyuprajitno B, Tu ML, Wahab AY, Hameed AA, Nishimura M, Procyshyn M, Chan YH, Group MS, (2011) Management of severe sepsis in patients admitted to Asian intensive care units: prospective cohort study. *BMJ* 342: d3245

8. Murthy S, Leligdowicz A, Adhikari NK, (2015) Intensive care unit capacity in low-income countries: a systematic review. *PLoS One* 10: e0116949
9. Cheng AC, West TE, Limmathurotsakul D, Peacock SJ, (2008) Strategies to reduce mortality from bacterial sepsis in adults in developing countries. *PLoS Med* 5: e175
10. Collaborators GBoD MaCoD, (2015) Global, regional, and national age-sex specific all-cause and cause-specific mortality for 240 causes of death, 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 385: 117-171
11. Global Burden of Disease Study C, (2015) Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 386: 743-800
12. Global Burden of Disease Study M, Causes of Death C, (2016) Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980-2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 388: 1459-1544
13. Rosenthal VD, Bijie H, Maki DG, Mehta Y, Apisarnthanarak A, Medeiros EA, Leblebicioglu H, Fisher D, Alvarez-Moreno C, Khader IA, Del Rocio Gonzalez Martinez M, Cuellar LE, Navoa-Ng JA, Abouqal R, Guanche Garcell H, Mitrev Z, Pirez Garcia MC, Hamdi A, Duenas L, Cancel E, Gurskis V, Rasslan O, Ahmed A, Kanj SS, Ugalde OC, Mapp T, Raka L, Yuet Meng C, Thu le TA, Ghazal S, Gikas A, Narvaez LP, Mejia N, Hadjieva N, Gamar Elanbya MO, Guzman Siritt ME, Jayatilleke K, members I, (2012) International Nosocomial

- Infection Control Consortium (INICC) report, data summary of 36 countries, for 2004-2009. *Am J Infect Control* 40: 396-407
14. Frikha N, Mebazaa M, Mnif L, El Euch N, Abassi M, Ben Ammar MS, (2005) [Septic shock in a Tunisian intensive care unit: mortality and predictive factors. 100 cases]. *Tunis Med* 83: 320-325
 15. Jacob ST, Moore CC, Banura P, Pinkerton R, Meya D, Opendi P, Reynolds SJ, Kenya-Mugisha N, Mayanja-Kizza H, Scheld WM, Promoting Resource-Limited Interventions for Sepsis Management in Uganda Study G, (2009) Severe sepsis in two Ugandan hospitals: a prospective observational study of management and outcomes in a predominantly HIV-1 infected population. *PLoS One* 4: e7782
 16. Siddiqui S, (2007) Not "surviving sepsis" in the developing countries. *J Indian Med Assoc* 105: 221
 17. Dunser MW, Bataar O, Tsenddorj G, Lundeg G, Torgersen C, Romand JA, Hasibeder WR, Helfen Beruhrt Study T, (2008) Differences in critical care practice between an industrialized and a developing country. *Wien Klin Wochenschr* 120: 600-607
 18. Vernet G, Mary C, Altmann DM, Doumbo O, Morpeth S, Bhutta ZA, Klugman KP, (2014) Surveillance for antimicrobial drug resistance in under-resourced countries. *Emerg Infect Dis* 20: 434-441
 19. Laxminarayan R, Matsoso P, Pant S, Brower C, Rottingen JA, Klugman K, Davies S, (2016) Access to effective antimicrobials: a worldwide challenge. *Lancet* 387: 168-175
 20. Bone RC, Balk RA, Cerra FB, Dellinger RP, Fein AM, Knaus WA, Schein RM, Sibbald WJ, (1992) Definitions for sepsis and organ failure and guidelines for

- the use of innovative therapies in sepsis. The ACCP/SCCM Consensus Conference Committee. American College of Chest Physicians/Society of Critical Care Medicine. Chest 101: 1644-1655
21. Levy MM, Fink MP, Marshall JC, Abraham E, Angus D, Cook D, Cohen J, Opal SM, Vincent JL, Ramsay G, International Sepsis Definitions C, (2003) 2001 SCCM/ESICM/ACCP/ATS/SIS International Sepsis Definitions Conference. Intensive Care Med 29: 530-538
 22. Kissoon N, Daniels R, van der Poll T, Finfer S, Reinhart K, (2016) Sepsis-The Final Common Pathway to Death From Multiple Organ Failure in Infection. Crit Care Med 44: e446
 23. Auma MA, Siedner MJ, Nyehangane D, Nalusaji A, Nakaye M, Mwanga-Amumpaire J, Muhindo R, Wilson LA, Boum Y, 2nd, Moore CC, (2013) Malaria is an uncommon cause of adult sepsis in south-western Uganda. Malar J 12: 146
 24. White NJ, Pukrittayakamee S, Hien TT, Faiz MA, Mokuolu OA, Dondorp AM, (2014) Malaria. Lancet 383: 723-735
 25. White NJ, Turner GD, Day NP, Dondorp AM, (2013) Lethal malaria: Marchiafava and Bignami were right. J Infect Dis 208: 192-198
 26. Chen Y, Maguire T, Hileman RE, Fromm JR, Esko JD, Linhardt RJ, Marks RM, (1997) Dengue virus infectivity depends on envelope protein binding to target cell heparan sulfate. Nat Med 3: 866-871
 27. Avirutnan P, Zhang L, Punyadee N, Manuyakorn A, Puttikhunt C, Kasinrerk W, Malasit P, Atkinson JP, Diamond MS, (2007) Secreted NS1 of dengue virus attaches to the surface of cells via interactions with heparan sulfate and chondroitin sulfate E. PLoS Pathog 3: e183

28. Adhikari NK, Fowler RA, Bhagwanjee S, Rubenfeld GD, (2010) Critical care and the global burden of critical illness in adults. *Lancet* 376: 1339-1346
29. Haniffa R, De Silva AP, Iddagoda S, Batawalage H, De Silva ST, Mahipala PG, Dondorp A, de Keizer N, Jayasinghe S, (2014) A cross-sectional survey of critical care services in Sri Lanka: a lower middle-income country. *J Crit Care* 29: 764-768
30. Mendsaikhan N, Begzjav T, Lundeg G, Brunauer A, Dunser MW, (2016) A Nationwide Census of ICU Capacity and Admissions in Mongolia. *PLoS One* 11: e0160921
31. Jochberger S, Ismailova F, Lederer W, Mayr VD, Luckner G, Wenzel V, Ulmer H, Hasibeder WR, Dunser MW, Helfen Beruhrt" Study T, (2008) Anesthesia and its allied disciplines in the developing world: a nationwide survey of the Republic of Zambia. *Anesth Analg* 106: 942-948, table of contents
32. Kwizera A, Dunser M, Nakibuuka J, (2012) National intensive care unit bed capacity and ICU patient characteristics in a low income country. *BMC Res Notes* 5: 475
33. Campos-Mino S, Sasbon JS, von Dessauer B, (2012) [Pediatric intensive care in Latin America]. *Med Intensiva* 36: 3-10
34. Kissoon N, (2016) Sepsis Care Differences Unlike Beauty Are Not Skin Deep. *Pediatr Crit Care Med* 17: 568-569
35. Divatia JV, Amin PR, Ramakrishnan N, Kapadia FN, Todi S, Sahu S, Govil D, Chawla R, Kulkarni AP, Samavedam S, Jani CK, Rungta N, Samaddar DP, Mehta S, Venkataraman R, Hegde A, Bande BD, Dhanuka S, Singh V, Tewari R, Zirpe K, Sathe P, Investigators IS, (2016) Intensive Care in India: The

- Indian Intensive Care Case Mix and Practice Patterns Study. *Indian J Crit Care Med* 20: 216-225
36. Austin S, Murthy S, Wunsch H, Adhikari NK, Karir V, Rowan K, Jacob ST, Salluh J, Bozza FA, Du B, An Y, Lee B, Wu F, Nguyen YL, Oppong C, Venkataraman R, Velayutham V, Duenas C, Angus DC, International Forum of Acute Care T, (2014) Access to urban acute care services in high- vs. middle-income countries: an analysis of seven cities. *Intensive Care Med* 40: 342-352
 37. Marshall JC, Bosco L, Adhikari NK, Connolly B, Diaz JV, Dorman T, Fowler RA, Meyfroidt G, Nakagawa S, Pelosi P, Vincent JL, Vollman K, Zimmerman J, (2016) What is an intensive care unit? A report of the task force of the World Federation of Societies of Intensive and Critical Care Medicine. *J Crit Care in press*
 38. Rhodes A, Ferdinande P, Flaatten H, Guidet B, Metnitz PG, Moreno RP, (2012) The variability of critical care bed numbers in Europe. *Intensive Care Med* 38: 1647-1653
 39. Baelani I, Jochberger S, Laimer T, Otieno D, Kabutu J, Wilson I, Baker T, Dunser MW, (2011) Availability of critical care resources to treat patients with severe sepsis or septic shock in Africa: a self-reported, continent-wide survey of anaesthesia providers. *Crit Care* 15: R10
 40. Baker T, Lugazia E, Eriksen J, Mwafongo V, Irestedt L, Konrad D, (2013) Emergency and critical care services in Tanzania: a survey of ten hospitals. *BMC Health Serv Res* 13: 140
 41. Bataar O, Lundeg G, Tsenddorj G, Jochberger S, Grander W, Baelani I, Wilson I, Baker T, Dunser MW, Helfen Beruhrt Study T, (2010) Nationwide

- survey on resource availability for implementing current sepsis guidelines in Mongolia. *Bull World Health Organ* 88: 839-846
42. Hsia RY, Mbembati NA, Macfarlane S, Kruk ME, (2012) Access to emergency and surgical care in sub-Saharan Africa: the infrastructure gap. *Health Policy Plan* 27: 234-244
 43. Arabi YM, Phua J, Koh Y, Du B, Faruq MO, Nishimura M, Fang WF, Gomersall C, Al Rahma HN, Tamim H, Al-Dorzi HM, Al-Hameed FM, Adhikari NK, Sadat M, Asian Critical Care Clinical Trials G, (2016) Structure, Organization, and Delivery of Critical Care in Asian ICUs. *Crit Care Med* 44: e940-948
 44. Dondorp AM, Iyer SS, Schultz MJ, (2016) Critical Care in Resource-Restricted Settings. *JAMA* 315: 753-754
 45. Mendsaikhan N, Begzjav T, Lundeg G, Dünser MW, (2016) Potentially preventable deaths by intensive care medicine in Mongolian hospitals. *Crit Care Research Practice*: in press
 46. Cubro H, Somun-Kapetanovic R, Thiery G, Talmor D, Gajic O, (2016) Cost effectiveness of intensive care in a low resource setting: A prospective cohort of medical critically ill patients. *World J Crit Care Med* 5: 150-164
 47. Phua J, Joynt GM, Nishimura M, Deng Y, Myatra SN, Chan YH, Binh NG, Tan CC, Faruq MO, Arabi YM, Wahjuprajitno B, Liu SF, Hashemian SM, Kashif W, Staworn D, Palo JE, Koh Y, Investigators AS, Asian Critical Care Clinical Trials G, (2016) Withholding and withdrawal of life-sustaining treatments in low-middle-income versus high-income Asian countries and regions. *Intensive Care Med* 42: 1118-1127

48. Kissoon N, (2016) Healthcare Costs to Poor Families: An Agonising Burden. *Indian J Pediatr* 83: 1063-1064
49. Dunser MW, Festic E, Dondorp A, Kissoon N, Ganbat T, Kwizera A, Haniffa R, Baker T, Schultz MJ, Global Intensive Care Working Group of European Society of Intensive Care M, (2012) Recommendations for sepsis management in resource-limited settings. *Intensive Care Med* 38: 557-574
50. Dellinger RP, Levy MM, Rhodes A, Annane D, Gerlach H, Opal SM, Sevransky JE, Sprung CL, Douglas IS, Jaeschke R, Osborn TM, Nunnally ME, Townsend SR, Reinhart K, Kleinpell RM, Angus DC, Deutschman CS, Machado FR, Rubenfeld GD, Webb SA, Beale RJ, Vincent JL, Moreno R, Surviving Sepsis Campaign Guidelines Committee including the Pediatric S, (2013) Surviving sepsis campaign: international guidelines for management of severe sepsis and septic shock: 2012. *Critical care medicine* 41: 580-637
51. Halpern NA, Pastores SM, Greenstein RJ, (2004) Critical care medicine in the United States 1985-2000: an analysis of bed numbers, use, and costs. *Crit Care Med* 32: 1254-1259
52. Pinsky MR, (2002) Understanding costs and cost-effectiveness in critical care: report from the second American Thoracic Society workshop on outcomes research. *Am J Respir Crit Care Med* 165: 540-550
53. Shann F, (2011) Role of intensive care in countries with a high child mortality rate. *Pediatr Crit Care Med* 12: 114-115
54. Approximating the GNI per capita used to differentiate between a lower-middle and upper-middle income country by the World Bank. In: Editor (ed)^(eds) Book Approximating the GNI per capita used to differentiate between a lower-middle and upper-middle income country by the World Bank. City, pp.

55. Assuncao MS, Teich V, Shiramizo SC, Araujo DV, Carrera RM, Serpa Neto A, Silva E, (2014) The cost-effectiveness ratio of a managed protocol for severe sepsis. *J Crit Care* 29: 692 e691-696
56. Harrison M, Collins CD, (2015) Is procalcitonin-guided antimicrobial use cost-effective in adult patients with suspected bacterial infection and sepsis? *Infect Control Hosp Epidemiol* 36: 265-272
57. Huang DT, Clermont G, Dremsizov TT, Angus DC, Pro CI, (2007) Implementation of early goal-directed therapy for severe sepsis and septic shock: A decision analysis. *Crit Care Med* 35: 2090-2100
58. Jones AE, Troyer JL, Kline JA, (2011) Cost-effectiveness of an emergency department-based early sepsis resuscitation protocol. *Crit Care Med* 39: 1306-1312
59. Karlsson S, Ruokonen E, Varpula T, Ala-Kokko TI, Pettila V, Finnsepsis Study G, (2009) Long-term outcome and quality-adjusted life years after severe sepsis. *Crit Care Med* 37: 1268-1274
60. Lehmann LE, Herpichboehm B, Kost GJ, Kollef MH, Stuber F, (2010) Cost and mortality prediction using polymerase chain reaction pathogen detection in sepsis: evidence from three observational trials. *Crit Care* 14: R186
61. Talmor D, Greenberg D, Howell MD, Lisbon A, Novack V, Shapiro N, (2008) The costs and cost-effectiveness of an integrated sepsis treatment protocol. *Crit Care Med* 36: 1168-1174
62. Negrini D, Sheppard L, Mills GH, Jacobs P, Rapoport J, Bourne RS, Guidet B, Csomos A, Prien T, Anderson G, Edbrooke DL, (2006) International Programme for Resource Use in Critical Care (IPOC)--a methodology and

- initial results of cost and provision in four European countries. *Acta Anaesthesiol Scand* 50: 72-79
63. Tan SS, Bakker J, Hoogendoorn ME, Kapila A, Martin J, Pezzi A, Pittoni G, Spronk PE, Welte R, Hakkaart-van Roijen L, (2012) Direct cost analysis of intensive care unit stay in four European countries: applying a standardized costing methodology. *Value Health* 15: 81-86
 64. Dräger S, Dal Poz MR, Evans DB, (2006) Health workers wages: an overview from selected countries. The WHO
 65. Parikh CR, Karnad DR, (1999) Quality, cost, and outcome of intensive care in a public hospital in Bombay, India. *Crit Care Med* 27: 1754-1759
 66. Kulkarni AP, Divatia JV, (2013) A prospective audit of costs of intensive care in cancer patients in India. *Indian J Crit Care Med* 17: 292-297
 67. Riviello ED, Kiviri W, Fowler RA, Mueller A, Novack V, Banner-Goodspeed VM, Weinkauff JL, Talmor DS, Twagirumugabe T, (2016) Predicting Mortality in Low-Income Country ICUs: The Rwanda Mortality Probability Model (R-MPM). *PLoS One* 11: e0155858
 68. Haniffa R, Lubell Y, Cooper BS, Mohanty S, Shamsul A, Karki A, Pattnaik R, Maswood A, Pangen R, Schultz MJ, Dondorp AM, Impact of a structured ICU training programme in resource-limited settings in Asia. *PLoS One*: submitted
 69. Penno EC, Baird SJ, Crump JA, (2015) Cost-Effectiveness of Surveillance for Bloodstream Infections for Sepsis Management in Low-Resource Settings. *Am J Trop Med Hyg* 93: 850-860
 70. Thukral A, Lodha R, Irshad M, Arora NK, (2006) Performance of Pediatric Risk of Mortality (PRISM), Pediatric Index of Mortality (PIM), and PIM2 in a

- pediatric intensive care unit in a developing country. *Pediatr Crit Care Med* 7: 356-361
71. Hutchings A, Durand MA, Grieve R, Harrison D, Rowan K, Green J, Cairns J, Black N, (2009) Evaluation of modernisation of adult critical care services in England: time series and cost effectiveness analysis. *BMJ* 339: b4353
 72. Rahman M, Fukui T, (2003) Biomedical publication--global profile and trend. *Public Health* 117: 274-280
 73. Bellani G, Laffey JG, Pham T, Fan E, Brochard L, Esteban A, Gattinoni L, van Haren F, Larsson A, McAuley DF, Ranieri M, Rubenfeld G, Thompson BT, Wrigge H, Slutsky AS, Pesenti A, Investigators LS, Group ET, (2016) Epidemiology, Patterns of Care, and Mortality for Patients With Acute Respiratory Distress Syndrome in Intensive Care Units in 50 Countries. *JAMA* 315: 788-800
 74. Fleischmann C, Scherag A, Adhikari NK, Hartog CS, Tsaganos T, Schlattmann P, Angus DC, Reinhart K, International Forum of Acute Care T, (2016) Assessment of Global Incidence and Mortality of Hospital-treated Sepsis. Current Estimates and Limitations. *Am J Respir Crit Care Med* 193: 259-272
 75. Vincent JL, Marshall JC, Namendys-Silva SA, Francois B, Martin-Loeches I, Lipman J, Reinhart K, Antonelli M, Pickkers P, Njimi H, Jimenez E, Sakr Y, investigators I, (2014) Assessment of the worldwide burden of critical illness: the intensive care over nations (ICON) audit. *Lancet Respir Med* 2: 380-386
 76. Kang KT, Chandler HK, Espinosa V, Kissoon N, (2014) Systems for Paediatric Sepsis: A Global Survey. *West Indian Med J* 63: 703-710

77. ISARIC (2016) International Severe Acute Respiratory and Emerging Infection Consortium (<https://isaric.tghn.org/about/>). In: Editor (ed)^(eds) Book International Severe Acute Respiratory and Emerging Infection Consortium (<https://isaric.tghn.org/about/>). City, pp.
78. Ashley EA, Dhorda M, Fairhurst RM, Amaratunga C, Lim P, Suon S, Sreng S, Anderson JM, Mao S, Sam B, Sopha C, Chuor CM, Nguon C, Sovannaroeth S, Pukrittayakamee S, Jittamala P, Chotivanich K, Chutasmit K, Suchatsoonthorn C, Runcharoen R, Hien TT, Thuy-Nhien NT, Thanh NV, Phu NH, Htut Y, Han KT, Aye KH, Mokuolu OA, Olaosebikan RR, Folaranmi OO, Mayxay M, Khanthavong M, Hongvanthong B, Newton PN, Onyamboko MA, Fanello CI, Tshefu AK, Mishra N, Valecha N, Phyo AP, Nosten F, Yi P, Tripura R, Borrmann S, Bashraheil M, Peshu J, Faiz MA, Ghose A, Hossain MA, Samad R, Rahman MR, Hasan MM, Islam A, Miotto O, Amato R, MacInnis B, Stalker J, Kwiatkowski DP, Bozdech Z, Jeeyapant A, Cheah PY, Sakulthaew T, Chalk J, Intharabut B, Silamut K, Lee SJ, Vihokhern B, Kunasol C, Imwong M, Tarning J, Taylor WJ, Yeung S, Woodrow CJ, Flegg JA, Das D, Smith J, Venkatesan M, Plowe CV, Stepniewska K, Guerin PJ, Dondorp AM, Day NP, White NJ, Tracking Resistance to Artemisinin C, (2014) Spread of artemisinin resistance in *Plasmodium falciparum* malaria. *N Engl J Med* 371: 411-423
79. Chantratita N, Tandhavanant S, Myers ND, Seal S, Arayawichanont A, Kiangsa-Ad A, Hittle LE, Ernst RK, Emond MJ, Wurfel MM, Day NP, Peacock SJ, West TE, (2013) Survey of innate immune responses to *Burkholderia pseudomallei* in human blood identifies a central role for lipopolysaccharide. *PLoS One* 8: e81617

80. Desjardins CA, Cohen KA, Munsamy V, Abeel T, Maharaj K, Walker BJ, Shea TP, Almeida DV, Manson AL, Salazar A, Padayatchi N, O'Donnell MR, Mlisana KP, Wortman J, Birren BW, Grosset J, Earl AM, Pym AS, (2016) Genomic and functional analyses of *Mycobacterium tuberculosis* strains implicate *ald* in D-cycloserine resistance. *Nat Genet* 48: 544-551
81. Kim JY, Farmer P, Porter ME, (2013) Redefining global health-care delivery. *Lancet* 382: 1060-1069
82. Scott KW, Jha AK, (2014) Putting quality on the global health agenda. *N Engl J Med* 371: 3-5
83. Binagwaho A, Kyamanywa P, Farmer PE, Nuthulaganti T, Umubyeyi B, Nyemazi JP, Mugeni SD, Asiimwe A, Ndagijimana U, Lamphere McPherson H, Ngirabega Jde D, Sliney A, Uwayezu A, Rusanganwa V, Wagner CM, Nutt CT, Eldon-Edington M, Cancedda C, Magaziner IC, Goosby E, (2013) The human resources for health program in Rwanda--new partnership. *N Engl J Med* 369: 2054-2059
84. Vukoja M, Kashyap R, Gavrilovic S, Dong Y, Kilickaya O, Gajic O, (2015) Checklist for early recognition and treatment of acute illness: International collaboration to improve critical care practice. *World J Crit Care Med* 4: 55-61
85. Baker T, Schell CO, Lugazia E, Blixt J, Mulungu M, Castegren M, Eriksen J, Konrad D, (2015) Vital Signs Directed Therapy: Improving Care in an Intensive Care Unit in a Low-Income Country. *PLoS One* 10: e0144801
86. Riviello ED, Kiviri W, Twagirimugabe T, Mueller A, Banner-Goodspeed VM, Officer L, Novack V, Mutumwinka M, Talmor DS, Fowler RA, (2016) Hospital Incidence and Outcomes of the Acute Respiratory Distress Syndrome Using

- the Kigali Modification of the Berlin Definition. *Am J Respir Crit Care Med* 193: 52-59
87. Chu KM, Jayaraman S, Kyamanywa P, Ntakiyiruta G, (2014) Building research capacity in Africa: equity and global health collaborations. *PLoS Med* 11: e1001612
 88. Singer M, Deutschman CS, Seymour CW, Shankar-Hari M, Annane D, Bauer M, Bellomo R, Bernard GR, Chiche JD, Coopersmith CM, Hotchkiss RS, Levy MM, Marshall JC, Martin GS, Opal SM, Rubenfeld GD, van der Poll T, Vincent JL, Angus DC, (2016) The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA* 315: 801-810
 89. Arabi YM, Schultz MJ, Salluh JI, (2016) Intensive Care Medicine in 2050: global perspectives. *Intensive Care Med*
 90. Maitland K, Kiguli S, Opoka RO, Engoru C, Olupot-Olupot P, Akech SO, Nyeko R, Mtove G, Reyburn H, Lang T, Brent B, Evans JA, Tibenderana JK, Crawley J, Russell EC, Levin M, Babiker AG, Gibb DM, Group FT, (2011) Mortality after fluid bolus in African children with severe infection. *N Engl J Med* 364: 2483-2495
 91. Riviello ED, Letchford S, Cook EF, Waxman AB, Gaziano T, (2015) Improving decision making for massive transfusions in a resource poor setting: a preliminary study in Kenya. *PLoS One* 10: e0127987
 92. Leligdowicz A, Fischer WA, 2nd, Uyeki TM, Fletcher TE, Adhikari NK, Portella G, Lamontagne F, Clement C, Jacob ST, Robinson L, Vanderschuren A, Hajek J, Murthy S, Ferri M, Crozier I, Ibrahima E, Lamah MC, Schieffelin JS, Brett-Major D, Bausch DG, Shindo N, Chan AK, O'Dempsey T, Mishra S,

- Jacobs M, Dickson S, Lyon GM, 3rd, Fowler RA, (2016) Ebola virus disease and critical illness. *Crit Care* 20: 217
93. Buist AS, Parry V, (2013) The American Thoracic Society methods in epidemiologic, clinical, and operations research program. A research capacity-building program in low- and middle-income countries. *Ann Am Thorac Soc* 10: 281-289
 94. Howitt P, Darzi A, Yang GZ, Ashrafian H, Atun R, Barlow J, Blakemore A, Bull AM, Car J, Conteh L, Cooke GS, Ford N, Gregson SA, Kerr K, King D, Kulendran M, Malkin RA, Majeed A, Matlin S, Merrifield R, Penfold HA, Reid SD, Smith PC, Stevens MM, Templeton MR, Vincent C, Wilson E, (2012) Technologies for global health. *Lancet* 380: 507-535
 95. Suarez D, Ferrer R, Artigas A, Azkarate I, Garnacho-Montero J, Goma G, Levy MM, Ruiz JC, Edusepsis Study G, (2011) Cost-effectiveness of the Surviving Sepsis Campaign protocol for severe sepsis: a prospective nationwide study in Spain. *Intensive Care Med* 37: 444-452

FIGURE LEGENDS

Figure 1. Most relevant infectious diseases as reported by the Global Burden of Diseases study stratified by the six World Health Organization regions. From [10, 11].

TABLES

Table 1 – Proposal for a categorization of intensive care units.

Table 2 – The cost–effectiveness of interventions for the management of severe sepsis.

FIGURE 1




Table 1. Proposal for a categorization of intensive care units*

Proposed categories	Category I – unrestricted	Category II – moderate restrictions	Category III – severe restrictions	Category IV – no formal ICU structure
Typical setting (not including private settings)	High-income countries	Higher-middle income countries	Lower-middle and low-income (major cities) countries	Rural areas of low-income countries
Formal ICU structure/service	Yes	Yes	Partly	No
Availability of specifically trained physicians and nurses	Widespread	Irregular	Rare	Unavailable
Availability of ICU equipment (i.e. patient monitor, mechanical ventilator, renal replacement therapy)	Unrestricted	Moderate restrictions (i.e. irregular maintenance of equipment, limited availability of advanced treatment modalities such as RRT)	Severe restrictions (i.e. basic monitoring typically available, limited number of mechanical ventilators, widespread unavailability of advanced treatment modalities such as RRT)	Unavailable
Availability of ICU drugs and disposable materials	Unrestricted	Mild restrictions	Moderate restrictions	Severe restrictions

*; The categories proposed here should not be seen as definite, but merely should serve as a starting point of future thinking

Abbreviations: ICU, intensive care unit; RRT, renal replacement therapy.

Table 2. The cost–effectiveness of interventions for the management of severe sepsis

Author, year [Reference]	Cohorts (Country)	Intervention	Conventional	Cost/QALY (in 2015 US \$)
Huang 2007 [57]	Monte Carlo simulation of patients with sepsis undergoing EGDT beginning either in the ED or ICU (US)	EGDT in ED or ICU	Standard care	\$3,506-8,953
Talmor 2008 [61]	Prospective cohort study of patients treated with an integrated sepsis protocol compared to controls (US)	Integrated sepsis protocol	Standard care	\$20,265 (2015)
Lehmann 2010 [60]	Monte Carlo simulation of sepsis episodes in ICU patients with use of PCR with cultures to identify the causative organism and initiate tailored antibiotic therapy (EU)	PCR and use cultures	Traditional cultures only	\$3,798
Karlsson 2009 [59]	Prospective cohort study of sepsis patients admitted to the ICU who were followed for 2 years (Finland)	ICU care for severe sepsis	Standard care	\$2,664
Jones 2011 [58]	Prospective before–after clinical trial to establish the initiation of EGDT in the ED prior to transfer to ICU for newly admitted sepsis patients (US)	EGDT for sepsis in ED	Standard ED care for sepsis	\$6,283 (2015)
Suarez 2011 [95]	Prospective before–after clinical trial to assess the impact of an educational program on adherence to ‘Surviving Sepsis Campaign Guidelines’ (Spain)	Surviving sepsis protocol for severe sepsis	Standard care	\$7,810 (2015)
Assuncao 2014 [55]	Prospective case–control analysis of patients treated with EGDT vs. standard care (Brazil)	EGDT	Standard care	Dominant
Harrison 2015 [56]	Markov model assessing cost-effectiveness of PCT as a diagnostic tool for bacterial infection in ICU patients (US)	PCT	Standard diagnostic techniques	Dominant

Abbreviations: EGDT, early goal directed therapy; ED, emergency department; ICU, intensive care unit; PCR, polymerase chain reaction; PCT, procalcitonin.

Textbox 1 – Availability of and indications for fluids in sepsis patients differ between high-income countries and low- and middle-income countries

Intravenous infusion of fluids is a key intervention in patients with sepsis as this syndrome frequently leads to intravascular hypovolemia from extravasation due to capillary leaking.

RESOURCE-RICH SETTINGS: physicians and nurses, independent from each other, have free and unlimited excess to fluids for intravenous infusion, which may facilitate a ‘too liberal’ fluid approach leading to ‘overzealous’ fluid resuscitation known to be associated with worse outcome of sepsis patients.

RESOURCE-POOR SETTINGS: usually there is a shortage of fluids for intravenous fluids, decision on the amounts of fluids to be given are restricted to attending physicians and not nurses, and the type of fluid chosen may depend on its price (e.g., dextrose containing fluids, such as 5% dextrose or 10% dextrose are cheap and thus readily available, opposite to crystalloid solutions); inadequate resuscitation, either because of too low or too late given, or wrong fluids infused, may worsen outcome of sepsis patients.

FLUID THERAPY IN BACTERIAL SEPSIS: the recommendations on fluid therapy in sepsis patients is largely built on evidence coming from randomized controlled trials in patients in RESOURCE-RICH SETTINGS, where most patients do have

BACTERIAL SEPSIS

FLUID THERAPY IN NON-BACTERIAL SEPSIS: Interestingly, randomized controlled trials including patients with severe falciparum malaria or dengue shock syndrome suggest more restricted fluid therapy than recommended for bacterial sepsis to be better. One major difference between BACTERIAL SEPSIS and NON-BACTERIAL SEPSIS is that fluid bolus resuscitation is not recommended in hyperlactatemic patients with severe falciparum malaria who are not hypotensive. Also, in Dengue Shock Syndrome proper fluid management is pivotal to ensure a good outcome; fluid therapy in Dengue Shock Syndrome should ensure adequate circulating volume to support tissue perfusion, but avoid overfilling, which will cause interstitial edema, which is in particular harmful in the lung.

Textbox 2 – Critical Care Expenditure in Indian Hospitals

In India critical care services are offered in four broad types of hospitals, with variable reimbursements. Variable reimbursements potentially leads to a great diversity in care of the critically ill, especially those who tend to have an extended ICU stay (i.e., longer than just a view days).

GOVERNMENTAL hospitals: the ICU bed, ventilation, basic medications may be free of costs, but patient will need to spend out of pocket for expensive medications like certain antibiotics, disposables and for the family to stay in a distinct place (far) away from their homes. Often in a few days they will exhaust their meager finances after which their care will be compromised

PUBLIC CHARITY TRUST hospitals, including hospitals of private medical colleges: these hospitals have a mandate to provide free treatment to patients below the poverty line. In addition, these hospitals also offer concessional treatment under various governmental insurance schemes for poor patients that, however, are mainly utilized for surgical patients and short stay patients like myocardial infarction and stroke. The insurance offered for sepsis is very meager and cannot cover more than the first few days in ICU. Patients thus will need to spend out of pocket for expensive medications like certain antibiotics, disposables and for the family to stay in a distinct place (far) away from their homes.

PRIVATE CORPORATE hospitals: These will treat affording patients either paying out of pocket or through some form of insurance. Except for the rich and very rich in a week or two such patient will exhaust their finances and will then either be transferred to GOVERNMENTAL hospitals or PUBLIC CHARITY TRUST hospitals. These transfers however are not easy given scarcity of beds. Public charitable hospitals will generally have an admission policy that will discourage patients that are deemed unsalvageable or may require some form of limitation of therapy.

GOVERNMENTAL hospitals have to accept such patients but here the beds crunch is often more severe. This leaves patient and care givers in a very difficult position and may lead to withdrawal of care or discharge against medical advice.

Smaller PRIVATE NURSING HOMES: these may offer critical services for selected patients but this too follows a similar trajectory as that of the PRIVATE CORPORATE hospitals, i.e., once patients' finances are exhausted they need to be transferred.