

## **Guidelines**

**Title: A proposal for standardized magnetic resonance imaging (MRI) descriptors of abnormally invasive placenta (AIP)- From the International Society for AIP**

**Short title: Standardized MRI descriptors for AIP**

**Abstract:** Abnormally invasive placenta (AIP) is used to describe a placenta that does not separate naturally after delivery and cannot be extirpated without causing abnormally high blood loss. Recently, the use of a standardized terminology for descriptors of AIP signs seen on ultrasound has been proposed but to date no such unified descriptors have been developed for magnetic resonance imaging (MRI). The purpose of this paper is to propose a unified terminology based on a consensus opinion from the members of the International Society for AIP (IS-AIP) that include obstetricians, gynecologists, radiologists, pathologists, anesthesiologists and basic science researchers. We assume that using these standardized MRI descriptors for AIP will be useful for clinical use, education, teaching and future research projects, thus assumably improving care of patients with this condition. In addition, using a uniform terminology for AIP should become the first step of a standardized MRI report.

**Index terms:** Abnormally invasive placenta (AIP); Magnetic resonance imaging (MRI); International Society for AIP;

## **Introduction**

Abnormally invasive placenta (AIP) is used to describe a placenta that does not separate naturally after delivery and cannot be extirpated without causing abnormally high blood loss. AIP is the consequence of an excessive invasion of the placenta within the myometrium. AIP may have different degrees of severity ranging histopathologically from placenta accreta (corresponding to abnormal adhesion of villous tissue that is attached directly to the

underlying myometrium without interposing decidua) and placenta increta (corresponding to a deeper invasion of the myometrium) to placenta percreta (corresponding to invasion of adjacent organs, most often the bladder). Whatever the severity of placental invasion, AIP can be complicated by major and potentially life-threatening peripartum bleeding [1-3]. AIP is the leading cause of emergency obstetric hysterectomy in developed countries [4, 5] and maternal mortality has been reported to be as high as 7% in the presence of AIP [6].

The principal risk factor for AIP is a pre-existing underlying uterine scar, which conveys a relative risk of 7 to 8. Uterine scar is predominantly the consequence of a previous cesarean section. Other risk factors have been identified, including placenta previa or prior uterine curettage [7-9]. The incidence of AIP increased tenfold over the last fifty years, as a consequence of a dramatic increase in cesarean section rates. Recent North-American studies have reported an incidence of AIP of 1 for 533 births [9-11]. With such a trend, it can be anticipated that the rate of cesarean sections in the United States of America will cause 4504 cases of AIP per year in 2020, and 130 annual maternal deaths [12].

Recently, the European working group on AIP (EW-AIP) has proposed the use of a standardized terminology for descriptors of AIP seen using ultrasound [13]. However, it is now well-established that magnetic resonance imaging (MRI) has an important role in the diagnosis of AIP [14-16]. Accordingly, the use of a standardized terminology for MRI is needed.

To improve comparability of future studies, to increase diagnostic capabilities and to facilitate international collaboration, the International Society for AIP (formerly the EW-AIP) proposes herein the use of standardized definitions of descriptors for AIP using MRI.

### **A plea for prenatal diagnosis of AIP**

Maternal prognosis can be substantially improved by prenatal diagnosis of AIP [1, 4]. The risk of bleeding is high in case of intra-partum diagnosis when forced placental extirpation is attempted. By contrast, when the diagnosis of AIP is made before delivery, birth can be planned in referral delivery ward where all necessary means (transfusion, interventional radiology, resuscitation and surgical skills) are available. Prenatal diagnosis also allows anticipating management strategies that will best meet the maternal expectations. Various surgical options can indeed be discussed depending on clinical specific situations and local

experience, including conservative approach with a placenta left in situ in the absence of major bleeding, cesarean-hysterectomy or local resection of the placenta and placental bed [5-7].

However, prenatal diagnosis can be falsely positive, resulting in important iatrogenic consequences for the mother and the fetus. In such cases, cesarean section may be performed by median laparotomy under general anesthesia at a premature gestational age. It is recommended that women at risk of AIP have a prenatal specialized screening in order to be referred to a multidisciplinary referral center [1, 8-10].

### **Imaging tools for the diagnosis of AIP**

Ultrasound is the first-line imaging tool for the screening and diagnosis of AIP. However, the diagnostic capabilities of ultrasound are impaired by anatomical or physiological limitations such as unfavorable placental location or high maternal body mass index. In addition, the absence of standardized descriptors, until the recent proposal published by our group [13], might also have affected the performance of ultrasound in previous studies [14].

By contrast to ultrasound, the performance of MRI is not limited by the depth of examination. MRI can be performed in addition to ultrasound in order to improve the results of prenatal imaging [12]. In addition, full coverage of the pelvis by MRI allows reevaluation at anytime by different observers, which is a possibility that is usually not allowed by ultrasound. MRI, however, is more expensive and less available than ultrasound. MRI has been recommended as a second-line imaging tool since 2011 by the Royal College of Obstetrics and Gynecology [17]. The overall performance of MRI for the diagnosis of AIP was analyzed in a systematic review that included 11 studies in 2015 [18]. Of note, studies reporting the use of intravenous contrast-agents were excluded from the analysis [18]. Reported performance of MRI for the prenatal diagnosis of AIP had sensitivities ranging from 75 to 100%, specificity from 65 to 100%, positive predictive value from 67 to 94.4% and negative predictive value from 79 to 92% [14, 19-28]. To date, MRI has not demonstrated superiority over ultrasound in the diagnosis of AIP [29]. However, it has been suggested that MRI might provide a more accurate evaluation of the degree of invasion for placenta percreta than ultrasound does, and especially for posterior AIP [30, 31]. Early knowledge of this would facilitate the most appropriate multidisciplinary and surgical care.

Suboptimal performances of MRI for the diagnosis of AIP might be due in part to a lack of consensus regarding the terminology used for descriptors of individual abnormalities in AIP [32]. As with ultrasound, researchers often use personal descriptions for signs, so different terms may apply to the same observed abnormality [18]. It is therefore urgent to standardize the terminology to avoid further confusion. Furthermore, with MRI, descriptors are highly dependent on the MRI sequence used at the time of image acquisition, with marked variations among equipment and vendors. MRI performances also depend on the magnetic field used. To date, most publications have reported the results of MRI using a 1.5-Tesla unit although MRI examination performed at 3-Tesla appears to be safe during pregnancy [33-36].

### **The end of the Tower of Babel**

The International Society for Abnormally Invasive Placenta ([www.IS-AIP.org](http://www.IS-AIP.org); formerly the EW-AIP) is an international society, currently consisting of obstetricians, gynecologists, radiologists, pathologists, anesthesiologists and basic science researchers from eleven European countries. The aim of the group is to advance diagnosis and treatment and to promote research and knowledge on AIP. To improve comparability of future studies, to increase diagnostic capabilities and to facilitate international collaboration, the IS-AIP here proposes standardized definitions of AIP-MRI descriptors.

Studies reporting the use of MRI for antenatal diagnosis of AIP were analyzed [25, 30, 31, 37-46]. The exact wording used for MRI descriptors of AIP was tabulated. These descriptions were grouped according to MRI sequences (T1- or T2-weighted images). The different terms used were examined, grouped according to sign and then replaced by a single, unambiguous, straightforward descriptor. These descriptors were discussed by a panel of experts present at the 9<sup>th</sup> EW-AIP meeting in Brussels in November 2015, who consensually agreed upon. During this meeting, and in further discussions among all EW-AIP members, importance was placed on unambiguously defining each sign, irrespective of opinions regarding the diagnostic values of each descriptor. These unified descriptors were augmented by typical MR images that all experts agreed upon their characteristic features.

Thirteen papers were examined for wording used to describe AIP signs. These were extracted and grouped by MRI sequences, and synonymous or identical terms unified under a common heading. Five common descriptors for signs seen on T2-weighted MR images and

four for T2-weighted and T1-weighted MR sequences were agreed by the expert group, meant to unambiguously describe the sign or finding (IS-AIP suggestion). The occurrence of each descriptor is listed in Table 1. Compound signs (e.g. “myometrial thinning < 1mm AND focal disruption of myometrium”) were grouped in the same descriptors. MR imaging examples of the unified descriptors are provided in Figures 1 to 8.

We assume that these MRI descriptors for AIP will be useful for clinical use, education, teaching and future research projects. Using a uniform terminology for AIP should become the first step of a standardized MRI report. In addition to using a uniform terminology, exact description of placental location and the placental portion suspicious for abnormal invasion (the location of AIP) should be considered a standard requirement for describing affected cases.

## **Conclusion**

Maternal mortality and morbidity associated with AIP are reduced when women with this condition deliver in a tertiary referral center with an experienced multi-disciplinary team [1, 8, 9, 47]. Referral to such a team depends on the prenatal diagnosis of AIP by the primary healthcare providers. By clearly defining the AIP descriptors on MRI we provide strong support to this referral process. The rarity of AIP needs close collaboration between centers both nationally and internationally. Ensuring that all investigators are reporting the same MRI findings when referring to a specific sign will improve homogeneity of data collection making results more valid. Based on these new descriptors we are currently collecting evidence and expert opinions for the predictive value of these signs, aiming to develop guidelines for diagnosing and managing AIP.

**Conflicts of interest:** The authors have no conflicts of interest to disclose.

## **References**

1. Chantraine F, Braun T, Gonser M, Henrich W, Tutschek B. Prenatal diagnosis of abnormally invasive placenta reduces maternal peripartum hemorrhage and morbidity. *Acta Obstet Gynecol Scand* 2013;92:439–44.

2. Soyer P, Dohan A, Dautry R, Guerrache Y, Ricbourg A, Gayat E, et al. Transcatheter arterial embolization for postpartum hemorrhage: indications, technique, results, and complications. *Cardiovasc Intervent Radiol* 2015;38:1068–81.
3. Soyer P, Morel O, Fargeaudou Y, Sirol M, Staub F, Boudiaf M, et al. Value of pelvic embolization in the management of severe postpartum hemorrhage due to placenta accreta, increta or percreta. *Eur J Radiol* 2011; 80:729–35.
4. Tikkanen M, Paavonen J, Loukovaara M, Stefanovic V. Antenatal diagnosis of placenta accreta leads to reduced blood loss. *Acta Obstet Gynecol Scand* 2011;90:1140–6.
5. Palacios-Jaraquemada JM. Caesarean section in cases of placenta praevia and accreta. *Best Pract Res Clin Obstet Gynaecol* 2013;27:221–32.
6. Chandrharan E, Rao S, Belli AM, Arulkumaran S. The Triple-P procedure as a conservative surgical alternative to peripartum hysterectomy for placenta percreta. *Int J Gynaecol Obstet* 2012;117:191–4.
7. Clausen C, Lönn L, Langhoff-Roos J. Management of placenta percreta: a review of published cases. *Acta Obstet Gynecol Scand* 2014;93:138–43.
8. Eller AG, Bennett MA, Sharshiner M, Masheter C, Soisson AP, Dodson M, et al. Maternal morbidity in cases of placenta accreta managed by a multidisciplinary care team compared with standard obstetric care. *Obstet Gynecol* 2011;117:331–7.
9. Al-Khan A, Gupta V, Illsley NP, Mannion C, Koenig C, Bogomol A, et al. Maternal and fetal outcomes in placenta accreta after institution of team-managed care. *Reprod Sci* 2014;21:761–71.
10. Wong HS, Hutton J, Zuccollo J, Tait J, Pringle KC. The maternal outcome in placenta accreta: the significance of antenatal diagnosis and non-separation of placenta at delivery. *N Z Med J* 2008;121:30–8.
11. D’Antonio F, Iacovella C, Bhide A. Prenatal identification of invasive placentation using ultrasound: systematic review and meta-analysis. *Ultrasound Obstet Gynecol* 2013;42:509–17.
12. Bour L, Placé V, Bendavid S, Fargeaudou Y, Portal JJ, Ricbourg A, et al. Suspected invasive placenta: evaluation with magnetic resonance imaging. *Eur Radiol* 2014;24:3150–60.
13. Collins SL, Ashcroft A, Braun T, Calda P, Langhoff-Roos J, Morel O, et al, European Working Group on Abnormally Invasive Placenta (EW-AIP): proposal for standardized ultrasound descriptors of abnormally invasive placenta (AIP). *Ultrasound Obstet Gynecol* 2016;47:271-5.

14. Derman AY, Nikac V, Haberman S, Zelenko N, Opsha O, Flyer M. MRI of placenta accreta: a new imaging perspective. *AJR Am J Roentgenol* 2011;197:1514–21.
15. Chen X, Shan R, Zhao L, Song Q, Zuo C, Zhang X, et al. Invasive placenta previa: Placental bulge with distorted uterine outline and uterine serosal hypervascularity at 1.5T MRI - useful features for differentiating placenta percreta from placenta accreta. *Eur Radiol* 2018;28:708-17.
16. Linduska N, Dekan S, Messerschmidt A, Kasprian G, Brugger PC, Chalubinski K, et al. Placental pathologies in fetal MRI with pathohistological correlation. *Placenta* 2009;30:555-9.
17. Royal College of Obstetricians and Gynaecologists. Placenta praevia, placenta praevia accreta and vasa praevia: diagnosis and management. In: RCOG Green-Top Guideline No 27 (3<sup>rd</sup> Edition), London RCOG 2011;1–26.
18. Rahaim NSA, Whitby EH. The MRI features of placental adhesion disorder and their diagnostic significance: systematic review. *Clin Radiol* 2015;70:917–25.
19. Peker N, Turan V, Ergenoglu M, Yeniel O, Sever A, Kazandi M, et al. Assessment of total placenta previa by magnetic resonance imaging and ultrasonography to detect placenta accreta and its variants. *Ginekol Pol* 2013;84:186–92.
20. Elhawary TM, Dabees NL, Youssef MA. Diagnostic value of ultrasonography and magnetic resonance imaging in pregnant women at risk for placenta accreta. *J Matern Fetal Neonatal Med* 2013;26:1443–9.
21. Maher MA, Abdelaziz A, Bazeed MF. Diagnostic accuracy of ultrasound and MRI in the prenatal diagnosis of placenta accreta. *Acta Obstet Gynecol Scand* 2013;92:1017–22.
22. Alamo L, Anaye A, Rey J, Denys A, Bongartz G, Terraz S, et al. Detection of suspected placental invasion by MRI: do the results depend on observer experience? *Eur J Radiol* 2013;82:51-7.
23. Shweel MAG, El Ameen NF, Ibrahiem MA, Kotib A. Placenta accreta in women with prior uterine surgery: diagnostic accuracy of Doppler ultrasonography and MRI. *Egypt J Radiol Nucl Med* 2012;43:473–80.
24. Mansour SM, Elkhyat WM. Placenta previa–accreta: do we need MR imaging? *Egypt J Radiol Nucl Med* 2011;42:433–42.
25. Lim PS, Greenberg M, Edelson MI, Bell KA, Edmonds PR, Mackey AM. Utility of ultrasound and MRI in prenatal diagnosis of placenta accreta: a pilot study. *AJR Am J Roentgenol* 2011;197:1506–13.

26. Teo TH, Law YM, Tay KH, Tan BS, Cheah FK. Use of magnetic resonance imaging in evaluation of placental invasion. *Clin Radiol* 2009; 64: 511–6.
27. Masselli G, Brunelli R, Casciani E, Poletti E, Piccioni MG, Anceschi M, Gualdi G. Magnetic resonance imaging in the evaluation of placental adhesive disorders: correlation with color Doppler ultrasound. *Eur Radiol* 2008;18:1292–9.
28. Dwyer BK, Belogolovkin V, Tran L, Rao A, Carroll I, Barth R, Chitkara U. Prenatal diagnosis of placenta accreta: sonography or magnetic resonance imaging? *J Ultrasound Med* 2008;27:1275–81.
29. Publications Committee, Society for Maternal-Fetal Medicine, Belfort MA. Placenta accreta. *Am J Obstet Gynecol* 2010;203:430–9.
30. Aitken K, Allen L, Pantazi S, Kingdom J, Keating S, Pollard L, Windrim R. MRI significantly improves disease staging to direct surgical planning for abnormal invasive placentation: A single centre experience. *J Obstet Gynaecol Can* 2016;38:246–51.
31. Kumar I, Verma A, Ojha R, Shukla RC, Jain M, Srivastava A. Invasive placental disorders: a prospective US and MRI comparative analysis. *Acta Radiol* 2017;58:121-8.
32. D'Antonio F, Iacovella C, Palacios-Jaraquemada J, Bruno CH, Manzoli L, Bhide A. Prenatal identification of invasive placentation using magnetic resonance imaging: systematic review and meta-analysis. *Ultrasound Obstet Gynecol* 2014;44:8-16
33. Wang PI, Chong ST, Kielar AZ, Kelly AM, Knoepp UD, Mazza MB, Goodsitt MM. Imaging of pregnant and lactating patients: part 1, evidence-based review and recommendations. *AJR Am J Roentgenol* 2012;198:778–84.
34. Wang PI, Chong ST, Kielar AZ, Kelly AM, Knoepp UD, Mazza MB, et al. Imaging of pregnant and lactating patients: part 2, evidence-based review and recommendations. *AJR Am J Roentgenol* 2012;198:785–92.
35. Expert Panel on MR Safety, Kanal E, Barkovich AJ, Bell C, Borgstede JP, Bradley WG, et al. ACR guidance document on MR safe practices: 2013. *J Magn Reson Imaging* 2013;37:501–30.
36. Patenaude Y, Pugash D, Lim K, Morin L; Diagnostic Imaging Committee, Lim K, et al; Society of Obstetricians and Gynaecologists of Canada. The use of magnetic resonance imaging in the obstetric patient. *J Obstet Gynaecol Can* 2014;36:349–63.
37. Budorick NE, Figueroa R, Vizcarra M, Shin J. Another look at ultrasound and magnetic resonance imaging for diagnosis of placenta accreta. *J Matern Fetal Neonatal Med* 2017;30:2422-7.

38. Lim G, Horowitz JM, Berggruen S, Ernst LM, Linn RL, Hewlett B, et al. Correlation of probability scores of placenta accreta on magnetic resonance imaging with hemorrhagic morbidity. *J Clin Anesth* 2016;34:261–9.
39. Balcacer P, Pahade J, Spektor M, Staib L, Copel JA, McCarthy S. Magnetic Resonance Imaging and Sonography in the diagnosis of placental invasion. *J Ultrasound Med* 2016;35:1445–56.
40. Kilcoyne A, Shenoy-Bhangle AS, Roberts DJ, Sisodia RC, Gervais DA, Lee SI. MRI of placenta accreta, placenta increta, and placenta percreta: pearls and pitfalls. *AJR Am J Roentgenol* 2017;208:214–21.
41. Thiravit S, Lapatikarn S, Muangsomboon K, Suvannarerg V, Thiravit P, Korpraphong P. MRI of placenta percreta: differentiation from other entities of placental adhesive disorder. *Radiol Med* 2017;122:61–8.
42. Ueno Y, Kitajima K, Kawakami F, Maeda T, Suenaga Y, Takahashi S, et al. Novel MRI finding for diagnosis of invasive placenta praevia: evaluation of findings for 65 patients using clinical and histopathological correlations. *Eur Radiol* 2014;24:881–8.
43. Cuthbert F, Teixidor Vinas M, Whitby E. The MRI features of placental adhesion disorder: a pictorial review. *Br J Radiol* 2016;89:20160284.
44. Riteau AS, Tassin M, Chambon G, Le Vaillant C, de Laveaucoupet J, Quéré MP, et al. Accuracy of ultrasonography and magnetic resonance imaging in the diagnosis of placenta accreta. *PloS One* 2014;9:948-66.
45. Azour L, Besa C, Lewis S, Kamath A, Oliver ER, Taouli B. The gravid uterus: MR imaging and reporting of abnormal placentation. *Abdom Radiol* 2016;41:2411–23.
46. Lax A, Prince MR, Mennitt KW, Schwebach JR, Budorick NE. The value of specific MRI features in the evaluation of suspected placental invasion. *Magn Reson Imaging* 2007;25:87–93.
47. Silver RM, Fox KA, Barton JR, Abuhamad AZ, Simhan H, Huls CK, Belfort MA, Wright JD. Center of excellence for placenta accreta. *Am J Obstet Gynecol* 2015;212:561–8.

## Figure legends

**Figure 1.** Heterogeneous placenta. 34-year-old woman with suspected abnormally invasive placenta (AIP) who had MR imaging of the pelvis at 36 weeks of pregnancy. T2-weighted HASTE MR image in the transverse plane shows ill-defined dark areas conferring heterogeneous appearance (arrows). AIP was confirmed intraoperatively.

**Figure 2.** Placental bulge. 39-year-old woman with suspected abnormally invasive placenta (AIP) who had MR imaging of the pelvis at 36 weeks of pregnancy. T2-weighted HASTE MR image of the pelvis in the sagittal plane shows placenta previa, abnormal anterior placental bulging with no visible uteroplacental interface (arrows) and placental protrusion through serosa. Intraoperative findings confirmed severe AIP.

**Figure 3.** Intraplacental dark bands. 29-year-old woman with suspected abnormally invasive placenta (AIP) who had MR imaging of the pelvis at 35 weeks of pregnancy. T2-weighted TrueFisp MR image of the pelvis in the sagittal plane shows placenta previa with intraplacental dark bands (arrowheads). Intraoperative findings confirmed AIP.

**Figure 4.** Loss of retroplacental dark zone. 35-year-old woman with suspected abnormally invasive placenta (AIP) who had MR imaging of the pelvis at 32 weeks of pregnancy. T2-weighted BLADE MR image of the pelvis in the coronal plane shows placenta previa with focal loss of retroplacental dark band (arrowhead). Intraoperative findings confirmed AIP.

**Figure 5.** Myometrial thinning. 31-year-old woman with suspected abnormally invasive placenta (AIP) who had MR imaging of the pelvis at 34 weeks of pregnancy. T2-weighted BLADE MR image of the pelvis in the sagittal plane shows placenta previa with myometrial thinning (less than 1 mm) (arrowhead). Intraoperative findings confirmed AIP.

**Figure 6.** Bladder wall interruption. 40-year-old woman with suspected abnormally invasive placenta (AIP) who had MR imaging of the pelvis at 34 weeks of pregnancy. T2-weighted HASTE MR image of the pelvis in the sagittal plane shows placenta previa with anterior bladder wall interruption (arrowhead). Intraoperative findings confirmed severe AIP.

**Figure 7.** Focal exophytic mass. 27-year-old woman with suspected abnormally invasive placenta (AIP) who had MR imaging of the pelvis at 35 weeks of pregnancy. T2-weighted HASTE MR image of the pelvis in the transverse plane shows right lateral focal exophytic mass (arrows). Intraoperative findings confirmed severe AIP.

**Figure 8.** Abnormal vascularisation of the placental bed. 39-year-old woman with suspected abnormally invasive placenta (AIP) who had MR imaging of the pelvis at 30 weeks of pregnancy. T2-weighted BLADE MR image of the pelvis in the sagittal plane shows abnormal vascularization of the placental bed (arrows). Intraoperative findings confirmed severe AIP.