SUPPORTING INFORMATION

Mixing natural and synthetic surfactants: co-adsorption of triterpenoid saponins and sodium dodecyl sulfate at the air-water interface.

I M Tucker¹, A Burley¹, R. E. Petkova¹, S. L. Hosking¹, J Penfold^{2,3#}, R K Thomas³, P X Li², J R P Webster², R Welbourn²

1. Unilever Research and Development, Port Sunlight Laboratory, Quarry Road East, Bebington, Wirral, UK
2. ISIS Facility, STFC, Rutherford Appleton Laboratory, Chilton, Didcot, OXON, UK
3. Physical and Theoretical Chemistry Laboratory, Oxford University, South Parks Road, Oxford, UK

Number of pages: 3

Number of figures: 2

Number of tables: 2

Figure S1. Surface tension versus concentration isotherms for SDS / escin mixtures, see legend for details

Figure S2. Variation in adsorbed amount versus surfactant concentration for 50/50 mole ratio SDS / escin in 0.1 M NaCl, for details see legend.

Table S1. Variation in mixed cmc for SDS / escin mixture.

Solution composition (mole% escin)	CMC (± 0.02 mM)	Surface tension above cmc (± 0.5 mN/m)
0.0	1.9	34
0.01	1.4	34
0.2	0.06	36
0.4	0.04	37
0.6	0.06	38
0.8	0.04	39
1.0	0.11	42

Table S2. Variation in adsorbed amount versus surfactant concentration for 50/50 mole ratio SDS / escin in 0.1 M NaCl.

Concentration (mM)	SDS		escin		Total ads ($\pm 0.05 \times 10^{-10}$ mol cm ⁻²)	Mole % escin (± 0.02)
	Area/molecule ($\pm 10 \text{ \AA}^2$)	Adsorbed amount ($\pm 0.05 \times 10^{-10}$ mol cm ⁻²)	Area/molecule ($\pm 10 \text{ \AA}^2$)	Adsorbed amount ($\pm 0.05 \times 10^{-10}$ mol cm ⁻²)		
0.3	209	0.79	85	1.95	2.74	0.71
0.6	109	1.52	144	1.15	2.67	0.43
1.0	107	1.55	127	1.31	2.86	0.46
1.5	98	1.69	205	0.81	2.50	0.32