



OPEN ACCESS

EDITED BY

Luigi Vetrugno,
Azienda Sanitaria Universitaria Friuli
Centrale, Italy

REVIEWED BY

Haoran Ma,
United Therapeutics Corporation,
United States
Glécia Virgolino da Silva Luz,
University of Brasília-UnB, Brazil

*CORRESPONDENCE

Patricia R. M. Rocco
✉ prmrocco@gmail.com

RECEIVED 09 December 2025

REVISED 15 January 2026

ACCEPTED 27 January 2026

PUBLISHED 12 February 2026

CITATION

Merola R, Battaglini D, Schultz MJ and
Rocco PRM (2026) Physiology-guided
personalized mechanical ventilation to
prevent ventilator-induced lung injury.
Front. Med. 13:1764151.
doi: 10.3389/fmed.2026.1764151

COPYRIGHT

© 2026 Merola, Battaglini, Schultz and
Rocco. This is an open-access article
distributed under the terms of the
[Creative Commons Attribution License
\(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or
reproduction in other forums is
permitted, provided the original author(s)
and the copyright owner(s) are credited
and that the original publication in this
journal is cited, in accordance with
accepted academic practice. No use,
distribution or reproduction is permitted
which does not comply with these terms.

Physiology-guided personalized mechanical ventilation to prevent ventilator-induced lung injury

Raffaele Merola¹, Denise Battaglini^{2,3}, Marcus J. Schultz^{4,5,6,7} and Patricia R. M. Rocco^{8*}

¹Anesthesia and Intensive Care Medicine, Department of Critical Care, AORN Ospedali Dei Colli, Naples, Italy, ²Department of Surgical Sciences and Integrated Diagnostics (DISC), University of Genoa, Genoa, Italy, ³Anesthesia and Intensive Care, IRCCS Ospedale Policlinico San Martino, Genoa, Italy, ⁴Department of Intensive Care, Amsterdam University Medical Centres, Amsterdam, Netherlands, ⁵Department of Anaesthesia, General Intensive Care and Pain Management, Medical University Wien, Vienna, Austria, ⁶Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom, ⁷Mahidol-Oxford Tropical Medicine Research Unit (MORU), Mahidol University, Bangkok, Thailand, ⁸Laboratory of Pulmonary Investigation, Carlos Chagas Filho Institute of Biophysics, Federal University of Rio de Janeiro, Rio de Janeiro, Brazil

Mechanical ventilation is essential for managing acute respiratory failure, yet it carries a significant risk of ventilator-induced lung injury (VILI). Lung-protective ventilation, most notably through the use of low tidal volumes, has improved outcomes in acute respiratory distress syndrome (ARDS), but these conventional strategies do not fully account for the profound heterogeneity of the injured lung or the variability in patient-specific physiology. Although tidal volumes of 4–8 ml/kg predicted body weight (PBW) provide a general reference for limiting strain, truly protective ventilation requires individualization based on regional aeration, compliance, and recruitability. Variability in these parameters leads to uneven distributions of stress and strain, while dynamic changes in respiratory drive, inspiratory effort, and cardiopulmonary interactions further complicate uniform ventilatory management. The mechanisms underlying VILI: barotrauma, volutrauma, atelectrauma, and biotrauma extend beyond the lung parenchyma and contribute to ventilator-associated diaphragm dysfunction and secondary organ injury. Bedside physiological tools, including esophageal manometry, electrical impedance tomography, and lung ultrasound, allow real-time evaluation of lung stress, regional ventilation, recruitability, and patient effort. When incorporated into clinical decision-making, these modalities facilitate individualized adjustments aimed at avoiding overdistension and collapse, limiting injurious pressures and volumes, and maintaining adequate gas exchange and hemodynamic stability. Advances in technology, such as closed-loop ventilation systems, adaptive control algorithms, and computational modeling, offer additional opportunities to refine personalized strategies and anticipate harmful mechanical patterns. Collectively, physiology-guided, personalized mechanical ventilation shifts practice from protocol-driven approaches to patient-centered care, with the overarching goal of mitigating VILI and improving outcomes in critically ill patients.

KEYWORDS

acute respiratory distress syndrome (ARDS), mechanical ventilation, respiratory mechanics, respiratory physiology, ventilator-induced lung injury (VILI)

Introduction

Mechanical ventilation (MV) is a cornerstone in the management of acute respiratory failure, including acute respiratory distress syndrome (ARDS) (1). While MV ensures adequate gas exchange, it may also aggravate or precipitate ventilator-induced lung injury (VILI) through excessive mechanical stress and strain (1, 2). The introduction of low tidal volume (VT) ventilation significantly reduced mortality in ARDS (3, 4), yet conventional protective strategies largely rely on global mechanical parameters that do not reflect the profound regional heterogeneity characteristic of ARDS (3).

In ARDS, loss of aerated lung volume, consolidation, gravitational gradients in pleural pressure, and non-uniform alveolar collapse generate steep regional disparities in stress and strain (5). As a result, ventilator settings that appear safe on a whole-lung basis may still impose injurious forces on vulnerable lung regions (6). These structural and functional asymmetries form the central limitation of uniform, population-based ventilation protocols (7, 8).

The increasing availability of bedside physiological monitoring: esophageal manometry, advanced mechanics, gas-exchange profiling, lung ultrasound (LUS), and electrical impedance tomography (EIT) (9, 10) reveals substantial inter- and in-patient variability in respiratory system behavior (11–13). These tools highlight that respiratory mechanics evolve dynamically with changes in disease severity, hemodynamics, sedation level, and systemic inflammation. Consequently, a physiology-guided, individualized approach to ventilation has gained traction as a strategy capable of addressing regional mechanical stresses more precisely (9).

It is now clear that VILI extends beyond the lungs. Mechanical stress and tissue deformation activate inflammatory pathways leading to systemic dissemination of cytokines, endothelial injury, and multiorgan dysfunction (1, 2, 14). This expanded understanding reinforces the need to refine ventilatory strategies that mitigate both local and systemic consequences of mechanical injury.

Despite substantial advances, fundamental questions remain: which physiological targets best predict benefit, how these measurements should guide bedside decision-making, and whether personalized ventilation improves clinical outcomes. This narrative review synthesizes contemporary physiological principles and emerging evidence supporting individualized mechanical ventilation in ARDS, with emphasis on regional mechanics, mechanical power, patient effort, and the systemic effects of VILI.

Ventilator-induced lung injury

Overview of VILI mechanisms

VILI encompasses several interrelated mechanisms: barotrauma, volutrauma, atelectrauma, and biotrauma, which together reflect the lung's response to excessive or unevenly distributed mechanical loads. More recent concepts, such as ergotrauma, viscoelastic stress, and regional mechanical

heterogeneity, extend beyond traditional global models and better capture the complex physiology of injured lungs (1, 2, 14).

Barotrauma and volutrauma: evolving concepts

Early animal studies shaped the foundational understanding of VILI. Webb and Tierney showed that extreme airway pressures caused severe lung injury in rats, but that applying PEEP prevented edema despite identical peak pressures, suggesting that pressure alone was insufficient to cause injury (15). Subsequently, Dreyfuss and colleagues demonstrated that excessive VT was the critical determinant of permeability injury, establishing excessive strain, rather than airway pressure, as the primary driver of volumetric trauma (16).

The “baby lung” model provided a conceptual breakthrough: ARDS reduces the size of the aerated lung available for ventilation (17). A fixed VT distributed over a smaller aerated lung volume results in disproportionately higher regional strain and inflammation. Subsequent computed tomography (CT) studies confirmed that reductions in respiratory system compliance (Crs) in ARDS primarily reflect loss of aerated volume, not increased intrinsic stiffness (18, 19).

Although historically treated as distinct entities, barotrauma and volutrauma are now understood as overlapping manifestations of excessive regional transpulmonary pressure. High airway pressures are not inherently harmful; the relevant distending force is the transpulmonary pressure, which depends on chest wall mechanics and pleural pressure (20, 21).

Atelectrauma and regional mechanical heterogeneity

Surfactant dysfunction, increased lung weight, and the supine position predispose ARDS patients to regional collapse (22). Cyclic opening and closing of unstable units produce shear forces that disrupt epithelial integrity and amplify inflammation (23). Stress may become several-fold greater at the interface between aerated and collapsed regions, where interdependence magnifies deformation (24).

Microscale heterogeneity is a defining feature of ARDS. Collapsed or partially aerated lung units that undergo intermittent reopening act as local stress multipliers, overdistingending adjacent aerated regions through mechanical interdependence and amplification of regional strain (25–27). This mechanism is closely related to the concept of stress concentrators in injured and heterogeneous lungs, whereby abrupt spatial transitions in aeration generate focal increases in stress and strain at the interfaces between collapsed and aerated tissue compartments (28). Such micro-atelectatic regions may remain undetectable on routine imaging yet contribute disproportionately to ventilator-induced lung injury.

PEEP modulates alveolar stability. Adequate PEEP prevents cyclic recruitment, whereas insufficient PEEP increases shear forces and promotes injury (29, 30). Clinically, low VT minimizes the

likelihood of exceeding local opening pressures, and appropriately set PEEP stabilizes alveoli throughout the respiratory cycle (31–34). However, excessive PEEP in non-recruitable or focal ARDS may preferentially ventilate already open regions, increasing overdistension and hemodynamic compromise (35, 36).

Biotrauma and systemic propagation of inflammation

Mechanical deformation triggers mechanotransduction cascades involving cytokine release, reactive oxygen species generation, and endothelial–epithelial disruption (37–39). These processes, collectively termed biotrauma, promote epithelial and endothelial injury, increased permeability, and extracellular matrix remodeling (37, 40–43). Lung-protective ventilation attenuates systemic inflammation and downstream organ dysfunction, underscoring the systemic implications of VILI (37, 40, 43, 44).

Ergotrauma: the central role of mechanical power

Mechanical power (MP), the energy delivered to the respiratory system per minute, integrates VT, respiratory rate (RR), pressures, and flow, offering a comprehensive measure of mechanical load (45). Although simplified bedside formulas exist, they vary in accuracy and their clinical applicability remains under evaluation (46–50).

Thresholds such as 17 J/min have been proposed but remain debated (51). Observational cohorts associate higher MP with mortality, with approximate thresholds of >17 J/min in mixed ICUs, >22 J/min in ARDS, and >14 J/min in the first 72 h of ECMO support (52–54). Experimental data indicate that MP above 25 J/min can cause structural injury (55).

Driving pressure (ΔP), defined as VT/Crs , remains a robust predictor of mortality and reflects functional lung size (6). The surrogate formula $(4 \times \Delta P) + RR$ has also been shown to predict outcomes after cardiac arrest (56, 57).

Viscoelasticity, strain dynamics, and time-dependent injury

The viscoelastic lung exhibits time-dependent stress–strain behavior. For a given VT, faster inflation increases strain rate and may be injurious. Strain is defined as VT normalized to end-expiratory lung volume (EELV); strain rate reflects the velocity of deformation (58). Regional time constants differ widely in ARDS. Units with short time constants fill rapidly, whereas slow units require prolonged inflation. High RR shorten inflation time, increasing heterogeneity and impairing gas exchange. RR has emerged as an independent contributor to VILI and mortality (56). In addition to inspiratory dynamics, emerging experimental evidence highlights the role of expiratory duration in modulating derecruitment.

Specifically, shortening expiratory time has been shown to limit cyclic alveolar collapse, preserve end-expiratory lung volume, and reduce regional stress amplification (59). In this context, time-controlled adaptive ventilation (TCAV), a form of airway pressure release ventilation (APRV) that dynamically constrains expiratory duration, has been associated in experimental and computational models with reduced derecruitment and attenuated lung injury (60, 61). Together, these findings support the concept that respiratory rate influences ventilator-induced lung injury not only through its effects on strain rate but also through time-dependent mechanisms governing recruitment–derecruitment dynamics.

The extracellular matrix accommodates small incremental strains, but abrupt increases in VT exceed its capacity to dissipate energy, increasing structural injury risk. Prolonged elevated strain leads to cumulative tissue damage (62).

Systemic effects of VILI

Diaphragm dysfunction

Controlled mechanical ventilation suppresses spontaneous effort and promotes ventilator-induced diaphragm dysfunction (VIDD). Oxidative stress, impaired mitochondrial biogenesis, proteolysis, and disrupted calcium handling contribute to atrophy and weakness, complicating liberation from MV (63, 64) (Figure 1).


Cardiopulmonary Interactions

Mechanical ventilation substantially alters heart–lung interactions (65). Increases in intrathoracic pressure (ITP) during positive-pressure ventilation decrease venous return, modify right ventricular (RV) preload, and may reduce cardiac output (65–68). These effects are particularly relevant in ARDS, where elevated pulmonary vascular resistance and varying degrees of RV dysfunction are common (69, 70) (Figure 1).

PEEP modulates these relationships in complex ways. Adequate PEEP can lower RV afterload by recruiting collapsed regions and reducing hypoxic vasoconstriction, whereas excessive PEEP increases ITP, limits venous return, and promotes overdistension, worsening RV performance. Fluid status further shapes these interactions: both hypovolemia and fluid overload can magnify adverse hemodynamic effects (71–73).

Systemic inflammation and organ dysfunction

Mechanical stress at the alveolar–capillary interface activates inflammatory pathways that propagate beyond the lung. Cytokines, chemokines, and damage-associated molecules {e.g. [interleukin (IL)-6, tumor necrosis factor (TNF)- α , IL-1 β , IL-8, C-X-C motif chemokine ligand [CXCL]10, CXCL12]} enter the circulation, promoting endothelial activation, glycocalyx disruption, and increased vascular permeability (74). These disturbances impair



microcirculatory flow and lead to tissue hypoperfusion. The resulting inflammatory profile resembles early sepsis, even without infection, and represents a key mechanism linking VILI to multiorgan dysfunction (75–77).

Renal, hepatic, and cardiovascular consequences

Circulating inflammatory mediators impair renal autoregulation, reduce cortical perfusion, and promote tubular injury, increasing the risk of acute kidney injury (78). The liver is similarly affected by inflammation and reduced blood flow, with potential consequences for metabolic and detoxification pathways (79). In the cardiovascular system, inflammatory signaling can depress myocardial contractility, raise pulmonary vascular resistance, and exacerbate RV strain. When combined with inadequate fluid management, these effects may result in pulmonary congestion and systemic venous hypertension (65) (Figure 1).

Neurological sequelae

Systemic inflammation also affects the brain. Disruption of the blood–brain barrier, alterations in cerebral blood flow related to PaCO₂ fluctuations, and exposure to sedatives contribute to

neuroinflammation and delirium (80). These disturbances may persist long after acute illness, adversely influencing long-term cognitive outcomes (81) (Figure 1).

Clinical implications

The extrapulmonary effects of VILI highlight the importance of physiology-based ventilation strategies. Approaches that reduce regional alveolar stress, such as low VT ventilation, careful PEEP titration, ΔP limitation, and preservation of spontaneous breathing when appropriate, may mitigate both pulmonary injury and its systemic consequences. Deeper understanding of how mechanical forces propagate through lung tissue and trigger multiorgan dysfunction will be crucial for developing targeted interventions.

The physiological foundations of personalized ventilation

Modern mechanical ventilation increasingly relies on individualized, physiology-based strategies. The heterogeneity of ARDS, characterized by reduced aerated lung volume, uneven regional compliance, and patchy consolidation, means that global ventilator settings often translate into markedly unequal regional forces. Even within conventionally “protective” ranges, pressures

and volumes may impose excessive strain on vulnerable units (1, 5, 7, 8).

Personalized ventilation is grounded in several principles: minimizing injurious mechanical loads, preventing both collapse and overdistension, preserving diaphragm function, and stabilizing cardiopulmonary interactions. These principles stem from an understanding that the “functional lung” in ARDS is considerably smaller than its anatomical volume and behaves as a highly heterogeneous, dynamically evolving structure (17). Achieving lung protection therefore requires tailoring ventilatory support to the individual patient’s mechanical properties, recruitability, spontaneous effort, and hemodynamic status (1, 5, 6, 65, 82).

This level of precision requires continuous physiological assessment. Bedside monitoring, via pressure–volume analysis, esophageal manometry, EIT, and LUS, helps characterize regional recruitment, overdistension, *pendelluft*, and ventilation–perfusion matching. These tools are especially valuable during transitions from controlled to assisted ventilation, when the risk of patient self-inflicted lung injury (P-SILI) is highest.

Respiratory system mechanics and dynamic indexes

Comprehensive evaluation of respiratory mechanics is essential for defining protective ventilator targets. Crs reflects the size of the aerated “baby lung” in ARDS more than total lung volume. Low Crs indicates a smaller ventilated lung and increased susceptibility to overdistension, even when VT is within the 4–8 ml/kg PBW range (5).

While VT normalized to PBW is reasonable for patients with structurally normal lungs, the height–lung size relationship underpinning PBW does not apply in ARDS. No reliable bedside technique exists to precisely quantify EELV, and its clinical utility remains uncertain. Crs may therefore provide a more physiologically meaningful estimate of functional lung size. Conceptually, normalizing VT to Crs is equivalent to normalizing it to ΔP . Airway resistance (R_{aw}) influences flow distribution and alveolar pressure dynamics. Elevations in R_{aw} worsen heterogeneity, increase intrinsic PEEP, and promote dynamic hyperinflation, especially at high RR (9, 83). Driving pressure has emerged as a robust marker of lung stress (84). Observational studies show that $\Delta P < 15$ cmH₂O, and ideally < 10 cmH₂O, is associated with improved survival (85). The STAMINA trial tested a ΔP -limited strategy incorporating Crs-optimized PEEP and VT titration, improving oxygenation, mechanics, and VILI biomarkers but not ventilator-free days (86). The trial’s early termination and brief intervention period limit interpretation, but its findings reinforce the central importance of early ΔP management (86). MP provides an integrative measure of mechanical load, but its clinical role continues to evolve (45).

Dynamic indices refine assessment of patient–ventilator interactions. Esophageal pressure monitoring estimates transpulmonary pressure, separating chest-wall from lung stress, especially relevant in patients with altered chest-wall mechanics. Measures of respiratory effort, including the esophageal pressure swing (ΔP_{es}), airway occlusion pressure (P0.1), and diaphragmatic thickening fraction on ultrasound, help identify

excessive spontaneous effort that could precipitate P-SILI (1, 82). Synchrony assessment identifies double triggering, breath stacking, and ineffective efforts, all linked to increased work of breathing, lung stress, and diaphragm injury (82, 87).

Recruitability assessment is central to individualized PEEP titration. Tools including pressure–volume curves, EIT-derived regional compliance, LUS re-aeration scoring, and gas-exchange changes during PEEP trials help determine whether additional PEEP will reopen collapsed tissue or induce overdistension (88, 89).

Heterogeneity in lung structure and function

Lung heterogeneity is a defining characteristic of ARDS and the primary rationale for individualized ventilation. Recruitability varies widely between patients. Those with highly recruitable, typically non-focal patterns may benefit from recruitment maneuvers and higher PEEP to reopen collapsed units and promote homogeneous ventilation. Conversely, individuals with focal or minimally recruitable disease are prone to overdistension and hemodynamic compromise when exposed to aggressive recruitment or high PEEP (90, 91).

The LIVE study illustrated this concept: misclassification of lung morphology resulted in inappropriate recruitment strategies and worse outcomes (92). Thus, bedside assessment of recruitability is essential. Tools such as the recruitment-to-inflation ratio, esophageal manometry, LUS, CT, and EIT offer complementary insights to balance recruitment against overdistension. The overarching goal is to achieve the most homogeneous ventilation pattern possible while minimizing regional stress and strain (73).

Respiratory drive and effort

Respiratory drive and inspiratory effort critically influence lung stress, synchrony, and diaphragm function during assisted ventilation. Excessive effort generates large negative pleural swings, increasing transpulmonary pressure and predisposing to P-SILI (13). Insufficient effort, in contrast, accelerates diaphragm disuse atrophy and prolongs weaning (93).

Monitoring neural, mechanical, and ventilatory parameters helps define an optimal “effort safe zone.” Diaphragm electrical activity (EAdi) provides continuous measurement of neural drive; although absolute values vary among patients, trends over time are clinically informative (94–98). Airway occlusion pressure (P0.1) reflects global neural drive and is independent of muscle fatigue or airway resistance (99, 100). Elevated P0.1 correlates with higher effort, dyspnea, prolonged ventilation, and increased mortality (101, 102).

Esophageal manometry allows direct quantification of pleural pressure swings, enabling estimation of inspiratory muscle pressure (P_{mus}) and identification of excessive or insufficient effort (45). The airway pressure deflection during an end-expiratory occlusion (ΔP_{occ}) provides a practical surrogate for P_{mus}, with extreme values associated with worse outcomes (103). Diaphragm ultrasound adds further insight; a thickening fraction

of approximately 30% often marks excessive effort, though considerable inter-individual variability persists (104).

Optimizing the alignment between patient effort and mechanical assistance is central to preserving diaphragm function and preventing occult lung stress. Achieving this balance requires integrating respiratory mechanics, recruitability, chest-wall compliance, and hemodynamics.

Physiology-guided protective strategies

Physiology-guided ventilation represents a shift from fixed, population-based targets toward individualized settings informed by mechanical, regional, and biological cues. Its goal is to define, for each patient, a protective window that minimizes VILI, diaphragm injury, and hemodynamic compromise.

PEEP and recruitment

PEEP stabilizes alveoli and reduces cyclic collapse, yet excessive levels can cause regional overdistension and reduce cardiac output. Trials comparing higher vs. lower PEEP strategies: ART (105), ALVEOLI (106), ExPress (107), and LOV (33), have produced inconsistent results, reflecting heterogeneity in patient characteristics, recruitability, and disease severity. A meta-analysis of ALVEOLI, ExPress, and LOV suggested a potential benefit in severe ARDS (108), although aggressive recruitment maneuvers remain associated with well-recognized risks.

Driving pressure provides a pragmatic tool for PEEP optimization, with the protective range often corresponding to the lowest ΔP (109). However, ΔP must be interpreted cautiously: alterations in lung volume, chest-wall elastance, airway closure, and intratidal recruitment can modify its relationship to true transpulmonary stress (9, 110).

Transpulmonary pressure estimated by esophageal manometry offers a more direct assessment of lung-distending forces and separates lung from chest-wall contributions (111, 112).

Imaging-based individualization

Imaging tools provide complementary insights for tailoring ventilator settings. CT remains the reference standard for characterizing lung morphology, quantifying recruitability, and identifying focal vs. non-focal patterns, though its use is limited by the need for transport and radiation exposure (19, 35, 113–123). EIT offers continuous bedside assessment of regional ventilation, enabling early detection of overdistension or collapse and supporting titration toward more homogeneous ventilation (124, 125). Evidence from clinical studies is mixed: some report reduced PEEP requirements or survival benefits compared with pressure–volume loop guidance, while broader use is constrained by equipment availability, operator dependency, and the need for consistent belt positioning (126–128). Lung ultrasound provides a practical, radiation-free assessment of aeration, consolidation, and

PEEP responsiveness, assists in classifying ARDS phenotypes, and can estimate opening and closing pressures during recruitment maneuvers (129–131).

Diaphragm-protective ventilation

Preserving diaphragmatic function has become a critical complement to lung protection. Maintaining inspiratory effort within a physiological range prevents both disuse atrophy from excessive unloading and injurious transpulmonary swings from uncontrolled spontaneous effort (63, 64). Continuous assessment of inspiratory effort supports fine-tuning of ventilatory assistance and helps balance respiratory drive with lung protection (132).

Early mobilization and structured reintroduction of spontaneous breathing, particularly with proportional modes, help retain contractility and maintain safe transpulmonary pressures. These strategies are especially important during the transition from controlled to assisted ventilation, when both diaphragm injury and occult lung stress are most likely (133–135).

Cardiovascular integration

Physiology-guided ventilation must incorporate cardiovascular considerations, recognizing the close interaction between intrathoracic pressures, preload, afterload, and RV function. Positive pressure ventilation alters venous return and pulmonary vascular resistance, effects that are amplified in ARDS due to elevated RV afterload and impaired RV-pulmonary arterial coupling.

Volume status critically modulates these interactions: hypovolemia exacerbates the reduction in venous return induced by positive pressure, whereas fluid overload increases RV preload and pulmonary vascular congestion, intensifying afterload on the failing RV. Thus, fluid stewardship is inseparable from ventilator titration. Adjustments to PEEP or ΔP must be interpreted in the context of hemodynamic reserve and preload responsiveness (8, 65–73).

Echocardiography and dynamic responsiveness indices (PPV, PLR-induced SV change) provide essential information on RV size, septal motion, and pulmonary pressures, helping identify patients at risk of hemodynamic compromise during ventilator titration (136–139).

Technological and conceptual advances enabling personalization

Recent technological innovations and growing understanding of ARDS pathophysiology have accelerated the shift from standardized protocols to individualized, physiology-driven ventilation. These advances support multimodal evaluation of lung mechanics, regional heterogeneity, inspiratory effort, and cardiopulmonary interactions to guide patient-specific adjustments.

Bedside monitoring tools, adaptive ventilation systems, integrated data platforms, and computational models now provide an unprecedented level of granularity for tailoring support to the evolving characteristics of the injured lung.

The technologies and monitoring approaches discussed in this review were selected to illustrate physiologically grounded and clinically relevant pathways toward personalized mechanical ventilation, rather than to provide an exhaustive inventory of available tools. As the field continues to evolve, additional methodologies and innovations will likely further expand and refine this framework.

Bedside monitoring and effort assessment

Tools for continuous assessment of Crs, ΔP , and inspiratory effort have matured considerably. Combined with advanced hemodynamic monitoring, they allow clinicians to quantify stress and strain, identify excessive respiratory drive, detect hemodynamic vulnerability, and titrate support within lung-, diaphragm-, and heart-protective domains (140). These features are particularly valuable in ARDS, where respiratory mechanics and cardiopulmonary status can evolve rapidly.

Closed-loop and adaptive ventilation systems

Closed-loop systems represent an important technological step forward. By integrating variables such as VT, RR, airway pressures, gas exchange, and dynamic respiratory mechanics, these systems adjust ventilator settings automatically in response to patient demand. Incorporation of parameters related to VILI risk enables maintenance of protective targets with fewer manual interventions (141, 142).

A systematic review of 51 randomized trials demonstrated that automated ventilation reduces clinician workload and performs comparably to expert clinicians in adjusting ventilator parameters (142). Automated oxygen-control systems increase time within target SpO₂ ranges and reduce staff burden relative to manual titration (143). Whether these technologies improve patient-centered outcomes remains uncertain and requires rigorous evaluation.

Importantly, in this section the term “adaptive ventilation” is used in a specific sense to denote algorithm-driven closed-loop systems, rather than clinician-directed ventilation strategies based on protocolized adjustments.

Computational modeling and machine learning

Emerging computational tools, including physiologic modeling, parameter estimation, and machine learning, offer

new opportunities to further refine individualized ventilatory care (144, 145). Recent advances have enabled models of varying complexity to be calibrated directly to patient-specific or experimental data, allowing a more detailed characterization of lung mechanics and time-dependent behavior. In principle, such approaches could support real-time assessment of regional mechanics, energy transfer, and spatial heterogeneity, thereby informing more precise ventilator adjustments and potentially reducing the risk of VILI (146, 147). However, clinical application remains at an early stage, and robust prospective validation is essential before widespread implementation.

Biological subphenotypes and precision medicine

Advances in ARDS subphenotyping also represent an important step toward precision medicine (148). Latent class analyses across multiple RCTs consistently identify biological subphenotypes, notably hyperinflammatory and hypoinflammatory, characterized by distinct inflammatory, endothelial, and coagulation profiles (149–153).

The hyperinflammatory subphenotype, marked by high cytokine levels, endothelial disruption, more severe shock, and metabolic acidosis, carries substantially higher mortality. Importantly, these subphenotypes respond differently to ventilatory, fluid, and pharmacologic strategies: patients with the hyperinflammatory phenotype appear to benefit from higher PEEP, conservative fluid management, and immunomodulatory therapies, including corticosteroids, whereas those with the hypoinflammatory phenotype derive less benefit and may be harmed by these same interventions (149–153).

These divergent responses parallel functional and radiographic distinctions between recruitable and non-recruitable lung morphologies, underscoring the importance of integrating physiologic assessment with biomarker-based classification. Prospective trials will be essential to determine whether subphenotype-guided strategies translate into improved outcomes.

Clinical translation and future perspectives

Although understanding of VILI has expanded substantially, key uncertainties persist. Lung-protective ventilation, particularly low VTs volumes and higher PEEP, remains central, yet survival benefits vary, especially regarding PEEP titration (33, 105–108). Newer approaches targeting lower ΔP and reduced MP offer additional protection (51–57, 84–86), but uniform application of low tidal volumes is unlikely to benefit all patients, and PEEP must be individualized to avoid overdistension or hemodynamic instability. These limitations reinforce the need for strategies grounded in each patient’s unique physiology.

Despite conceptual progress, several barriers impede clinical translation. Specialized monitoring tools, the need for precise calibration and placement, and cost considerations restrict




FIGURE 2

From monitoring to personalization: a physiology-driven approach to mechanical ventilation. Continuous monitoring of lung mechanics, patient respiratory effort, and advanced hemodynamic parameters provide real-time insight into pulmonary stress and strain, enabling early recognition of injurious breathing patterns and supporting ventilatory management within lung-, diaphragm-, and heart-protective ranges. The effective use of physiology-guided, individualized ventilation depends on coordinated interpretation of these data by the multidisciplinary team to balance patient effort with lung protection and to ensure safe transitions between controlled and assisted modes. Emerging technologies further strengthen this personalized approach: adaptive and closed-loop systems adjust ventilator settings in response to patient-specific changes, whereas machine-learning tools and digital-twin models integrate imaging, biomarkers, and physiologic signals to detect harmful trends, simulate treatment responses, and suggest optimal strategies. Together, these innovations shift clinical practice from reactive modification of ventilator settings to proactive, physiology-based optimization tailored to the individual patient. CO, cardiac output; Crs, respiratory system compliance; CT, chest computed tomography scan; EAdi, electrical activity of the diaphragm; EIT, electrical impedance tomography; LUS, lung ultrasound; MP, mechanical power; P0.1, airway occlusion pressure at 0.1 s; PL, transpulmonary pressure; PPV, pulse pressure variation; Raw, airway resistance; SVV, stroke volume variation; ΔP , driving pressure; ΔP_{occ} , airway pressure deflection during an end-expiratory occlusion; ΔP_{es} , esophageal pressure swing; and US, ultrasound.

widespread adoption. Many clinicians remain unfamiliar with interpreting advanced physiologic metrics or integrating multimodal data into moment-to-moment decisions. Continuous monitoring and frequent adjustments must also be coordinated with parallel ICU therapies, a challenge in high-acuity or resource-limited settings. Overcoming these barriers will require targeted training, improved user interfaces, and coordinated decision-support systems.

Personalized ventilation depends on integrated practice across the multidisciplinary ICU team. Intensivists, respiratory therapists, physiotherapists, and nurses must jointly interpret physiologic data, balance patient effort against lung protection, and manage transitions between ventilation modes safely (133, 134). Such collaboration is essential to avoid under- or over-assistance, minimize injury to both lung and diaphragm, and maintain hemodynamic stability.

Predictive modeling and digital-twin technologies, patient-specific simulations of lung mechanics and physiology, represent the next major step (154). By integrating demographic, imaging, biomarker, and real-time physiologic data, these tools may forecast regional stress patterns, guide ventilator selection, and anticipate

responses to therapeutic changes, enabling more anticipatory care (154, 155). Figure 2 shows an approach to mechanical ventilation based on physiology.

Ultimately, the goal is to prevent VILI rather than respond to its consequences. Achieving this will require the integration of advanced monitoring, predictive analytics, and adaptive ventilator adjustments to maintain regional stress and strain within safe limits while preserving diaphragmatic function and cardiopulmonary stability. Effective implementation will depend on rigorous trials, seamless incorporation into ICU workflows, and broad training to ensure that physiology-guided ventilation becomes standard practice rather than a specialized approach.

Conclusion

A comprehensive understanding of respiratory physiology remains fundamental to preventing iatrogenic injury during mechanical ventilation. Recognizing the heterogeneity of lung mechanics, patient effort, and cardiopulmonary interactions allows

clinicians to tailor ventilatory support that minimizes lung and diaphragm injury while ensuring adequate gas exchange. Personalized ventilation represents a transition from uniform, protocolized strategies to support that is dynamically adapted to the patient's evolving physiology. As multimodal monitoring, advanced analytics, and computational modeling become increasingly integrated into clinical care, they will strengthen the foundation for proactive, physiology-guided interventions that anticipate injury, preserve organ function, and ultimately improve outcomes in critically ill patients.

Author contributions

RM: Writing – review & editing, Writing – original draft. DB: Writing – review & editing, Supervision, Writing – original draft, Conceptualization. MS: Writing – original draft, Supervision, Writing – review & editing. PR: Supervision, Conceptualization, Writing – review & editing, Writing – original draft, Funding acquisition, Visualization, Investigation.

Funding

The author(s) declared that financial support was not received for this work and/or its publication.

References

- Merola R, Vargas M, Battaglini D. Ventilator-induced lung injury: the unseen challenge in acute respiratory distress syndrome management. *J Clin Med*. (2025) 14:3910. doi: 10.3390/jcm14113910
- Slutsky AS, Ranieri VM. Ventilator-induced lung injury. *N Engl J Med*. (2013) 369:2126–36. doi: 10.1056/NEJMra1208707
- Marini JJ. Evolving concepts in the ventilatory management of acute respiratory distress syndrome. *Clin Chest Med*. (1996) 17:555–75. doi: 10.1016/S0272-5231(05)70331-2
- Acute Respiratory Distress Syndrome Network, Brower RG, Matthay MA, Morris A, Schoenfeld D, Thompson BT. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. *N Engl J Med*. (2000) 342:1301–8. doi: 10.1056/NEJM200005043421801
- Gattinoni L, Tonetti T, Quintel M. Regional physiology of ARDS. *Crit Care*. (2017) 21:312. doi: 10.1186/s13054-017-1905-9
- Battaglini D, Fazzini B, Silva PL, Cruz FF, Ball L, Robba C, et al. Challenges in ARDS definition, management, and identification of effective personalized therapies. *J Clin Med*. (2023) 12:1381. doi: 10.3390/jcm12041381
- Pellegrini M, Del Sorbo L, Ranieri VM. Finding the optimal tidal volume in acute respiratory distress syndrome. *Intensive Care Med*. (2024) 50:1154–6. doi: 10.1007/s00134-024-07440-5
- Merola R, Vargas M, Battaglini D. Revisiting acute respiratory distress syndrome ventilation management: time for a paradigm shift focusing on tidal volume. *Respir Physiol Neurobiol*. (2025) 336:104454. doi: 10.1016/j.resp.2025.104454
- Pelosi P, Ball L, Barbas CSV, Bellomo R, Burns KEA, Einav S, et al. Personalized mechanical ventilation in acute respiratory distress syndrome. *Crit Care*. (2021) 25:250. doi: 10.1186/s13054-021-03686-3
- Pettenuzzo T, Zarantonello F, Zambianchi A, Lumetti G, Ruggerini D, Pacchiarini G, et al. Real-time stress and strain monitoring at the bedside: new frontiers in mechanical ventilation. *Front Med*. (2025) 12:1692488. doi: 10.3389/fmed.2025.1692488

Conflict of interest

The author(s) declared that this work was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Generative AI statement

The author(s) declared that generative AI was not used in the creation of this manuscript.

Any alternative text (alt text) provided alongside figures in this article has been generated by Frontiers with the support of artificial intelligence and reasonable efforts have been made to ensure accuracy, including review by the authors wherever possible. If you identify any issues, please contact us.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

- Kollisch-Singule MC, Jain SV, Andrews PL, Satalin J, Gatto LA, Villar J, et al. Looking beyond macroventilatory parameters and rethinking ventilator-induced lung injury. *J Appl Physiol* (1985). (2018) 124:1214–8. doi: 10.1152/jappphysiol.00412.2017
- Rixner M, Ludwig M, Lindner M, Frerichs I, Sablewski A, Wichmann KR, et al. Patient-specific prediction of regional lung mechanics in patients with ARDS with physics-based models: a validation study. *J Appl Physiol* (1985). (2025) 139:1029–49. doi: 10.1152/jappphysiol.00313.2025
- Marongiu I, Slobod D, Leali M, Spinelli E, Mauri T. Clinical and experimental evidence for patient self-inflicted lung injury (P-SILI) and bedside monitoring. *J Clin Med*. (2024) 13:4018. doi: 10.3390/jcm13144018
- Silva PL, Scharffenberg M, Rocco PRM. Understanding the mechanisms of ventilator-induced lung injury using animal models. *Intensive Care Med Exp*. (2023) 11:82. doi: 10.1186/s40635-023-00569-5
- Webb HH, Tierney DF. Experimental pulmonary edema due to intermittent positive pressure ventilation with high inflation pressures. Protection by positive end-expiratory pressure. *Am Rev Respir Dis*. (1974) 110:556–65. doi: 10.1164/arrd.1974.110.5.556
- Dreyfuss D, Soler P, Basset G, Saumon G. High inflation pressure pulmonary edema. Respective effects of high airway pressure, high tidal volume, and positive end-expiratory pressure. *Am Rev Respir Dis*. (1988) 137:1159–64. doi: 10.1164/ajrccm/137.5.1159
- Gattinoni L, Pesenti A. The concept of “baby lung”. *Intensive Care Med*. (2005) 31:776–84. doi: 10.1007/s00134-005-2627-z
- Protti A, Cressoni M, Santini A, Langer T, Mietto C, Febres D, et al. Lung stress and strain during mechanical ventilation: any safe threshold? *Am J Respir Crit Care Med*. (2011) 183:1354–62. doi: 10.1164/rccm.201010-1757OC
- Chiumello D, Marino A, Brioni M, Cigada I, Menga F, Colombo A, et al. Lung recruitment assessed by respiratory mechanics and computed tomography in patients with acute respiratory distress syndrome. What is the relationship? *Am J Respir Crit Care Med*. (2016) 193:1254–63. doi: 10.1164/rccm.201507-1413OC
- Bouhuys A. Physiology and musical instruments. *Nature*. (1969) 221:1199–204. doi: 10.1038/2211199a0

21. Gattinoni L, Protti A, Caironi P, Carlesso E. Ventilator-induced lung injury: the anatomical and physiological framework. *Crit Care Med.* (2010) 38:S539–48. doi: 10.1097/CCM.0b013e3181f1f27
22. Swenson KE, Swenson ER. Pathophysiology of acute respiratory distress syndrome and COVID-19 lung injury. *Crit Care Clin.* (2021) 37:749–76. doi: 10.1016/j.ccc.2021.05.003
23. Beitler JR, Malhotra A, Thompson BT. Ventilator-induced lung injury. *Clin Chest Med.* (2016) 37:633–46. doi: 10.1016/j.ccm.2016.07.004
24. Chen ZL, Song YL, Hu ZY, Zhang S, Chen YZ. An estimation of mechanical stress on alveolar walls during repetitive alveolar reopening and closure. *J Appl Physiol* (1985). (2015) 119:190–201. doi: 10.1152/jappphysiol.00112.2015
25. Cressoni M, Chiurazzi C, Gotti M, Amini M, Brioni M, Algieri I, et al. Lung inhomogeneities and time course of ventilator-induced mechanical injuries. *Anesthesiology.* (2015) 123:618–27. doi: 10.1097/ALN.0000000000000727
26. Retamal J, Hurtado D, Villarreal N, Bruhn A, Bugedo G, Amato MBP, et al. Does regional lung strain correlate with regional inflammation in acute respiratory distress syndrome during nonprotective ventilation? An Experimental Porcine Study. *Crit Care Med.* (2018) 46:e591–9. doi: 10.1097/CCM.0000000000003072
27. Albert K, Krischer JM, Pfaffenroth A, Wilde S, Lopez-Rodriguez E, Braun A, et al. Hidden microatelectases increase vulnerability to ventilation-induced lung injury. *Front Physiol.* (2020) 11:530485. doi: 10.3389/fphys.2020.530485
28. Gaver DP 3rd, Nieman GF, Gatto LA, Cereda M, Habashi NM, Bates JHT. The POOR Get POORer: a hypothesis for the pathogenesis of ventilator-induced lung injury. *Am J Respir Crit Care Med.* (2020) 202:1081–7. doi: 10.1164/rccm.202002-0453CP
29. Protti A, Andreis DT, Iapichino GE, Monti M, Comini B, Milesi M, et al. High positive end-expiratory pressure: only a dam against oedema formation? *Crit Care.* (2013) 17:R131. doi: 10.1186/cc12810
30. Halter JM, Steinberg JM, Gatto LA, DiRocco JD, Pavone LA, Schiller HJ, et al. Effect of positive end-expiratory pressure and tidal volume on lung injury induced by alveolar instability. *Crit Care.* (2007) 11:R20. doi: 10.1186/cc5695
31. Malhotra A. Low-tidal-volume ventilation in the acute respiratory distress syndrome. *N Engl J Med.* (2007) 357:1113–20. doi: 10.1056/NEJMct074213
32. Amato MB, Barbas CS, Medeiros DM, Magaldi RB, Schettino GP, Lorenzi-Filho G, et al. Effect of a protective-ventilation strategy on mortality in the acute respiratory distress syndrome. *N Engl J Med.* (1998) 338:347–54. doi: 10.1056/NEJM199802053380602
33. Meade MO, Cook DJ, Guyatt GH, Slutsky AS, Arabi YM, Cooper DJ, et al. Ventilation strategy using low tidal volumes, recruitment maneuvers, and high positive end-expiratory pressure for acute lung injury and acute respiratory distress syndrome: a randomized controlled trial. *JAMA.* (2008) 299:637–45. doi: 10.1001/jama.299.6.637
34. Caironi P, Cressoni M, Chiumello D, Ranieri M, Quintel M, Russo SG, et al. Lung opening and closing during ventilation of acute respiratory distress syndrome. *Am J Respir Crit Care Med.* (2010) 181:578–86. doi: 10.1164/rccm.200905-0787OC
35. Terragni PP, Rosboch G, Tealdi A, Corno E, Menaldo E, Davini O, et al. Tidal hyperinflation during low tidal volume ventilation in acute respiratory distress syndrome. *Am J Respir Crit Care Med.* (2007) 175:160–6. doi: 10.1164/rccm.200607-9150C
36. Grasso S, Stripoli T, De Michele M, Bruno F, Moschetta M, Angelelli G, et al. ARDSnet ventilatory protocol and alveolar hyperinflation: role of positive end-expiratory pressure. *Am J Respir Crit Care Med.* (2007) 176:761–7. doi: 10.1164/rccm.200702-1930C
37. Ranieri VM, Suter PM, Tortorella C, De Tullio R, Dayer JM, Brienza A, et al. Effect of mechanical ventilation on inflammatory mediators in patients with acute respiratory distress syndrome: a randomized controlled trial. *JAMA.* (1999) 282:54–61. doi: 10.1001/jama.282.1.54
38. Chiumello D, Pristine G, Slutsky AS. Mechanical ventilation affects local and systemic cytokines in an animal model of acute respiratory distress syndrome. *Am J Respir Crit Care Med.* (1999) 160:109–16. doi: 10.1164/ajrccm.160.1.9803046
39. Imai Y, Parodo J, Kajikawa O, de Perrot M, Fischer S, Edwards V, et al. Injurious mechanical ventilation and end-organ epithelial cell apoptosis and organ dysfunction in an experimental model of acute respiratory distress syndrome. *JAMA.* (2003) 289:2104–12. doi: 10.1001/jama.289.16.2104
40. Ranieri VM, Giunta F, Suter PM, Slutsky AS. Mechanical ventilation as a mediator of multisystem organ failure in acute respiratory distress syndrome. *JAMA.* (2000) 284:43–4. doi: 10.1001/jama.284.1.43
41. Parsons PE, Eisner MD, Thompson BT, Matthay MA, Ancukiewicz M, Bernard GR, et al. Lower tidal volume ventilation and plasma cytokine markers of inflammation in patients with acute lung injury. *Crit Care Med.* (2005) 33:1–6; discussion 230–2. doi: 10.1097/01.CCM.0000149854.61192.DC
42. Dolinay T, Kim YS, Howrylak J, Hunninghake GM, An CH, Fredenburgh L, et al. Inflammation-regulated cytokines are critical mediators of acute lung injury. *Am J Respir Crit Care Med.* (2012) 185:1225–34. doi: 10.1164/rccm.201201-0003OC
43. Agrawal A, Zhuo H, Brady S, Levitt J, Steingrub J, Siegel MD, et al. Pathogenetic and predictive value of biomarkers in patients with ALI and lower severity of illness: results from two clinical trials. *Am J Physiol Lung Cell Mol Physiol.* (2012) 303:L634–9. doi: 10.1152/ajplung.00195.2012
44. Barbata E, Ferrando C, López-Aladid R, Motos A, Bueno-Freire L, Fernández-Barat L, et al. Association between driving pressure, systemic inflammation and non-pulmonary organ dysfunction in patients with acute respiratory distress syndrome, a prospective pathophysiological study. *Anaesth Crit Care Pain Med.* (2025) 44:101458. doi: 10.1016/j.accpm.2024.101458
45. Gattinoni L, Tonetti T, Cressoni M, Cadringer P, Herrmann P, Moerer O, et al. Ventilator-related causes of lung injury: the mechanical power. *Intensive Care Med.* (2016) 42:1567–75. doi: 10.1007/s00134-016-4505-2
46. Cressoni M, Gotti M, Chiurazzi C, Massari D, Algieri I, Amini M, et al. Mechanical power and development of ventilator-induced lung injury. *Anesthesiology.* (2016) 124:1100–8. doi: 10.1097/ALN.0000000000001056
47. Tonetti T, Vasques F, Rapetti F, Maiolo G, Collino F, Romitti F, et al. Driving pressure and mechanical power: new targets for VILI prevention. *Ann Transl Med.* (2017) 5:286. doi: 10.21037/atm.2017.07.08
48. Paudel R, Trinkle CA, Waters CM, Robinson LE, Cassity E, Sturgill JL, et al. Mechanical power: a new concept in mechanical ventilation. *Am J Med Sci.* (2021) 362:537–45. doi: 10.1016/j.amjms.2021.09.004
49. Giosa L, Busana M, Pasticci I, Bonifazi M, Macri MM, Romitti F, et al. Mechanical power at a glance: a simple surrogate for volume-controlled ventilation. *Intensive Care Med Exp.* (2019) 7:61. doi: 10.1186/s40635-019-0276-8
50. Becher T, van der Staay M, Schädler D, Frerichs I, Weiler N. Calculation of mechanical power for pressure-controlled ventilation. *Intensive Care Med.* (2019) 45:1321–3. doi: 10.1007/s00134-019-05636-8
51. Gallardo A, Alcaráz M, Díaz-Cabrera A, Arriagada R, Rocco PRM, Battagliani D. Stepwise positive end-expiratory pressure titration modulates respiratory mechanics and mechanical power in mechanically ventilated adults. *Crit Care Sci.* (2025) 37:e20250252. doi: 10.62675/2965-2774.20250252
52. Serpa Neto A, Deliberato RO, Johnson AEW, Bos LD, Amorim P, Pereira SM, et al. Mechanical power of ventilation is associated with mortality in critically ill patients: an analysis of patients in two observational cohorts. *Intensive Care Med.* (2018) 44:1914–22. doi: 10.1007/s00134-018-5375-6
53. Parhar KKS, Zjadewicz K, Soo A, Sutton A, Zjadewicz M, Doig L, et al. Epidemiology, mechanical power, and 3-year outcomes in acute respiratory distress syndrome patients using standardized screening. An observational cohort study. *Ann Am Thorac Soc.* (2019) 16:1263–72. doi: 10.1513/AnnalsATS.201812-9100C
54. Chiu LC, Lin SW, Chuang LP, Li HH, Liu PH, Tsai FC, et al. Mechanical power during extracorporeal membrane oxygenation and hospital mortality in patients with acute respiratory distress syndrome. *Crit Care.* (2021) 25:13. doi: 10.1186/s13054-020-03428-x
55. Alkhalifah AS, Rumindo K, Brincat E, Blanchard F, Helleberg J, Clarke D, et al. Optimizing mechanical ventilation: personalizing mechanical power to reduce ICU mortality - a retrospective cohort study. *PLoS ONE.* (2025) 20:e0318018. doi: 10.1371/journal.pone.0318018
56. Costa ELV, Slutsky AS, Brochard LJ, Brower R, Serpa-Neto A, Cavalcanti AB, et al. Ventilatory variables and mechanical power in patients with acute respiratory distress syndrome. *Am J Respir Crit Care Med.* (2021) 204:303–11. doi: 10.1164/rccm.202009-3467OC
57. Robba C, Badenes R, Battagliani D, Ball L, Brunetti I, Jakobsen JC, et al. Ventilatory settings in the initial 72 h and their association with outcome in out-of-hospital cardiac arrest patients: a preplanned secondary analysis of the targeted hypothermia versus targeted normothermia after out-of-hospital cardiac arrest (T_{TM2}) trial. *Intensive Care Med.* (2022) 48:1024–38. doi: 10.1007/s00134-022-06756-4
58. Maeda Y, Fujino Y, Uchiyama A, Matsuura N, Mashimo T, Nishimura M. Effects of peak inspiratory flow on development of ventilator-induced lung injury in rabbits. *Anesthesiology.* (2004) 101:722–8. doi: 10.1097/0000542-200409000-00021
59. Al-Khalisy H, Nieman GF, Kollisch-Singule M, Andrews P, Camporota L, Shiber J, et al. Time-controlled adaptive ventilation (TCAV): a personalized strategy for lung protection. *Respir Res.* (2024) 25:37. doi: 10.1186/s12931-023-02615-y
60. Bates JHT, Kaczka DW, Kollisch-Singule M, Nieman GF, Gaver DP. Correction: Atelectrauma can be avoided if expiration is sufficiently brief: evidence from inverse modeling and oscillometry during airway pressure release ventilation. *Crit Care.* (2025) 29:14. doi: 10.1186/s13054-024-05227-0
61. Gaver DP. 3rd, Kollisch-Singule M, Nieman G, Satalin J, Habashi N, Bates JHT. Mechanical ventilation energy analysis: Recruitment focuses injurious power in the ventilated lung. *Proc Natl Acad Sci U S A.* (2025) 122:e2419374122. doi: 10.1073/pnas.2419374122
62. Felix NS, Samary CS, Cruz FF, Rocha NN, Fernandes MVS, Machado JA, et al. Gradually increasing tidal volume may mitigate experimental lung injury in rats. *Anesthesiology.* (2019) 130:767–77. doi: 10.1097/ALN.0000000000002630
63. Peñuelas O, Keough E, López-Rodríguez L, Carriedo D, Gonçalves G, Barreiro E, et al. Ventilator-induced diaphragm dysfunction: translational mechanisms lead to therapeutic alternatives in the critically ill. *Intensive Care Med Exp.* (2019) 7:48. doi: 10.1186/s40635-019-0259-9

64. Powers SK. Ventilator-induced diaphragm dysfunction: phenomenology and mechanism(s) of pathogenesis. *J Physiol.* (2024) 602:4729–52. doi: 10.1113/JP283860
65. Rocha NN, Silva PL, Battaglini D, Rocco PRM. Heart-lung crosstalk in acute respiratory distress syndrome. *Front Physiol.* (2024) 15:1478514. doi: 10.3389/fphys.2024.1478514
66. Mahmood SS, Pinsky MR. Heart-lung interactions during mechanical ventilation: the basics. *Ann Transl Med.* (2018) 6:349. doi: 10.21037/atm.2018.04.29
67. Marini JJ, Culver BH, Butler J. Mechanical effect of lung distention with positive pressure on cardiac function. *Am Rev Respir Dis.* (1981) 124:382–6. doi: 10.1164/arrd.1981.124.4.382
68. Corp A, Thomas C, Adlam M. The cardiovascular effects of positive pressure ventilation. *BJA Educ.* (2021) 21:202–9. doi: 10.1016/j.bjae.2021.01.002
69. Walkey AJ, Goligher EC, Del Sorbo L, Hodgson CL, Adhikari NKJ, Wunsch H, et al. Low tidal volume versus non-volume-limited strategies for patients with acute respiratory distress syndrome: a systematic review and meta-analysis. *Ann Am Thorac Soc.* (2017) 14:S271–9. doi: 10.1513/AnnalsATS.201704-337OT
70. McGuigan PJ, Bowcock EM, Barrett NA, Blackwood B, Boyle AJ, Cadamy AJ, et al. The effect of lower tidal volume ventilation facilitated by extracorporeal carbon dioxide removal compared with conventional lung protective ventilation on cardiac function. *Crit Care Explor.* (2024) 6:e1028. doi: 10.1097/CCE.0000000000001028
71. Ganerival S, Alves Dos Anjos G, Schleicher M, Hockstein MA, Tonelli AR, Duggal A, et al. Right ventricle-specific therapies in acute respiratory distress syndrome: a scoping review. *Crit Care.* (2023) 27:104. doi: 10.1186/s13054-023-04395-9
72. Joseph A, Petit M, Vieillard-Baron A. Hemodynamic effects of positive end-expiratory pressure. *Curr Opin Crit Care.* (2024) 30:10–9. doi: 10.1097/MCC.0000000000001124
73. Battaglini D, Roca O, Ferrer R. Positive end-expiratory pressure optimization in ARDS: physiological evidence, bedside methods and clinical applications. *Intensive Care Med.* (2024) 50:762–5. doi: 10.1007/s00134-024-07397-5
74. Ou D, Xu W, Feng Z, Yang Y, Xue W, Zhang Q, et al. Vascular endothelial glyocalyx shedding in ventilator-induced lung injury in rats. *Microvasc Res.* (2024) 153:104658. doi: 10.1016/j.mvr.2024.104658
75. Tutor JD, Mason CM, Dobard E, Beckerman RC, Summer WR, Nelson S. Loss of compartmentalization of alveolar tumor necrosis factor after lung injury. *Am J Respir Crit Care Med.* (1994) 149:1107–11. doi: 10.1164/ajrcm.149.5.8173748
76. Meduri GU, Headley S, Kohler G, Stentz F, Tolley E, Umberger R, et al. Persistent elevation of inflammatory cytokines predicts a poor outcome in ARDS. Plasma IL-1 beta and IL-6 levels are consistent and efficient predictors of outcome over time. *Chest.* (1995) 107:1062–73. doi: 10.1378/chest.107.4.1062
77. Del Sorbo L, Slutsky AS. Acute respiratory distress syndrome and multiple organ failure. *Curr Opin Crit Care.* (2011) 17:1–6. doi: 10.1097/MCC.0b013e3283427295
78. Hepokoski M, Englert JA, Baron RM, Crotty-Alexander LE, Fuster MM, Beitler JR, et al. Ventilator-induced lung injury increases expression of endothelial inflammatory mediators in the kidney. *Am J Physiol Renal Physiol.* (2017) 312:F654–60. doi: 10.1152/ajprenal.00523.2016
79. Beyer D, Hoff J, Sommerfeld O, Zipprich A, Gafner N, Press AT. The liver in sepsis: molecular mechanism of liver failure and their potential for clinical translation. *Mol Med.* (2022) 28:84. doi: 10.1186/s10020-022-00510-8
80. Mascia L, D'Albo R, Cavalli I, Giaccari L, Della Giovampaola M, Donati B. Organ crosstalk: brain-lung interaction. *Front Med.* (2025) 12:1655813. doi: 10.3389/fmed.2025.1655813
81. Millán Solano MV, Salinas Lara C, Sánchez-Garibay C, Soto-Rojas LO, Escobedo-Ávila I, Tena-Suck ML, et al. Effect of systemic inflammation in the CNS: a silent history of neuronal damage. *Int J Mol Sci.* (2023) 24:11902. doi: 10.3390/ijms241511902
82. Goligher EC, Dres M, Patel BK, Sahetya SK, Beitler JR, Telias I, et al. Lung- and diaphragm-protective ventilation. *Am J Respir Crit Care Med.* (2020) 202:950–61. doi: 10.1164/rccm.202003-0655CP
83. He Z, Wang C, Tian H. Dynamic nature of airway resistance in ARDS: methodological concerns in MPR calculations. *Intensive Care Med.* (2025) 51:450–1. doi: 10.1007/s00134-024-07771-3
84. Amato MB, Meade MO, Slutsky AS, Brochard L, Costa EL, Schoenfeld DA, et al. Driving pressure and survival in the acute respiratory distress syndrome. *N Engl J Med.* (2015) 372:747–55. doi: 10.1056/NEJMsa1410639
85. Urner M, Jüni P, Rojas-Saunero LP, Hansen B, Brochard LJ, Ferguson ND, et al. Limiting dynamic driving pressure in patients requiring mechanical ventilation. *Crit Care Med.* (2023) 51:861–71. doi: 10.1097/CCM.00000000000005844
86. Maia IS, Cavalcanti AB, Tramuja L, Veiga VC, Oliveira JS, Sady ERR, et al. Effect of a driving pressure-limiting strategy for patients with acute respiratory distress syndrome secondary to community-acquired pneumonia: the STAMINA randomised clinical trial. *Br J Anaesth.* (2025) 134:693–702. doi: 10.1016/j.bja.2024.10.012
87. de Haro C, Ochagavia A, López-Aguilar J, Fernandez-Gonzalo S, Navarra-Ventura G, Magrans R, et al. Patient-ventilator asynchronies during mechanical ventilation: current knowledge and research priorities. *Intensive Care Med Exp.* (2019) 7:43. doi: 10.1186/s40635-019-0234-5
88. Gattinoni L, Collino F, Maiolo G, Rapetti F, Romitti F, Tonetti T, et al. Positive end-expiratory pressure: how to set it at the individual level. *Ann Transl Med.* (2017) 5:288. doi: 10.21037/atm.2017.06.64
89. Gattinoni L, Collino F, Camporota L. Assessing lung recruitability: does it help with PEEP settings? *Intensive Care Med.* (2024) 50:749–51. doi: 10.1007/s00134-024-07351-5
90. Fosset M, von Wedel D, Redaelli S, Talmor D, Molinari N, Josse J, et al. Subphenotyping prone position responders with machine learning. *Crit Care.* (2025) 29:116. doi: 10.1186/s13054-025-05340-8
91. Pierrakos C, Smit MR, Pisani L, Paulus F, Schultz MJ, Constantin JM, et al. Lung ultrasound assessment of focal and non-focal lung morphology in patients with acute respiratory distress syndrome. *Front Physiol.* (2021) 12:730857. doi: 10.3389/fphys.2021.730857
92. Constantin JM, Jabaudon M, Lefrant JY, Jaber S, Quenot JP, Langeron O, et al. Personalised mechanical ventilation tailored to lung morphology versus low positive end-expiratory pressure for patients with acute respiratory distress syndrome in France (the LIVE study): a multicentre, single-blind, randomised controlled trial. *Lancet Respir Med.* (2019) 7:870–80. doi: 10.1016/S2213-2600(19)30138-9
93. Battaglini D, Rocco PRM. Beyond the ventilator: rethinking daily evaluation of respiratory drive and effort. *Lancet Respir Med.* (2025) 13:1045–6. doi: 10.1016/S2213-2600(25)00326-1
94. Piquilloud L, Beloncle F, Richard JM, Mancebo J, Mercat A, Brochard L. Information conveyed by electrical diaphragmatic activity during unstressed, stressed and assisted spontaneous breathing: a physiological study. *Ann Intensive Care.* (2019) 9:89. doi: 10.1186/s13613-019-0564-1
95. Spinelli E, Mauri T, Beitler JR, Pesenti A, Brodie D. Respiratory drive in the acute respiratory distress syndrome: pathophysiology, monitoring, and therapeutic interventions. *Intensive Care Med.* (2020) 46:606–18. doi: 10.1007/s00134-020-05942-6
96. Liu L, Liu H, Yang Y, Huang Y, Liu S, Beck J, et al. Neuroventilatory efficiency and extubation readiness in critically ill patients. *Crit Care.* (2012) 16:R143. doi: 10.1186/cc11451
97. Bellani G, Mauri T, Coppadoro A, Grasselli G, Patroniti N, Spadaro S, et al. Estimation of patient's inspiratory effort from the electrical activity of the diaphragm. *Crit Care Med.* (2013) 41:1483–91. doi: 10.1097/CCM.0b013e31827caba0
98. Stripoli T, Spadaro S, Di Mussi R, Volta CA, Trerotoli P, De Carlo F, et al. High-flow oxygen therapy in tracheostomized patients at high risk of weaning failure. *Ann Intensive Care.* (2019) 9:4. doi: 10.1186/s13613-019-0482-2
99. Whitelaw WA, Derenne JP, Milic-Emili J. Occlusion pressure as a measure of respiratory center output in conscious man. *Respir Physiol.* (1975) 23:181–99. doi: 10.1016/0034-5687(75)90059-6
100. Jonkman AH, de Vries HJ, Heunks LMA. Physiology of the respiratory drive in ICU patients: implications for diagnosis and treatment. *Crit Care.* (2020) 24:104. doi: 10.1186/s13054-020-2776-z. Erratum in: *Crit Care.* (2024) 28:94. doi: 10.1186/s13054-020-2776-z
101. Telias I, Junhasavasdikul D, Rittayamai N, Piquilloud L, Chen L, Ferguson ND, et al. Airway occlusion pressure as an estimate of respiratory drive and inspiratory effort during assisted ventilation. *Am J Respir Crit Care Med.* (2020) 201:1086–98. doi: 10.1164/rccm.201907-1425OC
102. Le Marec J, Hajage D, Decavèle M, Schmidt M, Laurent I, Ricard JD, et al. High airway occlusion pressure is associated with dyspnea and increased mortality in critically ill mechanically ventilated patients. *Am J Respir Crit Care Med.* (2024) 210:201–10. doi: 10.1164/rccm.202308-1358OC
103. Dianti J, Lovblom LE, Iftikhar MA, Sahetya S, Telias I, Urner M, et al. Association of respiratory drive and effort with mortality and time to discharge in patients on mechanical ventilation in Canada: a longitudinal, prospective, registry-based cohort study. *Lancet Respir Med.* (2025) 13:1087–95. doi: 10.1016/S2213-2600(25)00297-8
104. Dianti J, Fard S, Wong J, Chan TCY, Del Sorbo L, Fan E, et al. Strategies for lung- and diaphragm-protective ventilation in acute hypoxic respiratory failure: a physiological trial. *Crit Care.* (2022) 26:259. doi: 10.1186/s13054-022-04123-9
105. Writing Group for the Alveolar Recruitment for Acute Respiratory Distress Syndrome Trial (ART) Investigators, Cavalcanti AB, Suzumura EA, Laranjeira LN, Paisani DM, Damiani LP, et al. Effect of lung recruitment and titrated positive end-expiratory pressure (PEEP) vs low PEEP on mortality in patients with acute respiratory distress syndrome: a randomized clinical trial. *JAMA.* (2017) 318:1335–45. doi: 10.1001/jama.2017.14171
106. Brower RG, Lanken PN, MacIntyre N, Matthay MA, Morris A, Ancukiewicz M, et al. Higher versus lower positive end-expiratory pressures in patients with the acute respiratory distress syndrome. *N Engl J Med.* (2004) 351:327–36. doi: 10.1056/NEJMoa032193
107. Mercat A, Richard JC, Vielle B, Jaber S, Osman D, Diehl JL, et al. Positive end-expiratory pressure setting in adults with acute lung injury and acute respiratory distress syndrome: a randomized controlled trial. *JAMA.* (2008) 299:646–55. doi: 10.1001/jama.299.6.646
108. Briel M, Meade M, Mercat A, Brower RG, Talmor D, Walter SD, et al. Higher vs lower positive end-expiratory pressure in patients with acute lung injury and acute

- respiratory distress syndrome: systematic review and meta-analysis. *JAMA*. (2010) 303:865–73. doi: 10.1001/jama.2010.218
109. Chen L, Jonkman A, Pereira SM, Lu C, Brochard L. Driving pressure monitoring during acute respiratory failure in 2020. *Curr Opin Crit Care*. (2021) 27:303–10. doi: 10.1097/MCC.0000000000000827
110. Sakr Y, François B, Solé-Violan J, Kotfis K, Jaschinski U, Estella A, et al. Temporal changes in the epidemiology, management, and outcome from acute respiratory distress syndrome in European intensive care units: a comparison of two large cohorts. *Crit Care*. (2021) 25:87. doi: 10.1186/s13054-020-03455-8
111. Talmor D, Sarge T, Malhotra A, O'Donnell CR, Ritz R, Lisbon A, et al. Mechanical ventilation guided by esophageal pressure in acute lung injury. *N Engl J Med*. (2008) 359:2095–104. doi: 10.1056/NEJMoa0708638
112. Beitler JR, Sarge T, Banner-Goodspeed VM, Gong MN, Cook D, Novack V, et al. Effect of titrating positive end-expiratory pressure (PEEP) with an esophageal pressure-guided strategy vs an empirical high PEEP-Fio2 strategy on death and days free from mechanical ventilation among patients with acute respiratory distress syndrome: a randomized clinical trial. *JAMA*. (2019) 321:846–57. doi: 10.1001/jama.2019.0555
113. Giovanazzi S, Nocera D, Catozzi G, Collino F, Cressoni M, Ball L, et al. Assessment of recruitment from CT to the bedside: challenges and future directions. *Crit Care*. (2025) 29:64. doi: 10.1186/s13054-025-05263-4
114. Bitker L, Talmor D, Richard JC. Imaging the acute respiratory distress syndrome: past, present and future. *Intensive Care Med*. (2022) 48:995–1008. doi: 10.1007/s00134-022-06809-8
115. Cressoni M, Cadringer P, Chiurazzi C, Amini M, Gallazzi E, Marino A, et al. Lung inhomogeneity in patients with acute respiratory distress syndrome. *Am J Respir Crit Care Med*. (2014) 189:149–58. doi: 10.1164/rccm.201308-1567OC
116. Crotti S, Mascheroni D, Caironi P, Pelosi P, Ronzoni G, Mondino M, et al. Recruitment and derecruitment during acute respiratory failure: a clinical study. *Am J Respir Crit Care Med*. (2001) 164:131–40. doi: 10.1164/ajrccm.164.1.2007011
117. Pelosi P, Goldner M, McKibben A, Adams A, Eccher G, Caironi P, et al. Recruitment and derecruitment during acute respiratory failure: an experimental study. *Am J Respir Crit Care Med*. (2001) 164:122–30. doi: 10.1164/ajrccm.164.1.200710
118. Marini JJ. Hysteresis as an indicator of recruitment and ventilator-induced lung injury risk. *Crit Care Med*. (2020) 48:1542–3. doi: 10.1097/CCM.00000000000004533
119. Constantin JM, Grasso S, Chanques G, Aufort S, Futier E, Sebbane M, et al. Lung morphology predicts response to recruitment maneuver in patients with acute respiratory distress syndrome. *Crit Care Med*. (2010) 38:1108–17. doi: 10.1097/CCM.0b013e3181d451ec
120. Gattinoni L, Caironi P, Valenza F, Carlesso E. The role of CT-scan studies for the diagnosis and therapy of acute respiratory distress syndrome. *Clin Chest Med*. (2006) 27:559–70; Abstract VII. doi: 10.1016/j.ccm.2006.06.002
121. Gattinoni L, Pesenti A, Avalli L, Rossi F, Bombino M. Pressure-volume curve of total respiratory system in acute respiratory failure. Computed tomographic scan study. *Am Rev Respir Dis*. (1987) 136:730–6. doi: 10.1164/ajrccm/136.3.730
122. Protti A, Iapichino GE, Milesi M, Melis V, Pugni P, Comini B, et al. Validation of computed tomography for measuring lung weight. *Intensive Care Med Exp*. (2014) 2:31. doi: 10.1186/s40635-014-0031-0
123. Papazian L, Munshi L, Guérin C. Prone position in mechanically ventilated patients. *Intensive Care Med*. (2022) 48:1062–5. doi: 10.1007/s00134-022-06731-z
124. Battaglini D, Schultz MJ, Puentes GAC, Marini JJ, Rocco PRM. Imaging and pulmonary function techniques in ARDS diagnosis and management: current insights and challenges. *Crit Care*. (2025) 29:282. doi: 10.1186/s13054-025-05520-6
125. Frerichs I, Amato MB, van Kaam AH, Tingay DG, Zhao Z, Grychtol B, et al. Chest electrical impedance tomography examination, data analysis, terminology, clinical use and recommendations: consensus statement of the Translational EIT developmeNt stuDy group. *Thorax*. (2017) 72:83–93. doi: 10.1136/thoraxjnl-2016-208357
126. He H, Chi Y, Yang Y, Yuan S, Long Y, Zhao P, et al. Early individualized positive end-expiratory pressure guided by electrical impedance tomography in acute respiratory distress syndrome: a randomized controlled clinical trial. *Crit Care*. (2021) 25:230. doi: 10.1186/s13054-021-03645-y
127. Hsu HJ, Chang HT, Zhao Z, Wang PH, Zhang JH, Chen YS, et al. Positive end-expiratory pressure titration with electrical impedance tomography and pressure-volume curve: a randomized trial in moderate to severe ARDS. *Physiol Meas*. (2021) 42:014002. doi: 10.1088/1361-6579/abd679
128. Jimenez JV, Munroe E, Weirauch AJ, Fiorino K, Culter CA, Nelson K, et al. Electric impedance tomography-guided PEEP titration reduces mechanical power in ARDS: a randomized crossover pilot trial. *Crit Care*. (2023) 27:21. doi: 10.1186/s13054-023-04315-x
129. Cereda M, Xin Y, Goffi A, Herrmann J, Kaczka DW, Kavanagh BP, et al. Imaging the injured lung: mechanisms of action and clinical use. *Anesthesiology*. (2019) 131:716–49. doi: 10.1097/ALN.0000000000002583. Erratum in: *Anesthesiology*. (2019) 131:451. doi: 10.1097/ALN.0000000000002583
130. Boumans MMA, Aerts W, Pisani L, Bos LDJ, Smit MR, Tuinman PR. Diagnostic accuracy of lung ultrasound in diagnosis of ARDS and identification of focal or non-focal ARDS subphenotypes: a systematic review and meta-analysis. *Crit Care*. (2024) 28:224. doi: 10.1186/s13054-024-04985-1
131. Tusman G, Acosta CM, Costantini M. Ultrasonography for the assessment of lung recruitment maneuvers. *Crit Ultrasound J*. (2016) 8:8. doi: 10.1186/s13089-016-0045-9
132. Fu W, Guan L, Liu Q, Xie Z, You J, Chen R. Ventilator-induced diaphragmatic dysfunction: pathophysiology, monitoring and advances in potential treatment and prevention. *Eur Respir Rev*. (2025) 34:250069. doi: 10.1183/16000617.0069-2025
133. Battaglini D, Rocco PRM. Challenges in transitioning from controlled to assisted ventilation in acute respiratory distress syndrome (ARDS) management. *J Clin Med*. (2024) 13:7333. doi: 10.3390/jcm13237333
134. Arriagada R, San Martín C, Martí JD, Rocco PRM, Gosselink R, Schultz MJ, et al. Respiratory muscle dysfunction in older critically ill patients: implications for rehabilitation. *Anaesth Crit Care Pain Med*. (2025) 30:101657. doi: 10.1016/j.accpm.2025.101657
135. Pensier J, Battaglini D, Baedorf-Kassis E. Early mobilization in septic acute respiratory distress syndrome: the UNITE-COVID-II study. *Anaesth Crit Care Pain Med*. (2025) 44:101553. doi: 10.1016/j.accpm.2025.101553
136. Gavelli F, De Vita N, Lai C, Azzolina D, Pavot A, Jozwiak M, et al. Real-time changes in pulse pressure variation during a 10-second end-expiratory occlusion test reliably detect preload responsiveness. *Crit Care*. (2025) 29:241. doi: 10.1186/s13054-025-05483-8
137. Richard JC, Bayle F, Bourdin G, Leray V, Debord S, Delannoy B, et al. Preload dependence indices to titrate volume expansion during septic shock: a randomized controlled trial. *Crit Care*. (2015) 19:5. doi: 10.1186/s13054-014-0734-3
138. Monnet X, Teboul JL. Transpulmonary thermodilution: advantages and limits. *Crit Care*. (2017) 21:147. doi: 10.1186/s13054-017-1739-5
139. Mallat J, Siuba MT, Abou-Arab O, Kovacevic P, Ismail K, Duggal A, et al. Changes in pulse pressure variation induced by passive leg raising test to predict preload responsiveness in mechanically ventilated patients with low tidal volume in ICU: a systematic review and meta-analysis. *Crit Care*. (2025) 29:18. doi: 10.1186/s13054-024-05238-x
140. Battaglini D, Lassola S, Schultz MJ, Rocco PR. Innovations in protective mechanical ventilation for acute respiratory distress syndrome management. *Expert Rev Med Devices*. (2024) 21:789–92. doi: 10.1080/17434440.2024.2393773
141. Buiteman-Kruizinga LA, van Meenen DMP, Bos LDJ, van der Heiden PLJ, Paulus F, Schultz MJ, et al. closed-loop ventilation mode that targets the lowest work and force of breathing reduces the transpulmonary driving pressure in patients with moderate-to-severe ARDS. *Intensive Care Med Exp*. (2023) 11:42. doi: 10.1186/s40635-023-00527-1
142. Goossen RL, Schultz MJ, Tschernko E, Chew MS, Robba C, Paulus F, et al. Effects of closed loop ventilation on ventilator settings, patient outcomes and ICU staff workloads - a systematic review. *Eur J Anaesthesiol*. (2024) 41:438–46. doi: 10.1097/EJA.0000000000001972
143. Mól CG, Vieira AGDS, Garcia BMS, Dos Santos Pereira E, Eid RAC, Schultz MJ, et al. Closed-loop oxygen control for critically ill patients—a systematic review and meta-analysis. *PLoS ONE*. (2024) 19:e0304745. doi: 10.1371/journal.pone.0304745
144. Cecconi M, Greco M, Shickel B, Angus DC, Bailey H, Bignami E, et al. Implementing artificial intelligence in critical care medicine: a consensus of 22. *Crit Care*. (2025) 29:290. doi: 10.1186/s13054-025-05532-2
145. Fritsch SJ, Cecconi M. Setting the ventilator with AI support: challenges and perspectives. *Intensive Care Med*. (2025) 51:593–5. doi: 10.1007/s00134-024-07778-w
146. Lee JWW, Chiew YS, Wang X, Tan CP, Mat Nor MB, Cove ME, et al. Protocol conception for safe selection of mechanical ventilation settings for respiratory failure Patients. *Comput Methods Programs Biomed*. (2022) 214:106577. doi: 10.1016/j.cmpb.2021.106577
147. Lee JWW, Chiew YS, Wang X, Mat Nor MB, Chase JG, Desai T. Stochastic integrated model-based protocol for volume-controlled ventilation setting. *Biomed Eng Online*. (2022) 21:13. doi: 10.1186/s12938-022-00981-0
148. Merola R, Battaglini D. Redefining ARDS phenotypes: Challenges in the precision medicine era. *J Intensive Med*. (2025) 5:333–5. doi: 10.1016/j.jointm.2025.08.002
149. Calfee CS, Delucchi K, Parsons PE, Thompson BT, Ware LB, Matthay MA, et al. Subphenotypes in acute respiratory distress syndrome: latent class analysis of data from two randomised controlled trials. *Lancet Respir Med*. (2014) 2:611–20. doi: 10.1016/S2213-2600(14)70097-9
150. Famous KR, Delucchi K, Ware LB, Kangelaris KN, Liu KD, Thompson BT, et al. Acute respiratory distress syndrome subphenotypes respond differently to randomized fluid management strategy. *Am J Respir Crit Care Med*. (2017) 195:331–8. doi: 10.1164/rccm.201603-0645OC
151. Sinha P, Delucchi KL, Thompson BT, McAuley DF, Matthay MA, Calfee CS, et al. Latent class analysis of ARDS subphenotypes: a secondary analysis of the statins for acutely injured lungs from sepsis (SAILS) study. *Intensive Care Med*. (2018) 44:1859–69. doi: 10.1007/s00134-018-5378-3

152. Delucchi K, Famous KR, Ware LB, Parsons PE, Thompson BT, Calfee CS, et al. Stability of ARDS subphenotypes over time in two randomised controlled trials. *Thorax*. (2018) 73:439–45. doi: 10.1136/thoraxjnl-2017-211090
153. Calfee CS, Delucchi KL, Sinha P, Matthay MA, Hackett J, Shankar-Hari M, et al. Acute respiratory distress syndrome subphenotypes and differential response to simvastatin: secondary analysis of a randomised controlled trial. *Lancet Respir Med*. (2018) 6:691–8. doi: 10.1016/S2213-2600(18)30177-2
154. Silva A, Vale N. Digital twins in personalized medicine: bridging innovation and clinical reality. *J Pers Med*. (2025) 15:503. doi: 10.3390/jpm15110503
155. Caljé-van der Klei T, Sun Q, Chase JG, Zhou C, Tawhai MH, Knopp JL, et al. Pulmonary response prediction through personalized basis functions in a virtual patient model. *Comput Methods Programs Biomed*. (2024) 244:107988. doi: 10.1016/j.cmpb.2023.107988