

Preclinical models of endometriosis and interstitial cystitis/bladder pain syndrome: an Innovative Medicines Initiative-PainCare initiative to improve their value for translational research in pelvic pain

Paulina Nunez-Badinez^a, Bianca De Leo^{a,*}, Alexis Laux-Biehlmann^a, Anja Hoffmann^a, Thomas M. Zollner^a, Philippa T.K. Saunders^b, Ioannis Simitsidellis^b, Ana Charrua^c, Francisco Cruz^c, Raul Gomez^d, Miguel Angel Tejada^d, Stephen B. McMahon^e, Laure Lo Re^e, Florent Barthas^f, Katy Vincent^g, Judy Birch^h, Jane Meijlinkⁱ, Lone Hummelshøj^j, Patrick J. Sweeney^k, J. Douglas Armstrong^{k,l}, Rolf-Detlef Treede^m, Jens Nagel^a

Abstract

Endometriosis (ENDO) and interstitial cystitis/bladder pain syndrome (IC/BPS) are chronic pain conditions for which better treatments are urgently needed. Development of new therapies with proven clinical benefit has been slow. We have conducted a review of existing preclinical in vivo models for ENDO and IC/BPS in rodents, discussed to what extent they replicate the phenotype and pain experience of patients, as well as their relevance for translational research. In 1009 publications detailing ENDO models, 41% used autologous, 26% syngeneic, 18% xenograft, and 11% allogeneic tissue in transplantation models. Intraperitoneal injection of endometrial tissue was the subcategory with the highest construct validity score for translational research. From 1055 IC/BPS publications, most interventions were bladder centric (85%), followed by complex mechanisms (8%) and stress-induced models (7%). Within these categories, the most frequently used models were instillation of irritants (92%), autoimmune (43%), and water avoidance stress (39%), respectively. Notably, although pelvic pain is a hallmark of both conditions and a key endpoint for development of novel therapies, only a small proportion of the studies (models of ENDO: 0.5%-12% and models of IC/BPS: 20%-44%) examined endpoints associated with pain. Moreover, only 2% and 3% of publications using models of ENDO and IC/BPS investigated nonevoked pain endpoints. This analysis highlights the wide variety of models used, limiting reproducibility and translation of results. We recommend refining models so that they better reflect clinical reality, sharing protocols, and using standardized endpoints to improve reproducibility. We are addressing this in our project Innovative Medicines Initiative-PainCare/Translational Research in Pelvic Pain.

Keywords: Preclinical research, Endometriosis, Interstitial cystitis/bladder pain syndrome, Translational research, Pelvic pain, In vivo, Animal model, Rodent

1. Introduction

Animal models play a crucial role in research, helping to improve disease understanding and laying the foundation to develop new drug therapies. A wide spectrum of animal models has been

developed and proven to be a useful tool in translational research. Ideally, such models need to replicate the aetiology and symptoms of the disorder under investigation as closely as possible. However, achieving this ideal situation is no easy task and may be hampered by an incomplete understanding of the

Sponsorships or competing interests that may be relevant to content are disclosed at the end of this article.

^a Bayer AG, Research & Development, Pharmaceuticals, Berlin, Germany, ^b Centre for Inflammation Research, The University of Edinburgh, Queen's Medical Research Institute, Edinburgh, United Kingdom, ^c I3S—Instituto de Investigação e Inovação em Saúde, and Faculty of Medicine of Porto, Porto, Portugal, ^d Instituto de Investigación Sanitaria INCLIVA, Valencia, Spain, ^e Neurorestoration Group, Wolfson Centre for Age Related Diseases, King's College London, London, United Kingdom, ^f Grünenthal GmbH, Aachen, Germany, ^g Nuffield Department of Women's and Reproductive Health, University of Oxford, Oxford, United Kingdom, ^h Pelvic Pain Support Network, Poole, United Kingdom, ⁱ International Painful Bladder Foundation, Naarden, the Netherlands, ^j Endometriosis.org, London, United Kingdom, ^k Actual Analytics, Edinburgh, United Kingdom, ^l School of Informatics, University of Edinburgh, Edinburgh, United Kingdom, ^m Department of Neurophysiology, Centre for Biomedicine and Medical Technology Mannheim, University of Heidelberg, Mannheim, Germany

*Corresponding author. Address: Bayer AG, Research & Development, Pharmaceuticals Lab RH VI, Building S107, 09/R302, 13342 Berlin, Germany. Tel: +49 30 468 19823. E-mail address: bianca.deleo@bayer.com (B. De Leo).

Supplemental digital content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's Web site (www.painjournalonline.com).

PAIN 162 (2021) 2349–2365

Copyright © 2021 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the International Association for the Study of Pain. This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

<http://dx.doi.org/10.1097/j.pain.0000000000002248>

disorder targeted for evaluation. Depending on the primary goal, animal models might replicate a disease of interest to different degrees. On the one hand, some animal models focus on further understanding basic science by looking into specific aspects of the disease; in these cases, they may resemble the disease alone partially. On the other hand, when models are designed to be used for the development of new therapies, they need to replicate both aetiology, disease mechanisms (construct validity), and disease symptoms (face validity).

Endometriosis (ENDO) is a chronic, hormone-dependent, peripheral neuroinflammatory condition associated with devastating pelvic pain.⁵⁶ The defining hallmark of the disorder is the presence of tissue fragments (lesions), similar to the endometrium, outside the uterine cavity, rather than in the centre of the uterus (eutopic endometrium). Three major subtypes of ENDO have been defined based on lesion locations: superficial peritoneal, ovarian (endometrioma/cysts), and deep ENDO. There are no effective biomarkers for ENDO, and definitive diagnosis is based on imaging (deep, ovarian cyst) or laparoscopic visualisation of the pelvis. Endometriosis is estimated to affect 6% to 10% of women of reproductive age⁴⁴ and up to 50% of infertile women.⁹⁵ The natural history of the disease remains poorly understood although genetic, immunological, and hormonal factors are all believed to contribute to disease severity.⁴⁴ Notably, although research has identified changes in the central nervous system of patients with ENDO, investigators report a poor correlation between pain severity and the amount, location, and subtype of endometriotic lesions, with some affected patients even being asymptomatic.³² Currently, there is no cure for ENDO, and it is difficult to treat, with most patients offered surgery or drugs that symptomatically and inadequately treat pain or suppress oestrogen action/production causing temporary chemical castration. There are high recurrence rates after surgery, with estimates that lesions return in 40% to 50% of patients within 5 years.⁴⁹ Medical treatments are highly variable in their efficacy and acceptability,¹³³ with many patients having unwanted side effects including suppression of fertility and “menopausal” symptoms. Surveys of patients and health care professionals have consistently cited the need for development of new, effective, and noncontraceptive drugs to successfully control chronic pain symptoms.^{14,57}

Interstitial cystitis/bladder pain syndrome (IC/BPS) is a debilitating condition of unknown aetiology and without effective treatment. Interstitial cystitis/bladder pain syndrome is a symptom-based diagnosis requiring exclusion of other diseases and may have several different causes. A global consensus definition does not exist.⁸⁸ However, 2 distinct subtypes of patients have been distinguished: those with and those without Hunner lesions, which are diagnosed on the basis of cystoscopy. Management depends on the patient's subtype but is always symptomatic.⁸⁴ Recent epidemiological surveys estimated the prevalence of IC/BPS at around 45/100,000 in women and 8/100,000 in men.¹⁰³ To establish a diagnosis of IC/BPS, patients have to present chronic bladder pain and urinary symptoms such as urgency and frequency.⁸⁷ In addition, IC/BPS can be associated with a broad spectrum of comorbidities. These variations combined with the lack of knowledge concerning pathophysiology may explain why existing treatments have low efficacy.

Chronic pelvic pain (CPP)/urogenital pain is defined as “pain symptoms perceived to originate from pelvic organs/structures typically lasting more than 6 months. It is often associated with negative cognitive, behavioral, sexual, and emotional consequences as well as with symptoms suggestive of lower urinary

tract, sexual, bowel, pelvic floor, myofascial, or gynecological dysfunction.”⁵ Examples of conditions associated with CPP are irritable bowel syndrome, vulvodynia, prostatitis, ENDO, and IC/BPS; all of them with a high medical need for new, effective treatments.⁵

In ENDO and IC/BPS, particularly, it has been reported that these conditions can also occur simultaneously.²⁵ Furthermore, ENDO patients have approximately a 4 times higher risk of developing IC/BPS compared with non-ENDO controls.¹⁴¹ In addition, among patients with suspected IC/BPS, when subjected to cystoscopy, patients comorbid with ENDO have a stronger correlation with typical IC/BPS findings such as Hunner lesions, and reduced functional bladder capacity, compared with patients without ENDO.⁵⁴ All in all, because each condition individually has a strong impact on pelvic pain,^{41,61,123} the quality of life of patients with both conditions, or with CPP, is consequently markedly reduced.^{17,117,121,128}

Encouraging reports of progress towards new therapies based on preclinical research are tempered by relatively low success rates in subsequent clinical trials. A major factor for this discrepancy may relate to reproducibility, reliability, and robustness of the animal models used and whether they adequately reflect the full spectrum of symptoms and disease pathophysiology in patients,⁵⁰ especially regarding pain.¹⁶ In this review, we have addressed this issue by considering evidence of the endpoints that have been explored in preclinical models of 2 different pelvic pain disorders: ENDO and IC/BPS. Rigorous evaluation of these endpoints needs to be considered to improve translational research for these disorders. Reviews highlighting the latest advances using preclinical models for each of these 2 diseases have been performed and discussed previously.^{11,119} It is not the intention of this review to replicate findings, but rather to analyse, quantify, and compare the methodologies applied in the development of preclinical models and the outcome measures reported for each model type, with a critical view on parameters that may be of relevance in the clinic. We believe that by rigorously analysing these data, we can contribute to the development of improved preclinical models for both ENDO and IC/BPS, one of the main goals in our Innovative Medicines Initiative (IMI)-PainCare project, subproject Translational Research in Pelvic Pain (TRIIPP) (<https://www.imi-paincare.eu/PROJECT/TRIIPP/>).

2. Methods

2.1. Literature search

Two independent literature searches were performed using the advanced search engine from the Embase database (<https://www.embase.com/#advancedSearch/default>). For the first search, the terms “endometriosis” and “model” were used. For the second search, the terms “interstitial cystitis” or “bladder pain syndrome” were used. The advanced search settings included the following mapping options: “map to preferred term in Emtree,” “search also as free text in all fields,” “explode using narrower Emtree terms,” and “search as broadly as possible.” The first 2 searches were limited to show results published in the last 20 years (between 1998 and May 2019). Subsequently, 2 new searches with the same search criteria were performed in June 2020 covering all publications that arose between May 2019 and June 2020.

Overall, searches for ENDO and IC/BPS led to 3309 and 7266 results, respectively, which were exported into RIS (research information system) file format and opened in the software EndNote X7 (Clarivate, Philadelphia, PA) for further analyses.

Data from the search results were further automatically and manually analysed for systematic article selection and subcategory of preclinical models (Fig. 1). A detailed description of

article selection, definition of subcategories, and analysis of construct and face validity can be found in the supplementary data (available at <http://links.lww.com/PAIN/B319>).




Figure 1. Selection of relevant research articles using endometriosis and IC/BPS preclinical models. Articles from the Embase database were imported into the software EndNote and classified. The order of classification was defined as shown in the flow chart: first, "reviews and editorials," then "clinical studies," then "in vitro/ex vivo/in silico studies," then "not rodent preclinical models," and finally "not endometriosis" or "not IC/BPS models." To avoid duplication of data, all poster presentations that had led to publications were excluded from the analysis, as well as poster presentations of the same data published more than once. Articles remaining were categorized as "ENDO preclinical rodent models" or "IC/BPS preclinical rodent models" and consequently analysed. ENDO, endometriosis; IC/BPS, interstitial cystitis/bladder pain syndrome.

3. Results

3.1. Preclinical models of endometriosis

During the last 2 decades, 4 different transplantation techniques have mainly been used: xenograft transplantation (transplantation of endometriotic tissue of human origin into immunocompromised mice), syngeneic transplantation (transplantation of uterine tissue of murine origin into recipient mice), autologous transplantation (transplantation of uterine tissue (uterine horn/uterus fragments) of rodent origin into the diverse pelvic regions of the same animal), and allogeneic transplantation (transplantation of uterine tissue from one mouse into the same mouse).

In our search, we found that from a total of 1009 scientific publications using preclinical models of ENDO (Fig. 2A) 41% used autologous transplantation, 26% syngeneic transplantation, 18% xenograft transplantation, and 11% used allogeneic transplantation. A minority of publications (3%) did not specify the model used, and very few used other means to induce ENDO, eg, CRISPR/Cas gene editing. We evaluated the number of articles being published per year for each category, and it was apparent that the most commonly used ENDO model was also the first widely established methodology in rodents: publications using autologous transplantation increased every year until 2015, reaching a peak of 43 publications (Fig. 2B). Since then, there has been a slight reduction and a plateau, likely reflecting the adoption of other models as detailed below. Syngeneic transplantation is the second most widely used method, and starting in 2008, there was a rapid increase in the number of publications using this model, peaking in 2019 with 41 publications. The xenograft transplantation model (first described in 1999) has not been as widely used, likely reflecting limitations in access to human tissue/cells and the requirement for immunosuppressed recipients. Although the number of publications increased between 2008 and 2015, reaching a peak of 28 publications, there was a decrease in the following years. The allogeneic transplantation model is the least used model variant. This was first described in 2000 and was increasingly reported from 2010, reaching a peak in 2016 with 13 publications. It is important to note that during the analysis of the search results it became clear that a contributing factor to the high number of publications within each category comes from laboratory groups publishing their discoveries using the same model repeatedly over time (not shown), and that this is also of concern when reproducibility of data becomes an issue.

Furthermore, differences exist between the model subcategories in the methods used to deliver endometriotic/uterine tissue and the location of the tissue implants (Figs. 2C–J). In the autologous transplantation model, suture is the only method used to develop “lesions” (Figs. 2C and D). Considering the location of implanted tissue fragments, 33% of publications reported suture to the “peritoneum,” followed by the “abdominal wall” (26%). Since both refer to the same target tissue area, together they accounted for 59%, ie, the most targeted area in this model category. In addition, 19% of publications located tissue on mesenteric arteries/vessels, 3% on ovaries, and a minority (3% of total, 6 publications) in the colorectal region. Eight percent of publications sutured tissue fragments to other locations (eg, gastrocnemius muscle,⁴ bowel mesothelium,⁹ sciatic nerve,²³ and hypochondriac regions²⁰ or used dorsal skinfold chambers⁷²). Ten percent of publications did not specify the location of implanted tissue fragments.

In the syngeneic transplantation model (Figs. 2E and F), the most common route of delivery was by intraperitoneal injection of homogenized tissue fragments (62% of all publications in this

category), followed by suturing or gluing uterine fragments intraperitoneally (32% of publications), and, finally, a minority of publications reported subcutaneous injection of suspended tissue (less than 2% of publications³⁷). Four percent of publications within this category did not specify either the route of delivery or the location of the implanted tissue.

In the xenograft transplantation model (Fig. 2G and H), the most common route of delivery was intraperitoneal injection of suspended endometriotic tissue (53% of all publications in this category), followed by suturing or gluing endometriotic tissue fragments (24% of publications) and, finally, subcutaneous injection of suspended tissue (15% of publications). Nine percent of publications within this category did not specify either the delivery route or location of the implanted tissue.

In the allogeneic transplantation model (Figs. 2I and J), suturing was the most common method used to secure uterine tissue implants. The “peritoneum” (53%) was the most commonly targeted area, followed by mesenteric arteries/vessels (21%), and the “abdominal wall” (15%). Together, the “peritoneum” and the “abdominal wall” represent 68% of publications of the model category. Eleven percent of all publications did not disclose where tissue fragments were implanted.

3.2. Construct validity for preclinical models of endometriosis for translational research

We developed an algorithm with a 3-point scale to determine the models and delivery methods better suited for translational research in terms of construct validity. We found that the construct validity ranking for ENDO models was as follows (Fig. 3): the highest rank was reached by the syngeneic model performed by intraperitoneal injection of suspended tissue, with a score of 12 points. The second rank, with a score of 8 points, is shared by the allogeneic model and the syngeneic model, both performed by surgery. The third rank, with 6 points, is shared by the xenograft model by intraperitoneal injection and the autologous transplantation model, which is performed by surgery. The fourth rank, with 4 score points, is shared by xenograft performed by surgical implantation and syngeneic transplantation by subcutaneous injection. Finally, in the last rank with 2 points is xenograft transplantation performed by subcutaneous injection of endometriotic tissue. Interestingly, the model subcategory with the highest ranked score in our construct validity scale, syngeneic transplantation by intraperitoneal injection, is also the second overall most used preclinical model of ENDO, with 167 publications, and the most used model per year since 2017.

3.3. Face validity in preclinical models of endometriosis

All in all, the lesion size was the most addressed endpoint parameter in model subcategories of ENDO, being analysed in 53% of publications using the autologous transplantation model, 74% from syngeneic, 77% from xenograft, and 80% from the allogeneic transplantation model (Figs. 4A–D). The second most reported endpoint was inflammation, being measured in 31% of publications using autologous, 40% using syngeneic, 19% using xenograft, and 47% using allogeneic transplantation models. Although adhesions occur in a high proportion of patients, a specific evaluation of adhesions in the preclinical models was rarely evaluated (12%, 8%, 5%, and 14% of publications from each model subcategory as defined above).

Notably, one of the key clinical symptoms reported by those with ENDO is CPP, and this has not been rigorously and reproducibly analysed in preclinical models of ENDO. As pain




Figure 2. Preclinical models of endometriosis. (A) Four main categories of preclinical models of endometriosis were found. Blue fraction: autologous transplantation, turquoise fraction: syngeneic transplantation, green fraction: xenograft transplantation, and brown fraction: allogeneic transplantation. (B) Number of publications of the model categories published between 1998 and 2019. (C) Within the autologous transplantation model, tissue fragments may be implanted in different areas. Dark blue fraction: peritoneum, blue fraction: abdominal wall, cyan fraction: mesenteric arteries/vessels (A/V), light blue fraction: ovary, white fraction: colorectal region, lilac fraction: other regions, and gray fraction: not specified. (D) Scientific articles published per year using the autologous transplantation model between 1998 and 2019. (E) Variants of the syngeneic transplantation model. Dark turquoise fraction: publications using intraperitoneal injection of suspended tissue and turquoise fraction: publications using suture. Light turquoise fraction: publications using subcutaneous injection of suspended tissue. Gray fraction: articles where it was not specified. (F) Publications per year using the syngeneic transplantation model between 1998 and 2019. (G) Variants of the xenograft transplantation model. Dark green fraction: intraperitoneal injection of suspended tissue. Green fraction: suture of endometriotic tissue. Light green fraction: subcutaneous injection of suspended tissue. (H) Publications per year using the xenograft transplantation model between 1998 and 2019. (I) Allogeneic transplantation model variants. Brown fraction: peritoneum. Mustard fraction: mesenteric arteries/vessels (A/V). Light brown fraction: abdominal wall. Dark yellow: not specified. Gray: subcutaneous injection. (J) Publications per year using the allogeneic transplantation model between 1998 and 2019. N/S, not specified.




Figure 3. Construct validity for translational research of preclinical models of endometriosis. Based on the score results from our 3-point scale algorithm specifically developed to determine the construct validity for translational research purposes. The size of the dots represent the number of publications within the model, which are indicated on top of each dot as well. Colour of the dots indicates the model category; blue: autologous transplantation, green: xenograft transplantation, turquoise: syngeneic transplantation, and brown: allogeneic transplantation. The endometriosis model with the highest construct validity was syngeneic transplantation of endometriotic tissue by intraperitoneal injection of suspended tissue. AL-S, allogeneic transplantation models, suture; AT-S, autologous transplantation model, suture; SY-IP, syngeneic transplantation model performed by intraperitoneal injection; SY-S, syngeneic transplantation model performed by suture; SY-SC, syngeneic transplantation model performed by subcutaneous injection; XE-IP, xenograft transplantation model performed by intraperitoneal injection; XE-S, xenograft transplantation model performed by suture; XE-SC, xenograft transplantation model performed by subcutaneous injection.

perception cannot be reported by rodents, surrogate measures based on altered behaviours and/or altered response to stimuli are used.^{125,126} The model subcategory having the biggest proportion of publications addressing pain is autologous transplantation with 12%, followed by syngeneic 10%, allogeneic 6%, and xenograft transplantation with 0.5%. Within the autologous transplantation model, for example, a model of viscerovisceral hyperalgesia was developed. In this model, female rats with previously induced ENDO were also subjected to ureteral calculosis. Spontaneous pain behaviour was observed for several days in rats having ENDO and stone implantation, in contrast to ENDO alone, or sham ENDO controls.^{43,60,82} This model provides better insights into preclinical models regarding the occurrence of viscerovisceral hyperalgesia that also occurs in ENDO patients. However, it is unclear whether or not the reported observations may be due to the surgical induction of ureteral calculosis. From all publications studying nociception and pain, 87% used evoked-pain measures. By contrast, only 17% of publications measured nonevoked pain readouts. Within subcategories, the syngeneic transplantation model variant had the highest number of publications studying nonevoked pain: 11 publications accounting for 42% of the publications addressing pain in this subcategory.^{13,36,47,73,74,106} Sub/fertility, another common symptom reported by women with ENDO, was also surprisingly understudied, accounting for up to only 1%–6% of publications from all model subcategories.

3.4. Preclinical models of interstitial cystitis/bladder pain syndrome

In the last 22 years, the majority of preclinical models of IC/BPS were bladder-centric models, comprising 894 publications and representing 85% of the total. This was followed by models with complex mechanisms, which with 86 publications comprise 8% of total. Finally, the category of stress-induced models with 69 publications accounted for 7% of total (**Fig. 5A**). Because IC/BPS

is a disorder occurring in both men and women, these results include publications with both male and female rodents.

For each IC/BPS model category, the number of articles published per year was analysed (**Fig. 5B**). The largest model category, bladder-centric, was also the first methodology to be established, and publications using this model increased almost continuously every year until 2016, reaching a peak of 71 publications, followed by a plateau and subsequent decrease in 2019 with only 33 publications (**Fig. 5B**). In the bladder-centric category, 3 methods have been established: instillation of irritants (92% of publications), altered expression of urothelial targets (5% of publications), and radiation-induced cystitis (3% of publications, **Figs. 5C and D**). Instillation of irritants consists of infusing the bladder with an agent that damages the urothelium, generally achieved by transurethral catheterization under anaesthesia. Irritants commonly used are as follows: acetic acid,⁶⁶ protamine sulphate,³⁸ and lipopolysaccharide.⁸³ This category also includes the cyclophosphamide model which, in contrast to the other irritants, is administered systemically. This model variant accounts for up to 48% of all publications within the category (426 publications, **Figs. 5C and D**). Within the subcategory “altered expression of urothelial targets,” different techniques have been used aiming to understand the role of gene products in disease development, including gene therapy, transgenic mice, siRNA, and exogenous administration of specific peptides or protein inhibitors.^{22,29,64,68,98} The subcategory “radiation” consists of specifically irradiating the bladder one or more times, producing local tissue damage, and triggering haemorrhagic cystitis.¹³⁶

Within the “complex mechanisms” category, 5 different subcategories to induce bladder complications have been identified (**Figs. 5E and F**): autoimmune model, colonic instillation of 2,4,6-trinitrobenzenesulfonic acid (TNBS), injection of pseudorabies virus in the tail, pelvic pain comorbidities, and others. The autoimmune model accounted for 43% of publications within its category, with that being the biggest subcategory, followed by colonic instillation of TNBS. Colonic instillation of TNBS induces local inflammation and consequently bladder complications which are generally measured between 3 and 30 days after instillation.^{63,85,130} This subcategory accounted for 20% of publications in the category. Another model variant consists of injecting pseudorabies virus in the tail of the animals, inducing neurogenic cystitis characterized by progressive pelvic pain. The outcomes from model induction are generally measured 2 to 5 days after injection.^{21,113,142} This subcategory accounted for 14% of publications in the category. The next subcategory, pelvic pain comorbidities, includes experimental procedures in which other pelvic areas were impaired, development of bladder complications, eg, prostatic inflammation,^{97,116} constipation,⁵⁸ or uterine pain.⁷⁶ Bladder complications are measured between 7 days and 4 weeks after model induction. This subcategory accounted for 8% of publications. Finally, all other complex mechanisms that could not be related to any of the former subcategories were grouped together, eg, intake of tranilast, an antiallergy agent that has been reported to lead occasionally to IC/BPS in patients⁹⁹; and systemic administration of peptides/proteins that modulate the sensory nervous system and/or brain function.^{18,89,118} Together, this subcategory accounted for 15% of publications. In addition, publications using these techniques have shown a cyclic pattern with 3 different peaks of 6, 7, and 9 publications in the years 2007, 2012, and 2016, respectively, followed by a plateau and decrease in the respective following years (**Fig. 5F**).

Finally, stress-associated IC/BPS models are the newest and least used models, being first described in 1999 and becoming




Figure 4. Face validity of preclinical models of endometriosis. Analysis of the outcomes measured in the publications within each model category. (A) Autologous transplantation model. (B) Syngeneic transplantation model. (C) Xenograft transplantation model. (D) Allogeneic transplantation model. Dark blue bar: percentage of publications that measured macroscopic changes in either lesion size, volume, weight, or number. Blue bar: percentage of publications that measured of inflammatory markers. Light blue bar: percentage of publications that quantified adhesions. Purple bar: percentage of articles that measured fertility. Red bar: percentage of publications that assessed nociception/pain aspects. Coral bar: percentage of publications that assessed evoked-pain readouts. Green bar: percentage of publications that assessed nonevoked pain readouts. N/P, nociception/pain.

increasingly used as from 2006. The peak was reached in 2013 with 10 publications (Figs. 5G and H). Within the stress-induced model category, 5 different methods have been identified: water avoidance stress, manipulation of the environment, footshock stress, neonatal maternal separation, and restraint stress. The most published model variant is water avoidance stress, which consists of subjecting the animals to either acute (1 day) or chronic (10 days) stress by placing the animals for 1 hour on a platform inside a container full of water. Generally, bladder complications are measured one day after the last day of the test.^{91,107} This subcategory accounted for 39% of publications, followed by manipulation of the environment. In the latter, the model variant may be induced by chronic variable stress,⁵¹ early life stress by odour-shock conditioning,⁷⁸ cold stress,⁵⁹ or social stress.⁹⁶ Outcomes were measured in adulthood and vary depending on the stress protocol. This subcategory accounted for 32% of publications within stress-induced model subcategories. Next, the subcategory footshock stress accounted for 13% of publications and consisted of performing an electric foot shock of 1 mA for 1 second, the animals afterwards being subjected to functional experiments.¹¹² The subcategory restraint stress accounted for 10% of publications within stress-induced models and consisted of immobilizing the animals in a restrainer for 3 to 4 hours^{19,35} before measuring outcomes in bladder morphology and/or function. Finally, the subcategory neonatal maternal separation consisted of separating animal litters from their mother on postnatal days 1-5 to 21-25 daily for 3 hours. Subsequently, bladder complications were measured

when animals reached 1 to 5 months of age.^{90,108} The number of publications reporting this model accounted for 7% of publications.

3.5. Face validity in preclinical models of interstitial cystitis/bladder pain syndrome

In IC/BPS patients, several clinical features can be observed: bladder pain, micturition frequency, nocturia, perineal pain, glomerulations (although they are not IC/BPS specific), Hunner lesions, and urothelial denuded areas. These aspects have been mimicked partly in preclinical models of IC/BPS. In this study, we have categorized them as follows: “bladder morphology,” “bladder function,” “behaviour,” and “nociception/pain”; the latter being further classified as “evoked” and/or “nonevoked,” and finally “others” (Fig. 6).

The endpoint “bladder morphology” indicates that the microscopic bladder structure has been analysed. This allows evaluation of the degree to which the clinical hallmarks “glomerulations” and “Hunner lesions” are present in the animal model by detecting inflammatory markers in bladder tissue and areas of urothelial denudation. Likewise, with the endpoint “nociception/pain,” it is possible to evaluate to what degree bladder and perineal pain are seen in the animal models. Finally, with the endpoint “bladder function,” both nocturia and micturition frequency can be evaluated.

The most addressed endpoint was bladder function, being analysed in 55% of publications using bladder-centric models




Figure 5. Preclinical models of IC/BPS. (A) Preclinical models of IC/BPS. Three main categories were found. Dark purple fraction: bladder-centric models. Green fraction: complex mechanisms models. Orange fraction: stress-induced models of IC/BPS. (B) Number of publications of the model categories published per year during 1998 to 2019. (C) Variants of the bladder-centric model. Dark purple and fuchsia fractions: instillation of irritants. In fuchsia, the use of cyclophosphamide (CYP) is highlighted. Purple fraction: models with altered expression and/or function of urothelial targets. Light purple fraction: radiation-induced cystitis. (D) Publications per year using bladder-centric model variants between 1998 and 2019. (E) Variants of the complex mechanisms model. Dark emerald fraction: autoimmune models. Emerald fraction: colonic instillation of TNBS. Light emerald fraction: injection of PRV in tail. Mint fraction: chronic pelvic pain (CPP) comorbidities. Light mint fraction: others. (F) Publications per year using complex mechanisms models between 1998 and 2019. (G) Variants of the stress-induced model. Dark orange fraction: WAS. Orange fraction: manipulation of the environment. Light orange fraction: footshock stress. Sand fraction: Neonatal maternal separation (NMS). Light sand fraction: restraint stress. (H) Publications per year using stress-induced models between 1998 and 2019. IC/BPS, interstitial cystitis/bladder pain syndrome, PRV, pseudorabies virus; TNBS, 2,4,6-trinitrobenzenesulfonic acid; WAS, water avoidance stress.




Figure 6. Face validity of preclinical models of IC/BPS. Analysis of the outcomes measured in the publications within each IC/BPS model category. (A) Bladder-centric models. (B) Complex mechanisms models. (C) Stress-induced models. Blue bar: percentage of publications that assessed bladder function by cystometry, voiding spot, or metabolic cages. Light blue bar: percentage of publications that assessed bladder morphology by histopathology, bladder weight, or immunohistochemistry. Red bar: percentage of publications that assessed nociception/pain endpoints. Coral bar: percentage of publications that assessed evoked-pain readouts, such as abdominal von Frey or VMR. Green bar: percentage of publications that assessed nonevoked pain readouts, such as observing pain-related behaviours among individuals during a definite time frame. Purple bar: percentage of publications that assessed other endpoints such as changes in expression of different markers in tissues other than the bladder. IC/BPS, interstitial cystitis/bladder pain syndrome; N/P, nociception/pain; VMR, vasomotor reflex.

(Figs. 6A), 56% of publications using complex mechanisms models (Figs. 6B), and 74% of publications using stress-induced models (Fig. 6C). The second most evaluated endpoint was bladder morphology, which was evaluated in 50% of publications using bladder-centric models, 50% of publications using

complex mechanisms models, and 49% of publications using stress-induced models. Endpoints belonging to the category “others”—including gene expression/function in tissues other than the bladder or nonstructural gene expression analyses—were evaluated among categories almost in the same proportion, varying between 14% and 20% of total publications within each subcategory.

Nociception and pain endpoints were evaluated in a subset of publications investigating preclinical models of IC/BPS. The model subcategory most covering these outcomes is the stress-induced model subcategory with 44% of total publications, followed by complex mechanisms with 43% of publications, and finally bladder-centric models with 20% of publications. In other words, the largest model subcategory is also the one with the smallest proportion of studies including nociception and/or pain endpoints and vice-versa. However, of all publications studying nociception and pain within IC/BPS model categories, 91% used evoked-pain measures, and only 14% measured nonevoked pain readouts. Within subcategories, “bladder centric” was the model variant with the highest number and proportion of publications studying nonevoked pain: 32 publications accounting for 13% of the publications addressing pain and 4% of the total publications in this subcategory.

4. Discussion

Recently, the World Health Organisation recognised chronic pain as a disease in its own right. Within this framework, ENDO and IC/BPS are categorised as “chronic visceral pain from persistent inflammation in the pelvic region,” as described in the International Classification of Diseases 11th.⁶ The causes of both diseases remain unknown; there is no cure, and pain frequently persists and may worsen.

In this comprehensive literature review of all publications in the last 22 years using preclinical models of ENDO and IC/BPS in rodents, we have analysed which translational features have been studied more or less frequently to uncover aspects of high relevance to patients who have not, or been insufficiently, evaluated and need to be addressed to improve the development of new therapies for both conditions. In summary, our results indicate:

(1) The most frequently used preclinical model of ENDO is autologous transplantation, followed by syngeneic transplantation. Although the former seems to be declining in use, the latter has been increasingly used across the years. By contrast, the most frequently used preclinical models of IC/BPS were bladder centric. However, the number of publications in the last 3 years for bladder-centric models has been on the decline, whereas other model categories, such as complex mechanisms and stress-induced, are gaining traction.

(2) By adopting a 3-point scale to evaluate the construct validity of preclinical models of ENDO in the published literature, we found that syngeneic transplantation of endometriotic tissue, specifically performed by intraperitoneal injection, resembles the physiopathology of ENDO in a more reliable manner than the other subcategories. Unfortunately, with the current lack of knowledge of disease aetiology, it was not possible to develop a construct validity scale for preclinical models of IC/BPS as it remains unclear with which pathomechanisms the models need to be compared.

(3) In both pelvic pain disorders, the face validity has mostly been restricted to demonstrating tissue damage and organ dysfunction, despite the fact that patients suffer from CPP and other disease-associated traits (in ENDO: fertility issues, in IC/

BPS: urinary urgency). In addition, when assessing pain, preclinical models mostly used evoked pain measures rather than measuring ongoing pain. However, the latter is a significant element of the clinical symptoms of both disorders.

4.1. Rationale for the construct validity scale for translational research

Although all preclinical models of ENDO have been helpful in improving our understanding of lesion formation and disease progression, they also have a number of limitations when considering their relevance for translation to patients. We developed a scoring system for preclinical models of ENDO specifically designed to highlight which models are more suitable for translational research purposes. The scoring system is based on 3 criteria: first and foremost, it attempts to evaluate how closely each model mimics the pathomechanism of ENDO. Then, it aims to evaluate in each model to what degree the immunoinflammatory components of the animal model can be studied. Finally, it aims to evaluate which models have the ability to study the function of specific gene products.

Surgical attachment of endometriotic tissue fragments is routinely performed in all transplantation model subcategories. This technique offers the advantage of choosing the areas for disease establishment and progression. Examples of such areas include the ovaries, mesenteric arteries/vessels, or colorectal region.^{42,52,143} However, surgical implantation is a highly invasive procedure that encompasses administering general anaesthesia, opening and exposing tissue layers in the pelvic area, and suturing tissue implants. This procedure unavoidably induces postoperative pelvic pain, as well as a risk of infection. Nevertheless, when comparing allogeneic vs autologous transplantation, the advantage of autologous transplantation, which is usually performed in rats, rather than allogeneic transplantation, which is performed in mice, is that the size of the animal allows for surgery, that is, relatively easy to perform. In the allogeneic transplantation model, because mice are smaller, performing the surgery can be more challenging although the availability of large numbers of transgenic lines of mice opens up the possibility of studying the role of specific genes and cell types in the development of the disease.^{34,93}

In contrast to surgical procedures, injection of suspended tissue provides a considerably less invasive way of delivering endometriotic tissue without causing external or internal tissue injury. Specifically, intraperitoneal injection resembles more accurately what would occur during retrograde menstruation in humans. Retrograde menstruation is the physiological process involving the backward flux of menstrual debris containing viable endometrial cells through the fallopian tubes into the pelvic cavity. This process occurs naturally in over 90% of women of reproductive age with patent fallopian tubes and provides the material to potentially develop ENDO.¹⁴⁵ However, because only a fraction of patients with retrograde menstruation will develop ENDO, additional factors must be involved in the establishment and development of lesions that underpin disease progression. These factors likely include endocrine and metabolic (pelvic) environment, epithelial–mesenchymal transition, and altered immunity and inflammatory responses in those genetically susceptible.¹⁴⁵

Comparing xenograft and syngeneic transplantation models, the 2 models that have used intraperitoneal injection of suspended tissue, xenograft transplantation models are developed by transplanting human endometriotic tissue or cells directly from patients, bringing with them all the disease factors

that are actually present within them. The disadvantage of this model is that for mice to receive tissue of human origin, immunosuppression is required. Because ENDO is considered to be an estrogen-dependent inflammatory disorder, the contribution of inflammatory components of the recipient mice can only partially be studied. Nude mice have deficits in their reproductive system, including reduced circulating hormone levels.¹¹¹ Furthermore, because of the lack of T lymphocytes and/or B lymphocytes, depending on the phenotype, they are more susceptible to pathogens, and special care must be taken when performing behavioural studies. Most strikingly, in a model of colorectal pain, nude mice have shown abnormal pain sensation which is normalized after supplementation with T cells, demonstrating the link between T lymphocytes and visceral pain sensitivity.¹³⁵

4.2. Technical considerations on the syngeneic transplantation model performed by intraperitoneal injection model variant

Between research groups performing this model variant, differences in the methodology are apparent. For example, after extracting and splitting the uterus horn from donor mice, some groups report cutting the tissue fragments or mincing the uterine tissue to obtain tissue fragments,^{2,77,81,120,124,139} whereas others first remove the serosa layer and myometrium and use the remaining tissue as a source of tissue fragments for injection,^{7,33,46,53,55,104,114} and others have reported squeezing the uterine horn to obtain endometriotic tissue.²⁷ These differences in experimental protocols might seem slight but, in fact, provide recipient mice with different amounts and ratios of cell types. Another example of variation is to regulate the hormonal cycle of the animals and to induce menstruation in mice. These protocols include one or more of the following procedures: performing ovariectomy in mice to suppress the endogenous hormone activity, administering donor mice with oestrogen (17 β -oestradiol) for a period of time (usually 7 days), administering progesterone in donor mice to induce decidualisation before tissue harvest, as well as supplementing recipient mice with oestrogen before and/or after intraperitoneal injection of endometriotic tissue fragments. These artificial hormonal interventions in rodents might have positive effects in construct validity, but only when carefully calibrated to replicate menstruation as well as the hormone fluctuations that occur in ENDO patients. It is important to note that because rodents do not menstruate naturally, intraperitoneal injection of suspended tissue fragments from cycling mice may therefore introduce a different mix of cells to those introduced by retrograde menstruation in women.

In this regard, a new variant of the syngeneic transplantation was established in 2014 in which menstrual tissue was generated in and collected from donor mice and injected intraperitoneally in recipient mice.⁴⁶ Generation of menstrual tissue for use with the syngeneic mouse model of ENDO involves the use of donor mice which undergo a protocol of artificial decidualisation and progesterone withdrawal to induce menstruation.⁴⁶ Specifically, donor mice are ovariectomised to deplete endogenous steroid hormones and then treated with a hormone regime that mimics the normal menstrual cycle of women.⁴⁶ Decidualisation occurs by intrauterine oil administration which simulates blastocyst attachment followed by withdrawal of progesterone, which results in endometrial breakdown.⁴⁶ The “menstrual” tissue is injected into recipient mice, and progression of ENDO can be monitored. It is important to note that although this model resembles the physiological processes believed to be involved in

lesion formation as a consequence of retrograde menstruation, it also entails the added complexity of inducing artificial menstruation in donor mice, which needs to be taken into account. Both the frequency and the extent of decidualisation in donor mice can vary between experimental runs, with anecdotal evidence suggesting that stress levels of mice may have an impact on decidualisation. In addition, the decidualised endometrium is minced and injected into recipient mice, with no control over which tissue fragments will attach to form lesions, meaning that the cellular composition of the tissue fragments which will end up forming lesions can vary between animals, attachment sites, and experimental runs. Although this can be a drawback regarding controlled reproducibility, it does better resemble the lesion heterogeneity observed in those with ENDO. Interestingly, the established lesions in these mice demonstrated several hallmarks of the human disease, including lesion components, changes in oestrogen receptor expression, as well as macrophage infiltration and inflammation. Along the same lines, a new model of retrograde menstruation in rats was recently developed,¹⁰⁵ with promising insights into behavioural assessments of ongoing pain.

We recommend careful revision and discussion of these protocols, aiming to reach a consensus within the scientific community to improve understanding of the range of physiological parameters involved in these models. We hope to obtain greater insights into this regard in our IMI-PainCare project through controlled comparison of experimental protocols and performing multicentre staff training towards validated methods and comparable standard operating procedures.

4.3. Construct validity scale for preclinical models of interstitial cystitis/bladder pain syndrome

Interstitial cystitis/bladder pain syndrome is a difficult condition to diagnose, and there is limited understanding of the disease mechanism. Moreover, global consensus on definitions and treatment possibilities has not yet been achieved, and standard terminology is constantly evolving.¹³⁸ Based on this, it was not possible to establish a construct validity scale for IC/BPS. Given that the patient population is very heterogeneous in terms of detectable pathological changes, as well as local extension of the disease and comorbidities, we recommend that research focuses its efforts on deep patient phenotyping procedures. Patient phenotyping and subgrouping are likely to be essential to unlock major advances in the state of the art. Patients report a variety of different symptoms for both pain and urinary symptoms: In some patients, symptoms extend beyond the bladder, and in some, the disease is accompanied by different comorbidities. The multiplicity of symptoms and signs will require the use of different and appropriate animal models to progress translational studies of IC/BPS. In addition to the pain or the micturition phenotype, the molecular signature obtained from the analyses of urine, blood, or saliva samples of the patients will help to choose more appropriate animal models to address the putative diverse pathophysiological mechanisms of IC/BPS, pointing us towards to effective diagnostic tools and treatments.

Most of the bladder-centric models have limitations in reflecting the chronicity and comorbidities of IC/BPS. For example, models in the subcategory “instillation of irritants” generally induce an acute, short-term effect, and therefore, these experiments are often performed terminally. So far, there is no cure for IC/BPS,³⁰ which means that patients have to learn to live with the symptoms throughout their life, often with a tendency to accumulate comorbidities with time.¹¹⁰ It will be worth refining the preclinical models to obtain more insights into these aspects, eg,

by modulating the strength and duration of a trigger to induce a potentially milder but chronic disease state. A surprisingly high proportion of publications within the instillation of irritants subcategory used the cyclophosphamide model (48% of publications). Cyclophosphamide, which was originally administered in cancer patients as a chemotherapeutic agent, is metabolised into acrolein/ifosfamide, which accumulates in the bladder and in turn induces irritation of the urothelium, inflammation, and pain,^{31, 71} whereas it may also irritate many other tissues. In contrast with the rest of the irritants in the model category, this model variant can be administered systemically (generally injected intraperitoneally), which might explain why it has been preferred over the classic transurethral catheterization under anaesthesia commonly used for the rest of the irritants in the subcategory. In 2014, a chronic cyclophosphamide model was characterized,⁴⁵ and these mice, after repeated systemic injections of cyclophosphamide, have shown signs of mild inflammation, urothelium hyperplasia, and local changes in metabotropic glutamate receptor expression. However, these traits do not entirely represent what occurs in the bladder of IC/BPS patients, where fibrosis rather than hyperplasia may be observed in the urothelium of some patients,^{12, 65} and it is not clear to what extent these local changes in expression of glutamate receptors play a role in IC/BPS patients. Other traits that may occur in the cyclophosphamide model, such as increase in urinary frequency and/or urgency, can also be attributed to haemorrhagic cystitis, rather than IC/BPS. Therefore, the chronic cyclophosphamide administration model has limitations for translational research on IC/BPS.

A new bladder-centric model has been proposed as an alternative to normal irritants. This model is based on the breakdown of the urothelium's apical cell layer because of disruption of the glycosaminoglycan layer by enzymatic digestion.¹⁰⁰ The authors have presented 2 protocols of enzymatic deglycosilation: acute and chronic. They have found that after 2 hours of enzymatic digestion of the extracellular matrix of the urothelium, the mice replicate various pathological phenotypes found in patients as follows: disruption of the mucosal barrier with loss of the urothelial cell plaques, macrophage infiltration, increased urinary frequency and urgency, suprapubic hyperalgesia, and increased nociceptive activity.¹⁰⁰ A potential limitation of this model is that the effects induced by enzymatic digestion are reversed 7 days after model induction because of tissue repair which unfortunately does not occur in IC/BPS patients.

Acknowledgment of the limitations of the bladder-centric models in comparison with the complex clinical situation has probably driven the emergence of the complex mechanism models as well as the stress-induced models, as they apply stimuli beyond the bladder and thus shift the focus from the bladder to a more systemic origin of the disease, making it impossible to reliably score construct validity. Autoimmune models acknowledge the fact that in a significant number of patients IC/BPS occurs in association with autoimmune diseases such as Sjögren syndrome, rheumatoid arthritis, lupus erythematosus, and others.^{11, 79, 131} In this model variant, an autoimmune response is triggered in the animals by injecting immunogen agents and/or homogenate of bladder tissue combined with complete Freund's adjuvant. Bladder function and morphology are assessed between 10 days and 4 months after induction.^{10, 62, 144} The bladders from these animals show changes in both bladder tissue composition as well as bladder function, compared with control animals.⁷⁹

Inside the complex mechanism category, chronicity of the models varies greatly, lasting up to 16 weeks,¹²² with most of the models measuring their endpoints 1 to 12 weeks after model induction.^{3,62,67,79,80} In the same way, within the stress-induced model category, chronicity varies between 8 and 12 days^{1,40,91,137} and up to 8 weeks of age.¹⁰⁸ These long-term effects on the animals after model induction could be advantageous in construct validity for translational research; nevertheless, evaluation has to distinguish between acute and chronic effects, and those outcomes should then be carefully compared with patient outcomes. Nevertheless, these model categories might reflect the clinical situation only partially, and ultimately, it is unclear whether the pathophysiological alterations induced artificially in the animal models mimic the origin of the human disease. Moreover, the clinical situation does not easily allow separation of origin and consequence. Although there is evidence that traumatic experiences may be part of the disease history/origin in some patients,^{24,28,92} it is known that stress and psychological perturbations also occur as a consequence of a disease which has a significant psychosocial impact on daily life, may reduce quality of life, impair sleep, and lead to depression.^{70,132}

4.4. Face validity of preclinical models of endometriosis and interstitial cystitis/bladder pain syndrome

In over 2 decades of research using preclinical models of ENDO, most scientific articles (between 53% and 80%, depending on the model subcategory) measured the lesion size in their animal models as a primary endpoint. This endpoint has been used both as a proof-of-principle as well as to evaluate the effectiveness of a potential treatment. By contrast, only 0.5% to 12% of these scientific publications measured pain outcomes, and only 1% to 6% have measured fertility. This is a striking observation, considering that the most common symptoms of ENDO in patients are pain and infertility.¹⁵ Moreover, the lesion size does not correlate with pain levels and fertility.^{127,134} In IC/BPS, the most measured aspects in preclinical models were bladder function and bladder morphology, whereas measures of nociception and/or pain were only included in 20% to 44% of publications. Most strikingly, nonevoked pain outcomes comprised 0% to 4% of publications within the categories. Therefore, we highly recommend that the scientific community pays more attention to these aspects when pursuing development of new medicines, especially regarding pain-related outcomes.

Pain symptomatology in patients suffering from these chronic pain conditions is diverse: In ENDO, up to 80% of patients suffer from cyclic dysmenorrhea (pain during menstruation), up to 50% suffer from CPP, 40% to 50% suffer from deep dyspareunia (pain during intercourse), and a small percentage of patients (1%-2%) suffer from dysuria (pain during urination), and/or dyschezia (rectal pain during bowel motion or in some patients because of rectal bleeding during menstruation).^{15,115} In IC/BPS, the most severe pain symptoms are bladder pain and overall pain, pelvic pain, abdominal pain, bloating pain, dysmenorrhea, back pain, pain during bladder filling, and dyspareunia.²⁶ With sustained pain over time, central sensitization is also a phenomenon occurring in patients with ENDO⁴⁸ and IC/BPS.^{69,102} Both somatic and visceral pain should be studied in detail in preclinical models of ENDO and IC/BPS to refine them for improved face validity. Importantly, neurogenic inflammation should also be considered as a possible mechanism in the development of both conditions. It has been proposed that retrograde menstruation is the first step in the development of ENDO-associated pain.⁷⁵ In

this theory, ENDO-associated pain (EAP) would occur through local release of proinflammatory mediators by the same sensory neurons that respond to the presence of pronociceptive mediators. These, in turn, sustain the presence of immune cells, tissue degeneration, and release of more pronociceptive mediators, creating a positive feedback loop of sustained pain sensation, called neurogenic inflammation.⁷⁵ In IC/BPS, neurogenic inflammation may also manifest itself through inflammatory components in the bladder triggering pain sensation and sensitization of the sensory neurons innervating the bladder, contributing to chronic pain.

Pain research has evolved over the years, shifting from measuring physiological pain thresholds and determining hyperalgesia and allodynia to the study of chronic pain by measuring ongoing pain with nonevoked pain measures. Detection of pain thresholds and the existence of hyperalgesia and/or allodynia are helpful in the detection of acute and chronic pain states and also to help determine central sensitization. Nevertheless, these readouts may also exert stress and anxiety in animal models, and the lack of observation of these phenomena does not exclude that they do not exist but may be due to technical characteristics of the experimental design (eg, no difference in hyperalgesia levels between sham and verum after surgical induction of the model, in which the surgical intervention for both groups masks any difference in pain levels between groups), or even to the time when the test is performed. Nowadays, there is growing acknowledgement of the need for longitudinal studies to quantify and characterize the ongoing pain that occurs in chronic pain conditions.¹²⁶ In this regard, several alternatives to measure nonevoked pain have been proposed, including voluntary wheel running,^{39,86,109} recording of ultrasonic vocalisations in rodents,^{101,140} and recording spontaneous rodent behaviour in home-cage monitoring systems. In the latter, a few providers allow simultaneous recording of animals in social groups,^{8,129} opening the possibility of analysing the social effects of pain in rodents, and investigating how these might translate to the social effects of pain in patients.

We argue that further development of these ongoing observations of pain-related behaviours are highly valuable for monitoring outcomes in preclinical models of chronic pain, especially for ENDO and IC/BPS, and would substantially increase their face validity for translational research. Therefore, these methodologies should be taken into account in the experimental design of preclinical models aiming to discover future therapies to treat and hopefully cure ENDO and IC/BPS, as well as improve the patient's quality of life.

As part of our collaborative work between consortium members, the section below includes the perspective of 3 partners representing patient organizations: Judy Birch (The Pelvic Pain Support Network), Lone Hummelshoj (endometriosis.org), and Jane Meijlink (the International Painful Bladder Foundation).

4.5. Patients' perspective on

4.5.1. Pelvic pain and endometriosis

This analysis demonstrates that pain has rarely featured in published articles which aim to establish an effective mouse model for developing treatments for the pelvic pain conditions: ENDO and IC/BPS. It seems from the analysis that there are more articles reporting on pain in IC/BPS than in ENDO, which is puzzling. It is hugely disappointing and surprising/shocking that the importance of pain has received relatively little attention in the

translational research field in these pelvic pain conditions, despite several decades of awareness raising by patients/patient representatives highlighting pain as a priority among clinicians, academics, researchers, policy makers, and the public. This is especially so in the case of ENDO which, despite having attracted increased publicity in recent years, is woefully underfunded in terms of research investment. Regretfully, there has been no consensus to date around a suitable model to develop urgently needed new treatments, addressing not only the pain symptoms associated with ENDO, but also other symptoms including infertility and fatigue. We hope this analysis together with developing a way of accurately assessing pain in rodents may be addressed by the IMI-PainCare TRiPP project ultimately leading to the development of more effective treatments for ENDO and IC/BPS.

4.5.2. Interstitial cystitis/bladder pain syndrome

Interstitial cystitis/bladder pain syndrome covers a wide spectrum of severity, varying from very mild to devastatingly severe and can be an immensely stressful disease. Its hallmark symptoms of pain or discomfort associated with an urgent and frequent need to void day and night can cause great stress, anxiety, and depression, with some patients afraid to leave their home for fear of not finding a toilet in time. The impact on work and (sexual) relationships causes further stress. Many patients say that getting IC/BPS not only changed their life but also their whole character. The fact that it may take years and many doctors to obtain the right diagnosis creates more anxiety and loss of self-confidence and may have a long-lasting detrimental psychosocial effect on the patient. Even in today's world, women's urogenital complaints are still often dismissed in Freudian style as hysteria, and this can lead to long delays in getting treatment.

Because of the fact that in recent decades many patients with unidentifiable bladder pain and urinary symptoms have been piled onto the miscellaneous, heterogeneous heap now known as IC/BPS, treatment continues to be trial and error, whereas many patients never find a successful treatment to relieve their symptoms. Finding an effective treatment for the individual patient will continue to be difficult without meaningful subtyping and phenotyping for both research and treatment purposes.

Many IC/BPS patients suffer from one or multiple comorbid diseases, thereby increasing suffering and making treatment even more complex. More research is needed into the relationship between these comorbidities and IC/BPS and to ascertain whether the bladder disorder forms part of a systemic autoimmune disease in some of these patients.

Where research is concerned, in recent years, too much attention has been focused on the evoked pain aspect alone to the exclusion of urgency and frequency symptoms—which in practical terms may cause the most bother in many patients—as well as nonevoked pain readouts. This has led many drug trials based on evoked pain to fail or to new drugs helping only very few patients. Research can therefore be improved by (1) taking all key symptoms into account including nonevoked pain readouts, sensory urgency,⁹⁴ and urinary frequency and b) phenotyping the patients. For clarification, the term “evoked pain” means the use of sensory stimuli to detect and quantify pain thresholds. By contrast, the term “nonevoked pain” means to record and quantify pain-related behaviour without applying sensory stimuli.

Preclinical research with rodents has tended to look at stress-induced pain behaviour, whereas little research has been performed into the stress and anxiety caused to the patient by the disease itself, by inadequate treatment, by a lack of

understanding and indeed empathy regarding the consequences of the disease and by failure to take a serious look at what is needed to limit the impact of these consequences so as to improve the patient's (psychosocial) quality of life.

4.6. Future perspectives

Chronic pelvic pain/urogenital pain conditions as a whole are underresearched in comparison with other chronic pain conditions. The IMI PainCare with the subproject TRiPP comprises a large public–private partnership between the European Union and the European Federation of Pharmaceutical Industries and Associations. The TRiPP subproject aims to improve disease understanding of 2 specific conditions associated with CPP in women: ENDO because of its prevalence and IC/BPS, a less common condition but associated with a significant impact on quality of life.

The subproject TRiPP will be adopting new approaches to stratify patients by identifying underlying mechanistic pathways, leveraging cross-disciplinary knowledge of pain mechanisms with state-of-the-art biomarker discovery. We will implement a back-translational approach to these conditions, characterising the women suffering with these 2 forms of CPP in our clinical work package and then refining preclinical models of these conditions in light of these findings in our preclinical work packages.

For this purpose, the team is conducting one noninterventional trial (TRiPP NIT1) on deep phenotyping of women with EAP and IC/BPS and will use the findings to evaluate whether women with EAP and IC/BPS can be stratified and to back translate into more relevant preclinical models and improve the translational path to clinical studies.

Thus, our review focuses on these 2 conditions as relevant to this project. Although it would be interesting to expand these analyses of preclinical models of other CPP conditions and of BPS in males, this is beyond the scope of this project/review.

Conflict of interest statement

All coauthors are members of the IMI-PainCare consortium. A. Hoffmann, A. Laux-Biehlmann, B. De Leo, J. Nagel, P. Nunez-Badinez, and T.M. Zollner report personal fees from Bayer AG during the conduct of the study. A. Hoffmann, A. Laux-Biehlmann, B. De Leo, J. Nagel, and T.M. Zollner report personal fees from Bayer AG outside the submitted work. F. Barthas reports personal fees from Grünenthal GmbH during the conduct of the study and outside the submitted work. F. Cruz reports grants and personal fees from Allergan, personal fees from Astellas, personal fees from Bayer, and personal fees from Recordati, outside the submitted work. K. Vincent reports grants from Innovative Medicines Initiative during the conduct of the study; grants and personal fees from Bayer Healthcare, personal fees from Grünenthal GmbH, personal fees from Eli Lilly, and personal fees from AbbVie outside the submitted work. R-D. Treede reports grants from the IMI2 PainCare project of the European Union during the conduct of the study and personal fees from Bayer, Grünenthal, GSK, Sanofi outside the submitted work. J.D. Armstrong reports personal fees and others from Actual Analytics Ltd during the conduct of the study. The remaining authors have no conflicts of interest to declare.

Acknowledgments

This project has received funding from the Innovative Medicines Initiative 2 Joint Undertaking under grant agreement No [777500].

This joint undertaking receives support from the European Union's Horizon 2020 research and innovation programme and the European Federation of Pharmaceutical Industries and Associations.

The statements and opinions presented here reflect the authors' view and neither IMI nor the European Union, the European Federation of Pharmaceutical Industries and Associations, nor any associated partners are responsible for any use that may be made of the information contained herein. www.imi.europa.eu; www.imi-paincare.eu.

Appendix A. Supplemental digital content

Supplemental digital content associated with this article can be found online at <http://links.lww.com/PAIN/B319>.

Article history:

Received 1 October 2020

Received in revised form 15 February 2021

Accepted 18 February 2021

Available online 3 March 2021

References

- Ackerman AL, Jellison FC, Lee UJ, Bradesi S, Rodriguez LV. The Glutamate receptor mediates the establishment and perpetuation of chronic visceral pain in an animal model of stress-induced bladder hyperalgesia. *Am J Physiol Ren Physiol* 2016;310:F628–36.
- Altan ZM, Denis D, Kagan D, Grund EM, Palmer SS, Nataraja SG. A long-acting tumor necrosis factor alpha-binding protein demonstrates activity in both in vitro and in vivo models of endometriosis. *J Pharmacol Exp Ther* 2010;334:460–6.
- Altuntas CZ, Daneshgari F, Sakalar C, Goksoy E, Gulen MF, Kavran M, Qin J, Li X, Tuohy VK. Autoimmunity to uroplakin II causes cystitis in mice: a novel model of interstitial cystitis. *Eur Urol* 2012;61:193–200.
- Alvarez P, Chen X, Hendrich J, Irwin JC, Green PG, Giudice LC, Levine JD. Ectopic uterine tissue as a chronic pain generator. *Neuroscience* 2012;225:269–82.
- American College of O, Gynecologists' Committee on Practice B-G. Chronic pelvic pain: ACOG Practice Bulletin, number 218. *Obstet Gynecol* 2020;135:e98–e109.
- Aziz Q, Giamberardino MA, Barke A, Korwisi B, Baranowski AP, Wessellmann U, Rief W, Treede RD, Pain ITfCoC. The IASP classification of chronic pain for ICD-11: chronic secondary visceral pain. *PAIN* 2019;160:69–76.
- Bacci M, Capobianco A, Monno A, Cottone L, Di Puppo F, Camisa B, Mariani M, Brignole C, Ponzoni M, Ferrari S, Panina-Bordignon P, Manfredi AA, Rovere-Querini P. Macrophages are alternatively activated in patients with endometriosis and required for growth and vascularization of lesions in a mouse model of disease. *Am J Pathol* 2009;175:547–56.
- Bains RS, Cater HL, Sillito RR, Chartsias A, Sneddon D, Concas D, Keskivali-Bond P, Lukins TC, Wells S, Acevedo Arozana A, Nolan PM, Armstrong JD. Analysis of individual mouse activity in group housed animals of different inbred strains using a novel automated home cage analysis system. *Front Behav Neurosci* 2016;10:106.
- Bengochea TS, Olivares CN, Ricci AG, Abramovich D, Parborell F, Meresman GF, Barañao RI, Bilotas MA. Endometriosis impairs ovarian reserve and ovulation in a rat model. *Biocell* 2016;40.
- Bicer F, Altuntas CZ, Izgi K, Ozer A, Kavran M, Tuohy VK, Daneshgari F. Chronic pelvic allodynia is mediated by CCL2 through mast cells in an experimental autoimmune cystitis model. *Am J Physiol Ren Physiol* 2015;308:F103–113.
- Birder L, Andersson KE. Animal modelling of interstitial cystitis/bladder pain syndrome. *Int Neurourol J* 2018;22(suppl 1):S3–9.
- Birder LA. Pathophysiology of interstitial cystitis. *Int J Urol* 2019;26:12–15.
- Boyken J, Laux-Biehlmann A, Nagel J, Zollner TM, Schmidt N. Mimicking early stage endometriosis by intraperitoneal inoculation of uterine tissue induces an acute sterile peritonitis with pelvic pain in mice. *Reprod Sci* 2016;23:249A.
- Brady PC, Horne AW, Saunders PTK, Thomas AM, Missmer SA, Farland LV. Research priorities for endometriosis differ among patients, clinicians, and researchers. *Am J Obstet Gynecol* 2020;222:630–2.
- Bulletti C, Coccia ME, Battistoni S, Borini A. Endometriosis and infertility. *J Assist Reprod Genet* 2010;27:441–7.
- Burma NE, Leduc-Pessah H, Fan CY, Trang T. Animal models of chronic pain: advances and challenges for clinical translation. *J Neurosci Res* 2017;95:1242–56.
- Cagnacci A, Della Vecchia E, Xholli A. Chronic pelvic pain improvement: impact on quality of life and mood. *Gynecol Endocrinol* 2019;35:502–5.
- Campeau L, Soler R, Pareta R, Sivanandane S, Opara E, Andersson KE. Rat bone marrow derived mesenchymal stem cell therapy in a parkinsonian animal model of detrusor overactivity. *J Urol* 2012;187:e44–5.
- Cao J, Boucher W, Kempuraj D, Donelan JM, Theoharides TC. Acute stress and intravesical corticotropin-releasing hormone induces mast cell dependent vascular endothelial growth factor release from mouse bladder explants. *J Urol* 2006;176:1208–13.
- Cetinkaya N, Attar R, Yildirim G, Ficioglu C, Ozkan F, Yilmaz B, Yesildaglar N. The effects of different doses of melatonin treatment on endometrial implants in an oophorectomized rat endometriosis model. *Arch Gynecol Obstet* 2015;291:591–8.
- Chen MC, Keshavan P, Gregory GD, Klumpp DJ. RANTES mediates TNF-dependent lamina propria mast cell accumulation and barrier dysfunction in neurogenic cystitis. *Am J Physiol* 2007;292:F1372–9.
- Chen MC, Mudge CS, Klumpp DJ. Urothelial lesion formation is mediated by TNFR1 during neurogenic cystitis. *Am J Physiol* 2006;291:F741–9.
- Chen S, Xie W, Strong JA, Jiang J, Zhang JM. Sciatic endometriosis induces mechanical hypersensitivity, segmental nerve damage, and robust local inflammation in rats. *Eur J Pain* 2016;20:1044–57.
- Chen WC, Lee MH, Chiu C, Wu HC. The importance of biopsychological care in women with interstitial cystitis/bladder pain syndrome (IC/BPS) who have child abuse experience and psychiatric dysfunction. *Neurourol Urodyn* 2013;32:601–2.
- Cheng C, Rosamilia A, Healey M. Diagnosis of interstitial cystitis/bladder pain syndrome in women with chronic pelvic pain—a prospective observational study. *Int Urogynecol J Pelvic Floor Dysfunct* 2011;22:S1785–6.
- Cheng C, Rosamilia A, Healey M. Diagnosis of interstitial cystitis/bladder pain syndrome in women with chronic pelvic pain: a prospective observational study. *Int Urogynecol J Pelvic Floor Dysfunct* 2012;23:1361–6.
- Cheng CW, Licence D, Cook E, Luo F, Arends MJ, Smith SK, Print CG, Charnock-Jones DS. Activation of mutated K-ras in donor endometrial epithelium and stroma promotes lesion growth in an intact immunocompetent murine model of endometriosis. *J Pathol* 2011;224:261–9.
- Chiu CD, Lee MH, Chen WC, Ho HL, Wu HC. Childhood trauma perpetrated by close others, psychiatric dysfunction, and urological symptoms in patients with interstitial cystitis/bladder pain syndrome. *J Psychosomatic Res* 2017;93:90–5.
- Chuang YC, Yang LC, Chiang PH, Kang HY, Ma WL, Wu PC, DeMiguel F, Chancellor MB, Yoshimura N. Gene gun particle encoding preproenkephalin cDNA produces analgesia against capsaicin-induced bladder pain in rats. *Urology* 2005;65:804–10.
- Cox A. Management of interstitial cystitis/bladder pain syndrome. *Can Urol Assoc J* 2018;12:S157–60.
- Cox PJ. Cyclophosphamide cystitis—identification of acrolein as the causative agent. *Biochem Pharmacol* 1979;28:2045–9.
- Coxon L, Horne AW, Vincent K. Pathophysiology of endometriosis-associated pain: a review of pelvic and central nervous system mechanisms. *Best Pract Res Clin Obstet Gynaecol* 2018;51:53–67.
- Dabrosin C, Gyorffy S, Margetts P, Ross C, Gauldie J. Therapeutic effect of angiostatin gene transfer in a murine model of endometriosis. *Am J Pathol* 2002;161:909–18.
- Delsouc MB, Ghersa F, Ramirez D, Della Vedova MC, Gil RA, Vallcaneras SS, Casais M. Endometriosis progression in tumor necrosis factor receptor p55-deficient mice: impact on oxidative/nitrosative stress and metalloprotein profile. *J Trace Elem Med Biol* 2019;52:157–65.
- Ercan F, Cetinel S, Erin N, Aydin H, Hürdağ C, Parker T, Parker K, Mayhew T. Volume of nerve fibers in the stress-induced bladder of adult rats following capsaicin treatment. *Urol Int* 2003;71:393–8.
- Fattori V, Franklin NS, Gonzalez-Cano R, Peterse D, Madrian E, Verri WA, Andrews N, Woolf CJ, Rogers MS. A non-surgical murine model of endometriosis-associated pain that predicts response to clinically active drugs. *Reprod Sci* 2019;26:86A.

- [37] Ferrero H, Buigues A, Martinez J, Simon C, Pellicer A, Gomez R. A novel homologous model for noninvasive monitoring of endometriosis progression. *Biol Reprod* 2017;96:302–12.
- [38] Fraser MO, Chuang YC, Lavelle JP, Yoshimura N, de Groat WC, Chancellor MB. A reliable, nondestructive animal model for interstitial cystitis: intravesical low-dose protamine sulfate combined with physiological concentrations of potassium chloride. *Urology* 2001;57(6 suppl 1):112.
- [39] Fuentes I, Pierce A, Eller-Smith O, Christianson J. Voluntary wheel running can improve increased urogenital sensitivity and function resulting from neonatal maternal separation in male mice. *J Pain* 2017;18:S10.
- [40] Fuentes IM, Pierce AN, Di Silvestro ER, Maloney MO, Christianson JA. Differential influence of early life and adult stress on urogenital sensitivity and function in male mice. *Front Syst Neurosci* 2017;11:97.
- [41] Gater A, Wichmann K, Seitz C, Gerlinger C, Taylor F. Assessing the patient-reported symptoms and impacts of endometriosis—development of the endometriosis symptom diary and endometriosis impact scale. *J Endometriosis* 2013;5:S40.
- [42] Ghera F, Delsouc MB, Goyeneche AA, Vallcaneras SS, Meresman G, Telleria CM, Casais M. Reduced inflammatory state promotes reinnervation of endometriotic-like lesions in TNFRp55 deficient mice. *Mol Hum Reprod* 2019;25:385–96.
- [43] Giamberardino MA, Berkley KJ, Affaitati G, Lerza R, Centurione L, Lapenna D, Vecchiet L. Influence of endometriosis on pain behaviors and muscle hyperalgesia induced by a ureteral calculosis in female rats. *PAIN* 2002;95:247–57.
- [44] Giudice LC. Clinical practice. Endometriosis. *N Engl J Med* 2010;362:2389–98.
- [45] Golubeva AV, Zhdanov AV, Mallel G, Dinan TG, Cryan JF. The mouse cyclophosphamide model of bladder pain syndrome: tissue characterization, immune profiling, and relationship to metabotropic glutamate receptors. *Physiol Rep* 2014;2:e00260.
- [46] Greaves E, Cousins FL, Murray A, Esnal-Zufiaurre A, Fassbender A, Horne AW, Saunders PT. A novel mouse model of endometriosis mimics human phenotype and reveals insights into the inflammatory contribution of shed endometrium. *Am J Pathol* 2014;184:1930–9.
- [47] Greaves E, Horne AW, Jerina H, Mikolajczak M, Hilferty L, Mitchell R, Fleetwood-Walker SM, Saunders PT. EP2 receptor antagonism reduces peripheral and central hyperalgesia in a preclinical mouse model of endometriosis. *Sci Rep* 2017;7:44169.
- [48] Guo S, He W, Liu X. Evidence for central sensitization in women with endometriosis. *Int J Gynecol Obstet* 2009;107:S196–7.
- [49] Guo SW. Recurrence of endometriosis and its control. *Hum Reprod Update* 2009;15:441–61.
- [50] Guo SW. Methodological issues in preclinical mouse efficacy studies of adenomyosis. *Curr Obstet Gynecol Rep* 2012;1:138–45.
- [51] Han DY, Jeong HJ, Lee MY. Bladder hyperactivity induced by chronic variable stress in rats. *Low Urin Tract Symptoms* 2015;7:56–61.
- [52] He Y, Zeng JF, Liu YH. Comparison of autotransplantation sites in endometriosis rat model. *Acad J Sec Mil Med Univ* 2016;37:513–16.
- [53] Heard ME, Velarde MC, Giudice LC, Simmen FA, Simmen RC. Kruppel-like factor 13 deficiency in uterine endometrial cells contributes to defective steroid hormone receptor signaling but not lesion establishment in a mouse model of endometriosis. *Biol Reprod* 2015;92:140.
- [54] Heaton RL, Walid MS. Endometriosis/adenomyosis is associated with more typical cystoscopic findings of interstitial cystitis in patients with elevated PUF scores. *Gynecol Surg* 2010;7:353–7.
- [55] Hirata T, Osuga Y, Yoshino O, Hirota Y, Harada M, Takemura Y, Morimoto C, Koga K, Yano T, Tsutsumi O, Taketani Y. Development of an experimental model of endometriosis using mice that ubiquitously express green fluorescent protein. *Hum Reprod* 2005;20:2092–6.
- [56] Horne AW, Saunders PTK. SnapShot: endometriosis. *Cell* 2019;179:1677–e1671.
- [57] Horne AW, Saunders PTK, Abokhras IM, Hogg L. Endometriosis priority setting partnership steering G. Top ten endometriosis research priorities in the UK and Ireland. *Lancet* 2017;389:2191–2.
- [58] Iguchi N, Pineda R, Carrasco A, Dönmez MI, Malykhina A, Wilcox D. Constipation induces overactive bladder through alterations of nicotinic and purinergic pathways in mouse bladders. *J Urol* 2018;199:e506.
- [59] Imamura T, Ishizuka O, Aizawa N, Zhong C, Ogawa T, Nakayama T, Tanabe T, Nishizawa O. Cold environmental stress induces detrusor overactivity via resniferatoxin-sensitive nerves in conscious rats. *Neurourol Urodyn* 2008;27:348–52.
- [60] Iuvone T, Affaitati G, De Filippis D, Lopopolo M, Grassia G, Lapenna D, Negro L, Costantini R, Vaia M, Cipollone F, Ialenti A, Giamberardino MA. Ultramicronized palmitoylethanolamide reduces viscerovisceral hyperalgesia in a rat model of endometriosis plus ureteral calculosis: role of mast cells. *PAIN* 2016;157:80–91.
- [61] Jia SZ, Leng JH, Shi JH, Sun PR, Lang JH. Health-related quality of life in women with endometriosis: a systematic review. *J Ovarian Res* 2012;5:861–70.
- [62] Jin XW, Liu BK, Zhang X, Zhao ZH, Shao Y. Establishment of a novel autoimmune experimental model of bladder pain syndrome/interstitial cystitis in C57BL/6 mice. *Inflammation* 2017;40:861–70.
- [63] Kawamori N, Yoshikawa S, Kashyap M, Tyagi P, Arai Y, Chancellor MB, Yoshimura N. Liposome based intravesical therapy targeting nerve growth factor ameliorates bladder hypersensitivity in rats with experimental colitis. *J Urol* 2016;195:1920–6.
- [64] Keay S, Leitzell S, Ochrczin A, Clements G, Zhan M, Johnson D. A mouse model for interstitial cystitis/painful bladder syndrome based on APF inhibition of bladder epithelial repair: a pilot study. *BMC Urol* 2012;12:17.
- [65] Kim A, Han JY, Ryu CM, Yu HY, Lee S, Kim Y, Jeong SU, Cho YM, Shin DM, Choo MS. Histopathological characteristics of interstitial cystitis/bladder pain syndrome without Hunner lesion. *Histopathology* 2017;71:415–24.
- [66] Kitta T, Nonomura K, Yoshimura N. Effect of Adenosine A2A receptor antagonist on bladder overactivity in rats with chemical cystitis. *J Urol* 2010;183:e651.
- [67] Kogan P, Xu S, Wang Y, O'Donnell MA, Lutgendorf SK, Bradley CS, Schrepf A, Kreder KJ, Luo Y. Sub-noxious intravesical lipopolysaccharide triggers bladder inflammation and symptom onset in A transgenic autoimmune cystitis model: a MAPP network animal study. *Sci Rep* 2018;8:6573.
- [68] Kouzoukas DE, Ma F, Meyer-Siegler KL, Westlund KN, Hunt DE, Vera PL. Protease-activated receptor 4 induces bladder pain through high mobility group Box-1. *PLoS One* 2016;11:e0152055.
- [69] Lai HH, Gardner V, Ness TJ, Gereau IRW. Segmental hyperalgesia to mechanical stimulus in interstitial cystitis/bladder pain syndrome: evidence of central sensitization. *J Urol* 2014;191:1294–9.
- [70] Lai HH, Thu JHL, Moh FV, Paradis A, Vetter J. Clustering of patients with interstitial cystitis/bladder pain syndrome and chronic prostatitis/chronic pelvic pain syndrome. *J Urol* 2019;202:546–51.
- [71] Lanteri-Minet M, Bon K, de Pommery J, Michiels JF, Menetrey D. Cyclophosphamide cystitis as a model of visceral pain in rats: model elaboration and spinal structures involved as revealed by the expression of c-Fos and Krox-24 proteins. *Exp Brain Res* 1995;105:220–32.
- [72] Laschke MW, Elitzsch A, Vollmar B, Vajkoczy P, Menger MD. Combined inhibition of vascular endothelial growth factor (VEGF), fibroblast growth factor and platelet-derived growth factor, but not inhibition of VEGF alone, effectively suppresses angiogenesis and vessel maturation in endometriotic lesions. *Hum Reprod* 2006;21:262–8.
- [73] Laux-Biehlmann A, Boyken J, Koch M, Dahlhof H, Zollner TM, Nagel J, Schmidt N. Retrograde menstruation causes pain and inflammation in mice. *Reprod Sci* 2015;22:207A.
- [74] Laux-Biehlmann A, Buiaakova O, Hayward MD, Zollner TM, Nagel J. Anxiety related behaviors in a mouse model of endometriosis. *Reprod Sci* 2016;23:163A.
- [75] Laux-Biehlmann A, d'Hooghe T, Zollner TM. Menstruation pulls the trigger for inflammation and pain in endometriosis. *Trends Pharmacol Sci* 2015;36:270–6.
- [76] Li J, Micevych P, McDonald J, Rapkin A, Chaban V. Inflammation in the uterus induces phosphorylated extracellular signal-regulated kinase and substance P immunoreactivity in dorsal root ganglia neurons innervating both uterus and colon in rats. *J Neurosci Res* 2008;86:2746–52.
- [77] Liang B, Wu L, Xu H, Cheung CW, Fung WY, Wong SW, Wang CC. Efficacy, safety and recurrence of new progestins and selective progesterone receptor modulator for the treatment of endometriosis: a comparison study in mice. *Reprod Biol Endocrinol* 2018;16:32.
- [78] Ligon C, Mohammadi E, Ge P, Hannig G, Higgins C, Greenwood-Van Meerveld B. Linacotide inhibits colonic and urinary bladder hypersensitivity in adult female rats following unpredictable neonatal stress. *Neurogastroenterol Motil* 2018;30:e13375.
- [79] Lin YH, Liu G, Kavan M, Altuntas CZ, Gasbarro G, Tuohy VK, Daneshgari F. Lower urinary tract phenotype of experimental autoimmune cystitis in mouse: a potential animal model for interstitial cystitis. *BJU Int* 2008;102:1724–30.
- [80] Liu W, Deyoung BR, Chen X, Evanoff DP, Luo Y. RDP58 inhibits T cell-mediated bladder inflammation in an autoimmune cystitis model. *J Autoimmun* 2008;30:257–65.
- [81] Long Q, Liu X, Guo SW. Surgery accelerates the development of endometriosis in mice. *Am J Obstet Gynecol* 2016;215:320.e321–320.e315.

- [82] Lopopolo M, Affaitati G, Fabrizio A, Massimini F, Lapenna D, Giamberardino MA, Costantini R. Effects of tramadol on viscerovisceral hyperalgesia in a rat model of endometriosis plus ureteral calculosis. *Fundam Clin Pharmacol* 2014;28:331–41.
- [83] Lv J, Huang Y, Zhu S, Yang G, Zhang Y, Leng J, Bo J, Liu D. MCP-1-induced histamine release from mast cells is associated with development of interstitial cystitis/bladder pain syndrome in rat models. *Mediators Inflamm* 2012;2012:358184.
- [84] Malde S, Palmisani S, Al-Kaisy A, Sahai A. Guideline of guidelines: bladder pain syndrome. *BJU Int* 2018;122:729–43.
- [85] Malykhina AP, Qin C, Lei Q, Pan XQ, Greenwood-Van Meerveld B, Foreman RD. Differential effects of intravesical resiniferatoxin on excitability of bladder spinal neurons upon colon-bladder cross-sensitization. *Brain Res* 2013;1491:213–24.
- [86] Manzanares G, Brito-da-Silva G, Gandra PG. Voluntary wheel running: patterns and physiological effects in mice. *Braz J Med Biol Res* 2018;52:e7830.
- [87] Marcu I, Campian EC, Tu FF. Interstitial cystitis/bladder pain syndrome. *Semin Reprod Med* 2018;36:123–35.
- [88] Marcu I, Westrich DJ, Hammami MB, Vareedayah A, Mogalapalli A, Mogalapalli N, McFerrin CA, Campian EC, Marsicano EV. Relationship between interstitial cystitis (IC) with or without cystoscopic findings and gastroenterological (GI) functional and mucosal disorders. *Female Pelvic Med Reconstr Surg* 2018;24:S146.
- [89] Matos R, Cordeiro JM, Coelho A, Ferreira S, Silva C, Igawa Y, Cruz F, Charrua A. Bladder pain induced by prolonged peripheral alpha 1A adrenoceptor stimulation involves the enhancement of transient receptor potential vanilloid 1 activity and an increase of urothelial adenosine triphosphate release. *Acta Physiol* 2016;218:265–75.
- [90] Matos R, Cruz F, Charrua A. Childhood stressful events induce chronic bladder pain in adulthood through a TRPV1 dependent mechanism. *J Urol* 2017;197:e380–1.
- [91] Matos R, Serrão P, Rodrigues L, Birder LA, Cruz F, Charrua A. The water avoidance stress induces bladder pain due to a prolonged adrenergic (alpha1A) stimulation of the bladder. *Eur Urol Supplements* 2017;16:e350–1.
- [92] McKernan LC, Johnson BN, Reynolds WS, Williams DA, Cheavens JS, Dmochowski RR, Crofford LJ. Posttraumatic stress disorder in interstitial cystitis/bladder pain syndrome: relationship to patient phenotype and clinical practice implications. *Neurourol Urodyn* 2019;38:353–62.
- [93] Mei J, Zhou WJ, Zhu XY, Lu H, Wu K, Yang HL, Fu Q, Wei CY, Chang KK, Jin LP, Wang J, Wang YM, Li DJ, Li MQ. Suppression of autophagy and HCK signaling promotes PTGS2high FCGR3–NK cell differentiation triggered by ectopic endometrial stromal cells. *Autophagy* 2018;14:1376–97.
- [94] Meijlink J. An urgent case for sensory urgency: a patient perspective. *Neurourol Urodyn* 2020;39:2008–10.
- [95] Meuleman C, Vandenabeele B, Fieuws S, Spiessens C, Timmerman D, D'Hooghe T. High prevalence of endometriosis in infertile women with normal ovulation and normospermic partners. *Fertil Steril* 2009;92:68–74.
- [96] Mingin GC, Heppner TJ, Tykocki NR, Erickson CS, Vizzard MA, Nelson MT. Social stress in mice induces urinary bladder overactivity and increases TRPV1 channel-dependent afferent nerve activity. *Am J Physiol* 2015;309:R629–38.
- [97] Mizoguchi S, Wolf-Johnson AS, Suzuki T, Takaoka E, DeFranco DB, Wang Z, Birder LA, Yoshimura N. Bladder urothelial and afferent dysfunctions underlying bladder overactivity in rats with chemically induced prostatic inflammation. *J Urol* 2018;199:e599.
- [98] Mo L, Cheng J, Lee EYHP, Sun TT, Wu XR. Gene deletion in urothelium by specific expression of Cre recombinase. *Am J Physiol* 2005;289:F562–8.
- [99] Nishijima S, Sugaya K, Kadekawa K, Ashitomi K, Ueda T, Yamamoto H. High-dose tranilast administration to rats creates interstitial cystitis-like symptoms with increased vascular permeability. *Life Sci* 2013;93:897–903.
- [100] Offiah I, Didangelos A, O'Reilly BA, McMahon SB. Manipulating the extracellular matrix: an animal model of the bladder pain syndrome. *PAIN* 2017;158:161–70.
- [101] Oliveira AR, Barros HM. Ultrasonic rat vocalizations during the formalin test: a measure of the affective dimension of pain? *Anesth analgesia* 2006;102:832–9.
- [102] Orr N, Wahl K, Noga H, Albert A, Smith K, Yong P. 072 increased central sensitization among women with painful bladder syndrome and deep dyspareunia. *J Sex Med* 2019;16:S26–7.
- [103] Patnaik SS, Laganà AS, Vitale SG, Buttice S, Noventa M, Gizzo S, Valenti G, Rapisarda AMC, la Rosa VL, Magno C, Triolo O, Dandolu V. Etiology, pathophysiology and biomarkers of interstitial cystitis/painful bladder syndrome. *Arch Gynecol Obstet* 2017;295:1341–59.
- [104] Paul S, Sharma AV, Mahapatra PD, Bhattacharya P, Reiter RJ, Swarnakar S. Role of melatonin in regulating matrix metalloproteinase-9 via tissue inhibitors of metalloproteinase-1 during protection against endometriosis. *J Pineal Res* 2008;44:439–49.
- [105] Persoons E, De Clercq K, Van den Eynde C, Pinto SJPC, Luyten K, Van Bree R, Tomassetti C, Voets T, Vriens J. Mimicking Sampson's retrograde menstrual theory in rats: a new rat model for ongoing endometriosis-associated pain. *Int J Mol Sci* 2020;21:2326.
- [106] Persoons E, De Clercq K, Vriens J, Pinto SPC. A new rat model to investigate endometriosis-associated pain using menstrual endometrial tissue. *Reprod Sci* 2019;26:223A–4A.
- [107] Pierce AN, DiSilvestro ER, Eller OC, Wang R, Ryals JM, Christianson JA. Urinary bladder hypersensitivity and dysfunction in female mice following early life and adult stress. *Brain Res* 2016;1639:58–73.
- [108] Pierce AN, Eller-Smith OC, Christianson JA. Voluntary wheel running attenuates urinary bladder hypersensitivity and dysfunction following neonatal maternal separation in female mice. *Neurourol Urodyn* 2018;37:1623–32.
- [109] Pitzer C, Kuner R, Tappe-Theodor A. Voluntary and evoked behavioral correlates in inflammatory pain conditions under different social housing conditions. *Pain Rep* 2016;1:e564.
- [110] Porru D, Bobbi V, Di Franco C, Viglio A, Novario M, Gardella B, Nappi RE, Tinelli C, Spinillo A, Paulli M, Rovereto B. Time of symptom onset and histologic findings in bladder pain syndrome/interstitial cystitis (IC/BPS): new findings on useful correlations. *Neurourol Urodyn* 2018;37:S57–9.
- [111] Rebar RW, Morandini IC, Erickson GF, Petze JE. The hormonal basis of reproductive defects in athymic mice: diminished gonadotropin concentrations in prepubertal females. *Endocrinology* 1981;108:120–6.
- [112] Robbins MT, Ness TJ. Footshock-induced urinary bladder hypersensitivity: role of spinal corticotropin-releasing factor receptors. *J Pain* 2008;9:991–8.
- [113] Rudick CN, Schaeffer AJ, Klumpp DJ. Pharmacologic attenuation of pelvic pain in a murine model of interstitial cystitis. *BMC Urol* 2009;9:16.
- [114] Ruiz A, Rockfield S, Taran N, Haller E, Engelman RW, Flores I, Panina-Bordignon P, Nanjundan M. Effect of hydroxychloroquine and characterization of autophagy in a mouse model of endometriosis. *Cell Death Dis* 2016;7:e2059.
- [115] Schliep KC, Mumford SL, Peterson CM, Chen Z, Johnstone EB, Sharp HT, Stanford JB, Hammoud AO, Sun L, Buck Louis GM. Pain typology and incident endometriosis. *Hum Reprod* 2015;30:2427–38.
- [116] Schwartz ES, La JH, Young EE, Feng B, Joyce S, Gebhart GF. Chronic prostatitis induces bladder hypersensitivity and sensitizes bladder afferents in the mouse. *J Urol* 2016;196:892–901.
- [117] Sewell M, Churilov L, Mooney S, Ma T, Maher P, Grover SR. Chronic pelvic pain—pain catastrophizing, pelvic pain and quality of life. *Scand J Pain* 2018;18:441–8.
- [118] Shimizu S, Shimizu T, Nakamura K, Aratake T, Zou S, Hamada T, Nagao Y, Yamamoto M, Honda M, Hiagashi Y, Saito M. Involvement of brain gabaa receptor for angiotensin II-induced frequent urination in the rat. *Neurourol Urodyn* 2017;36:S485–6.
- [119] Simitsidellis I, Gibson DA, Saunders PTK. Animal models of endometriosis: replicating the aetiology and symptoms of the human disorder. *Best Pract Res Clin Endocrinol Metab* 2018;32:257–69.
- [120] Somigliana E, Viganò P, Rossi G, Carinelli S, Vignali M, Panina-Bordignon P. Endometrial ability to implant in ectopic sites can be prevented by interleukin-12 in a murine model of endometriosis. *Hum Reprod* 1999;14:2944–50.
- [121] Stojanow K, Leeners B, Bartley J, Mechsner S, Rauchfuss M. Psychological well-being in women with endometriosis: differences between patients with and without chronic pelvic pain. *J Endometriosis* 2013;5:S43–4.
- [122] Sugino Y, Nishikawa N, Yoshimura K, Kuno S, Hayashi Y, Yoshimura N, Okazaki T, Kanematsu A, Ogawa O. BALB/c-Fcgr2bPdc1 mouse expressing anti-urothelial antibody is a novel model of autoimmune cystitis. *Sci Rep* 2012;2:317.
- [123] Suskind AM, Berry SH, Sutorp MJ, Elliott MN, Hays RD, Ewing BA, Clemens JQ. Health-related quality of life in patients with interstitial cystitis/bladder pain syndrome and frequently associated comorbidities. *Qual Life Res* 2013;22:1537–41.
- [124] Takai E, Taniguchi F, Nakamura K, Uegaki T, Iwabe T, Harada T. Parthenolide reduces cell proliferation and prostaglandin estradiol synthesis in human endometriotic stromal cells and inhibits development of endometriosis in the murine model. *Fertil Steril* 2013;100:1170–8.

- [125] Tappe-Theodor A, King T, Morgan MM. Pros and cons of clinically relevant methods to assess pain in rodents. *Neurosci Biobehav Rev* 2019;100:335–43.
- [126] Tappe-Theodor A, Kuner R. Studying ongoing and spontaneous pain in rodents—challenges and opportunities. *Eur J Neurosci* 2014;39:1881–90.
- [127] Thomassin I, Bazot M, Detchev R, Barranger E, Cortez A, Darai E. Symptoms before and after surgical removal of colorectal endometriosis that are assessed by magnetic resonance imaging and rectal endoscopic sonography. *Am J Obstet Gynecol* 2004;190:1264–71.
- [128] Tripoli TM, Sato H, Sartori MG, De Araujo FF, Girão MJ, Schor E. Evaluation of quality of life and sexual satisfaction in women suffering from chronic pelvic pain with or without endometriosis. *J Sex Med* 2011;8:497–503.
- [129] Tse K, Sillito R, Keerie A, Collier R, Grant C, Karp NA, Vickers C, Chapman K, Armstrong JD, Redfern WS. Pharmacological validation of individual animal locomotion, temperature and behavioural analysis in group-housed rats using a novel automated home cage analysis system: a comparison with the modified Irwin test. *J Pharmacol Toxicol Methods* 2018;94:1–13.
- [130] Ustinova EE, Gutkin DW, Pezzone MA. Sensitization of pelvic nerve afferents and mast cell infiltration in the urinary bladder following chronic colonic irritation is mediated by neuropeptides. *Am J Physiol Ren Physiol* 2007;292:F123–30.
- [131] van de Merwe JP, Yamada T, Sakamoto Y. Systemic aspects of interstitial cystitis, immunology and linkage with autoimmune disorders. *Int J Urol* 2003;10(suppl):S35–38.
- [132] Vasudevan V, Moldwin R. Addressing quality of life in the patient with interstitial cystitis/bladder pain syndrome. *Asian J Urol* 2017;4:50–4.
- [133] Vercellini P, Buggio L, Frattaruolo MP, Borghi A, Dridi D, Somigliana E. Medical treatment of endometriosis-related pain. *Best Pract Res Clin Obstet Gynaecol* 2018;51:68–91.
- [134] Vercellini P, Fedele L, Aimi G, Pietropaolo G, Consonni D, Crosignani PG. Association between endometriosis stage, lesion type, patient characteristics and severity of pelvic pain symptoms: a multivariate analysis of over 1000 patients. *Hum Reprod* 2007;22:266–71.
- [135] Verma-Gandhu M, Bercik P, Motomura Y, Verdu EF, Khan WI, Blennerhassett PA, Wang L, El-Sharkawy RT, Collins SM. CD4+ T-cell modulation of visceral nociception in mice. *Gastroenterology* 2006;130:1721–8.
- [136] Vianello A, Soler R, Fuhllase C, Soker S, Atala A, Yoo JJ, Williams K, Porena M. Endothelial cell therapy to prevent radiation-induced cystitis in a rat model. *Eur Urol Suppl* 2010;9:293.
- [137] Wang Z, Chang HH, Gao Y, Zhang R, Guo Y, Holschneider DP, Rodríguez LV. Effects of water avoidance stress on peripheral and central responses during bladder filling in the rat: a multidisciplinary approach to the study of urologic chronic pelvic pain syndrome (MAPP) research network study. *PLoS One* 2017;12:e0182976.
- [138] Whitmore KE, Fall M, Sengiku A, Tomoe H, Logadottir Y, Kim YH. Hunner lesion versus non-Hunner lesion interstitial cystitis/bladder pain syndrome. *Int J Urol* 2019;26:26–34.
- [139] Wieser F, Wu J, Shen Z, Taylor RN, Sidell N. Retinoic acid suppresses growth of lesions, inhibits peritoneal cytokine secretion, and promotes macrophage differentiation in an immunocompetent mouse model of endometriosis. *Fertil Steril* 2012;97:1430–7.
- [140] Williams WO, Riskin DK, Mott AK. Ultrasonic sound as an indicator of acute pain in laboratory mice. *J Am Assoc Lab Anim Sci* 2008;47:8–10.
- [141] Wu CC, Chung SD, Lin HC. Endometriosis increased the risk of bladder pain syndrome/interstitial cystitis: a population-based study. *Neurourol Urodyn* 2018;37:1413–18.
- [142] Yang W, Yaggie R, Jiang M, Rudick C, Done J, Heckman C, Schaeffer A, Klumpp D. Acyloxyacyl hydrolase modulates pelvic pain severity. *J Urol* 2017;197:e1155.
- [143] Yu J, Zhao L, Zhang D, Zhai D, Shen W, Bai L, Liu Y, Cai Z, Li J, Yu C. The effects and possible mechanisms of puerarin to treat endometriosis model rats. *Evid Based Complement Altern Med* 2015;2015:269138.
- [144] Zhang L, Ihsan AU, Cao Y, Khan FU, Cheng Y, Han L, Zhou X. An immunogenic peptide, T2 induces interstitial cystitis/painful bladder syndrome: an autoimmune mouse model for interstitial cystitis/painful bladder syndrome. *Inflammation* 2017;40:2033–41.
- [145] Zondervan KT, Becker CM, Koga K, Missmer SA, Taylor RN, Viganò P. Endometriosis. *Nat Rev Dis Primers* 2018;4:9.