



ScienceDirect

Contents lists available at sciencedirect.com
Journal homepage: www.elsevier.com/locate/jval

Economic Evaluation

Cost-Effectiveness Analysis of Treating Patients With NTRK-Positive Cancer With the Histology-Independent Therapy Entrectinib



Simone Huygens, PhD,* Heleen Vellekoop, MSc, Matthijs Versteegh, PhD, Irene Santi, PhD, László Szilberhorn, PhD, Tamás Zelei, PhD, Balázs Nagy, PhD, Apostolos Tsiachristas, PhD, Rositsa Koleva-Kolarova, PhD, Sarah Wordsworth, PhD, Maureen Rutten-van Mölken, PhD, on behalf of the HEcoPerMed Consortium

ABSTRACT

Objectives: This study tackles several challenges of evaluating histology-independent treatments using entrectinib as an example. Histology-independent treatments are provided based on genetic marker(s) of tumors, regardless of the tumor type. We evaluated the lifetime cost-effectiveness of testing all patients for NTRK fusions and treating the positive cases with entrectinib compared with no testing and standard of care (SoC) for all patients.

Methods: The health economic model consisted of a decision tree reflecting the NTRK testing phase followed by a micro-simulation model reflecting treatment with either entrectinib or SoC. Efficacy of entrectinib was based on data from basket trials, whereas historical data from NTRK-negative patients were corrected for the prognostic value of NTRK fusions to model SoC.

Results: “Testing” (testing for NTRK fusions, with subsequent entrectinib treatment in NTRK-positive patients and SoC in NTRK-negative patients) had higher per-patient quality-adjusted life-years (QALYs) and costs than “No testing” (SoC for all patients), with a difference of 0.0043 and €732, respectively. This corresponded to an incremental cost-effectiveness ratio (ICER) of €169 957/QALY and, using a cost-effectiveness threshold of €80 000/QALY, an incremental net monetary benefit of –€388. When excluding the costs of genetic testing for NTRK fusions, the ICER was reduced to €36 290/QALY and the incremental net monetary benefit increased to €188.

Conclusions: When treatment requires the identification of a genetic marker, the associated costs and effects need to be accounted for. Because of the low prevalence of NTRK fusions, the number needed-to-test to identify patients eligible for entrectinib is large. Excluding the testing phase reduces the ICER substantially.

Keywords: histology-independent treatment, tumor-agnostic therapy, tissue-independent therapy.

VALUE HEALTH. 2023; 26(2):193–203

Introduction

Recently, the European Medicines Agency (EMA) approved the first histology-independent treatments, entrectinib and larotrectinib, for tumors with NTRK gene fusions.^{1,2} Histology-independent (also called “tumor-agnostic”) treatments are prescribed based on a genetic marker of the tumor, whereas most other oncology treatments are prescribed based on the tumor type. Evaluating the efficacy and cost-effectiveness of existing histology-independent treatments has proven challenging for various reasons.³

First, clinical trials for entrectinib and larotrectinib were basket trials where patients with different tumor types were pooled together.^{4–6} Because of the small number of patients per tumor type, tumor-specific effectiveness was not provided and marketing authorization for the pharmaceuticals was granted for all

NTRK-positive (NTRK+) tumors, assuming similar efficacy across tumor types. Nevertheless, there may be heterogeneity in treatment effect across the tumor types.⁷ Second, the trials were single-arm trials. The lack of randomized controlled trial (RCT) data for entrectinib and larotrectinib creates additional uncertainty around their effectiveness. Although historical data might be used to construct a synthetic control arm to the trial arm, it can be highly difficult to ascertain that the patient populations in control and intervention arm are sufficiently comparable.⁸ A key issue for histology-independent treatments is that all patients in the trial have tumors with a specific genetic marker, whereas the available historical data is likely for a mixed patient population with tumors with and without the genetic marker. Given that genetic markers may affect disease prognosis, historical data from patients without the genetic marker should be corrected for the prognostic value of the genetic marker.⁹

*Simone Huygens and Heleen Vellekoop contributed equally as first authors.

Third, for most tumor types the standard of care (SoC) does not include testing for NTRK fusions, meaning that the introduction of TRK inhibitors would also require the introduction of NTRK testing. Evaluating the cost-effectiveness of a new treatment requires a comparison between the new situation, in which the intervention is implemented, and the current situation.⁹ To accurately reflect the new situation, all changes to the care pathway that are needed to identify and treat eligible patients need to be accounted for. That is, the health and cost consequences associated with introducing NTRK testing should be included in the cost-effectiveness analysis of TRK inhibitors. Various topics are to be considered when modeling tests, including the expected testing procedures in clinical practice, test properties (eg, sensitivity and specificity), and mortality during the testing phase.⁹

In this article, we estimate the cost-effectiveness of entrectinib compared with SoC in cancer patients in The Netherlands. We compare a strategy in which patients are tested for NTRK fusions and receive subsequent treatment (entrectinib for NTRK+ patients and SoC for NTRK-negative [NTRK-] patients) to a strategy in which no additional testing is used and all patients receive SoC. We illustrate how some of the challenges arising from single-arm trial data may be addressed (the second challenge mentioned above) and how testing pathways can be incorporated in cost-effectiveness analyses (third challenge).

Methods

Intervention and Comparator

The intervention comprised NTRK gene fusion testing for all patients with locally advanced or metastatic solid tumors followed by treatment with entrectinib in NTRK+ patients and SoC in NTRK- patients. Patients in the comparator arm were not tested for NTRK fusions and all patients received SoC. National tumor-specific treatment guidelines were used to identify the treatments provided in SoC for each tumor type (Appendix Table 1 in Supplemental Materials found at <https://doi.org/10.1016/j.jval.2022.08.006>). We included treatments indicated for patients with locally advanced or metastatic solid tumors who had already received (at least) 1 systemic anticancer therapy.^{2,10} Experimental treatments (ie, treatments outside current clinical guidelines) were excluded.

Larotrectinib also targets oncology patients with NTRK gene fusions and could be an alternative to entrectinib. Nevertheless, we were unable to include both pharmaceuticals in the model because of key differences in the trial populations, including the distribution of tumor types and the presence of pediatric patients in the larotrectinib trial but not the entrectinib trial. Publicly available data are insufficient to adjust the estimated effectiveness in the entrectinib and larotrectinib trials for differences in the trial populations. We opted to include entrectinib in our analysis because we had data available on the fitted distribution for overall survival (OS) and treatment discontinuation for entrectinib, whereas this data was not available for larotrectinib.

Study Population

The study population included adult patients with locally advanced or metastatic solid tumors who have received one or more lines of treatment and are willing to undergo further testing and treatment. Although entrectinib also received EMA approval for pediatric patients ≥ 12 years, there were no pediatric patients included in the ALKA-372-001, STARTRK-1, and STARTRK-2 trials for entrectinib (hereafter called entrectinib trials, N = 54), so we

opted to focus on the adult population. Based on the patient characteristics in the trials, patients were assumed to be 58 years old upon entering the model, with 59% of patients being female.⁶ The included cancers were breast, bile duct (ie, cholangiocarcinoma), colorectal, endometrial, ovarian, pancreatic, and thyroid cancer, as well as neuroendocrine tumor, non-small cell lung cancer, sarcoma, secretory carcinoma of the breast, and secretory carcinoma of the salivary gland.

To model the testing phase of patients, the tumor types were categorized into 3 groups based on a consensus report of Dutch experts, which outlines the envisioned NTRK testing policy in Dutch clinical practice¹¹: (1) tumor types with high NTRK fusion prevalence ($> 90\%$), (2) tumor types with low NTRK fusion prevalence but wild-type TRK protein expression, and (3) tumor types with low NTRK fusion prevalence and no/very little wild-type TRK protein expression (Appendix Table 2 in Supplemental Materials found at <https://doi.org/10.1016/j.jval.2022.08.006>).

The assumed distribution of patients across tumor types in the model was based on the distribution in the pooled data set from the entrectinib trials. As the trials included only NTRK+ patients, the trial distribution of tumor types was combined with the tumor-specific NTRK prevalence to obtain the distribution of tumor types among the total group of patients eligible for NTRK fusion testing (ie, both NTRK+ and NTRK- patients; Appendix Table 2 in Supplemental Materials found at <https://doi.org/10.1016/j.jval.2022.08.006>). Appendix Table 2 in Supplemental Materials found at <https://doi.org/10.1016/j.jval.2022.08.006> also shows the tumor distribution in Dutch clinical practice among our study population, as observed in data from the Hartwig Medical Foundation (HMF), which cover 44 hospitals in The Netherlands. Nonetheless, because we are using the pooled effectiveness estimate from the entrectinib trials, we used the tumor distribution based on the entrectinib trials in our model.

Model Structure

The model consists of a decision tree (reflecting the testing phase) and a microsimulation (reflecting treatment) as shown in Figure 1. We used a lifetime time horizon and a cycle length of 1 month in the microsimulation. Analyses were performed from a Dutch societal perspective and effects and costs were discounted at 1.5% and 4%, respectively.¹²

Decision Tree

Patients enter the model in a decision tree that compares “NTRK fusion testing” with a “no testing” strategy.

For patients receiving testing, the decision tree reflects the period from the decision to test for potential eligibility for entrectinib until the start of treatment. All patients were tested using immunohistochemistry (IHC) and/or next-generation sequencing of RNA (RNA-NGS). Patients with tumor types in groups 1 (high NTRK fusion prevalence) and 2 (wild-type TRK protein expression) were tested only through RNA-NGS. Patients in group 3 (low NTRK fusion prevalence, low TRK protein expression) first received an IHC test to identify those with elevated levels of TRK proteins. Patients who tested positive on the IHC test subsequently underwent confirmatory RNA-NGS testing. The latter strategy may save costs, because IHC tests tend to be much cheaper than RNA-NGS tests. Nevertheless, using IHC as a first screening tool has little added value in groups 1 and 2 because most patients will test positive on the IHC test, either because of the high prevalence of NTRK gene fusions (group 1) or wild-type TRK protein expression (ie, TRK protein expression that is not due to an NTRK fusion) (group 2). See Appendix Figures 1 and 2 in Supplemental Materials found at

Decision tree

The decision tree starts with a yellow node at the beginning of the timeline. It branches into 'TESTING' and 'NO TESTING'. 'TESTING' leads to 'Low fusion, low protein', which then branches into 'Biopsy IHC' and 'No biopsy IHC'. 'Biopsy IHC' leads to 'NTRK + (IHC)' and 'NTRK - (IHC)'. 'NTRK + (IHC)' leads to 'Biopsy NGS' and 'No biopsy NGS'. 'Biopsy NGS' leads to 'Rebiopsy NGS' and 'No rebiopsy NGS'. 'Rebiopsy NGS' leads to 'NTRK + (NGS)' and 'NTRK - (NGS)'. 'NTRK + (NGS)' leads to 'TRK inhibitor' and 'Die'. 'NTRK - (NGS)' leads to 'SOC' and 'Die'. 'No biopsy NGS' leads to 'TN' and 'FN'. 'TN' leads to 'TRK inhibitor' and 'Die'. 'FN' leads to 'SOC' and 'Die'. 'No biopsy IHC' leads to 'NTRK + (IHC)' and 'NTRK - (IHC)'. 'NTRK + (IHC)' leads to 'Biopsy NGS' and 'No biopsy NGS'. 'Biopsy NGS' leads to 'Rebiopsy NGS' and 'No rebiopsy NGS'. 'Rebiopsy NGS' leads to 'NTRK + (NGS)' and 'NTRK - (NGS)'. 'NTRK + (NGS)' leads to 'TRK inhibitor' and 'Die'. 'NTRK - (NGS)' leads to 'SOC' and 'Die'. 'No biopsy NGS' leads to 'TN' and 'FN'. 'TN' leads to 'TRK inhibitor' and 'Die'. 'FN' leads to 'SOC' and 'Die'. 'NTRK - (IHC)' leads to 'TN' and 'FN'. 'TN' leads to 'TRK inhibitor' and 'Die'. 'FN' leads to 'SOC' and 'Die'. 'NO TESTING' leads to 'NTRK + (NGS)' and 'NTRK - (NGS)'. 'NTRK + (NGS)' leads to 'TRK inhibitor' and 'Die'. 'NTRK - (NGS)' leads to 'SOC' and 'Die'.

Microsimulation model

The microsimulation model shows a flow from 'Alive On treatment' to 'Alive Off treatment' or 'Dead'. 'Alive On treatment' leads to 'Alive Off treatment' and 'Dead'. 'Alive Off treatment' leads to 'Dead'.

Timeline: Decision to receive additional treatment, Start of treatment, Death

As mentioned above, the entrectinib trials were single-arm trials. To be able to assess the relative effectiveness of entrectinib compared with SoC, we created a synthetic comparator arm. We used the HMF database, containing data from cancer patients with

Table 1. Input parameters

Input parameter	Value	Source	OWSA	PSA
Testing phase				
NTRK prevalence (by tumor type)	Appendix Table 2 in Supplemental Materials found at https://doi.org/10.1016/j.jval.2022.08.006		± 20%	β
IHC sensitivity (by tumor type)	Appendix Table 3 in Supplemental Materials found at https://doi.org/10.1016/j.jval.2022.08.006		± 20%	β
IHC specificity (by tumor type)	Appendix Table 3 in Supplemental Materials found at https://doi.org/10.1016/j.jval.2022.08.006		± 20%	β
Probability biopsy NGS	0.098	13	± 20%	β
Probability rebiopsy NGS	0.159	13	± 20%	β
Probability biopsy IHC	equal to NGS	Assumption	± 20%	β
Duration waiting time - Biopsy requested until biopsy done - Rebiopsy requested until rebiopsy done - (Re-)biopsy done until IHC test results - (Re-)biopsy done until RNA-NGS results - Final test results available until start of treatment	10.5 days 14 days 3.5 days 10 days 5.5 days	Expert judgment by 3 clinical geneticists and 1 oncologist (mean estimate was used)	± 20%	Multivariate normal
Cost biopsy	Appendix Table 6 in Supplemental Materials found at https://doi.org/10.1016/j.jval.2022.08.006		± 20%	γ*
Cost RNA-NGS	Appendix Table 6 in Supplemental Materials found at https://doi.org/10.1016/j.jval.2022.08.006		± 20%	γ*
Costs IHC	Appendix Table 6 in Supplemental Materials found at https://doi.org/10.1016/j.jval.2022.08.006		± 20%	γ*
Treatment phase				
Starting age in years	58	6	± 20%	Normal*
Proportion females	0.59	6	± 20%	β
OS NTRK– (by tumor type)	Appendix Table 5 in Supplemental Materials found at https://doi.org/10.1016/j.jval.2022.08.006	HMF	-	Asymptomatic normal ¹⁴
TTD NTRK– (by tumor type)	Appendix Table 5 in Supplemental Materials found at https://doi.org/10.1016/j.jval.2022.08.006	HMF	-	Asymptomatic normal ¹⁴
OS entrectinib	exponential	Roche	-	Multivariate normal
TTDisc entrectinib	exponential	Roche	-	Multivariate normal
HR NTRK+ OS adjusted	1.44	HMF	95% CI	Lognormal
HR NTRK+ TTDisc adjusted	1.37	HMF	95% CI	Lognormal
Tumor distribution [†]	Appendix Table 2 in Supplemental Materials found at https://doi.org/10.1016/j.jval.2022.08.006	6	-	-
Utilities – time to death	-	Versteegh et al	± 20%	Multivariate normal

continued on next page

Table 1. Continued

Input parameter	Value	Source	OWSA	PSA
Informal care use	-	Santi et al (submitted)	± 20%	Multivariate normal
Treatment costs SoC (per month)	Appendix Table 7 in Supplemental Materials found at https://doi.org/10.1016/j.jval.2022.08.006		± 20%	γ^*
Treatment costs entrectinib (per month)	€5913	Roche	± 20%	γ^*
Adverse event costs (per event)	Appendix Table 8 in Supplemental Materials found at https://doi.org/10.1016/j.jval.2022.08.006		± 20%	γ^*
Informal care costs (per hour)	€14.77	12	± 20%	γ^*
Hospital costs related and unrelated to cancer per year (not year before death) at age 58	€4453	PAID	± 20%	-
Hospital costs related and unrelated to cancer in the year before death at age 58	€58 064	PAID	± 20%	-

CI indicates confidence interval; HMF, Hartwig Medical Foundation; HR, hazard ratio; IHC, immunohistochemistry; NGS, next-generation sequencing; NTRK–, NTRK negative; NTRK+, NTRK positive; OS, overall survival; OWSA, one-way sensitivity analysis; PAID, Practical Application to Include Disease Costs; PSA, probabilistic sensitivity analysis; SoC, standard of care; TTD, time-to-treatment-discontinuation.

*Assumption SD is 10% of the mean.

†Calculated using the following formula: tumor type distribution before test = tumor type distribution in entrectinib trials / NTRK prevalence, rescaled to proportions.

locally advanced or metastatic tumors who had genomic profiling between 2012 and 2020 (CPCT-02 study – NCT01855477).^{10,17} Patients were eligible for enrolment in the CPCT-02 study if (1) their age \geq 18 years, (2) they had a locally advanced or metastatic solid tumor, (3) they had an indication for new line of systemic treatment with registered anticancer agents, and (4) performing a tumor biopsy was safe according to the treating physician.¹⁰ There are no publicly available patient-level data from the entrectinib trials, meaning that statistical methods to match the study populations from the HMF database and the entrectinib trials (eg, propensity score matching) could not be applied.¹⁸

OS and TTD in NTRK– patients receiving SoC were based on data from 1596 NTRK– patients who received SoC (Appendix Fig. 3 in Supplemental Materials found at <https://doi.org/10.1016/j.jval.2022.08.006>). For each tumor type, parametric distributions were fitted to data on OS and TTD, using the Akaike Information Criterion to determine the parametric distribution with the best fit (Appendix Table 5 in Supplemental Materials found at <https://doi.org/10.1016/j.jval.2022.08.006>). Subsequently, tumor-specific monthly transition probabilities to “death” and “discontinuation of treatment” were extracted to be used in the model. The transition probabilities to discontinuation of treatment (entrectinib or SoC) and death were estimated independently.

To estimate transition probabilities to “death” and “treatment discontinuation” in NTRK+ patients receiving SoC, we applied a hazard ratio (HR) reflecting the prognostic value of carrying an oncogenic NTRK gene fusion to the OS and TTD estimates for NTRK– patients.

Prognostic value of NTRK fusions

We used the HMF database to estimate the prognostic value of an NTRK fusion. Patients in the database were classified into 2 cohorts: NTRK+ and NTRK–. A subgroup of the NTRK– cohort was matched to the patients in the NTRK+ cohort, using the optimal matching method.¹⁹ Within each tumor type, patients

were matched based on baseline patient characteristics, including age, sex, number of previous lines of therapy, and year of biopsy, using a ratio of 1:4 (NTRK+:NTRK–). A total of 24 NTRK+ patients were matched with 96 NTRK– patients, with a successful covariate balance between the 2 groups. OS and TTD analyses were performed using the Kaplan-Meier method (Fig. 2) and Cox regressions, with the date of the first postbiopsy treatment as the index date and age, sex, and number of previous lines of treatment as covariates. If a subject was known to be alive before the cutoff date, the subject was censored at the last known alive date. A HR of 1.44 (95% confidence interval [CI] 0.81–2.56) was estimated for OS among NTRK+ patients and 1.37 (95% CI 0.86–2.18) for TTD (Appendix Fig. 4 in Supplemental Materials found at <https://doi.org/10.1016/j.jval.2022.08.006>). This suggests that in SoC, NTRK+ patients have worse prognosis than NTRK– patients. We assumed that the HRs were constant across tumor types.

Costs


All costs are expressed in 2020 euros.

Decision tree

For each tumor type, we calculated the mean costs of biopsy, IHC testing, and RNA-NGS testing. Cost variations across tumor types were caused by differences in the type of biopsy needed (eg, more resources are needed for a lung biopsy than for a skin biopsy) and differences in price-setting among the main treatment centers for the various tumor types. The cost of an IHC test varied between €67 and €328 and the cost of an RNA-NGS test varied between €870 and €2137 (Appendix Table 6 in Supplemental Materials found at <https://doi.org/10.1016/j.jval.2022.08.006>).

Microsimulation model

We included the costs of oncology drugs and their administration (including admissions at a day care unit if needed). For

Figure 2. Prognostic value of NTRK fusion on OS.

CI indicates confidence interval; HR, hazard ratio; OS, overall survival; pos, positive.

entrectinib, the monthly treatment cost was €5913 for all tumor types. The treatment costs for the various SoC treatments were combined into an average cost per tumor type with a maximum of €5501 per month for colorectal cancer (Appendix Table 7 in Supplemental Materials found at <https://doi.org/10.1016/j.jval.2022.08.006>). In addition, costs for the treatment of adverse events (AEs) were included. Because of a lack of data on AEs with entrectinib, we assumed that the occurrence of AEs when receiving entrectinib was equal to AE occurrence in SoC. The tumor-specific prevalence of AEs was multiplied by the cost of treating the AEs, with input values for both variables based on estimates from the literature (Appendix Table 8 in Supplemental Materials found at <https://doi.org/10.1016/j.jval.2022.08.006>). The costs of AEs were applied for patients in the “alive and on treatment” health state.

As per the Dutch health technology assessment (HTA) guideline, we also included the costs of non-hospital care related to cancer (eg, psychosocial care, home nursing, and general practitioner check-ups), as well as hospital and non-hospital care unrelated to cancer. We obtained age-dependent cost estimates through the “Practical Application to Include Disease Costs” tool.²⁰ At baseline, the total annual other healthcare costs derived from “Practical Application to Include Disease Costs” (ie, excluding drugs, administration of drugs, and AE costs) were €4453. In the year before death, they increased to €58 064.

Regarding societal costs, we included informal care costs but no productivity costs. Because the patients in our study population are in an advanced stage of cancer, we assumed that they are already out of the workforce upon entering our model in order that no additional productivity losses occur (ie, the friction period has already passed).²¹ Informal care estimates were provided by a regression analysis that estimated the impact of proximity to death on the use of informal care, correcting for covariates, such as age and sex, using the Survey of Health, Ageing, and Retirement in Europe data.²² The analysis provided the probability of using

informal care, as well as the amount of hours used. We valued informal care at the standard rate recommended by the national HTA guideline (€14.77 per hour).¹²

Utilities

Based on previous research findings that the quality of life decreases as patients approach death, we incorporated patient utility in our model as a function of time to death.²³ We used as model inputs the regression coefficients from a study that estimated the relationship between proximity to death and utility.²³ In the study, proximity-to-death values were obtained using OS functions and utility values based on the SF-6D method were used. Age and sex were also included as covariates, allowing us to include age- and sex-specific patient utility in our model.²³ We included the utility during the testing pathway, as well as during treatment.

Analyses

The decision tree and the microsimulation model were programmed in line with the Decision Analysis in R for Technologies in Health (DARTH) modeling framework in R 3.6.1 using RStudio 1.2.1335 (RStudio, Boston, MA).²⁴⁻²⁶

Base-Case Analysis

The base-case analysis reflects the cost-effectiveness of testing cancer patients with locally advanced or metastatic solid tumors for NTRK gene fusions and subsequently treating NTRK+ patients with entrectinib and NTRK- patients with SoC compared with no testing and treating all patients with SoC. The incremental net monetary benefit (INMB) was calculated using a cost-effectiveness threshold of €80 000 per quality-adjusted life-year (QALY), which is the recommended value by Dutch HTA guidelines, given the calculated disease burden.^{27,28} We opted to simulate 5000 patients after evaluating the stability of model outcomes at various

cohort sizes (Appendix Fig. 5 in Supplemental Materials found at <https://doi.org/10.1016/j.jval.2022.08.006>).

Scenario Analyses

The first scenario analysis aimed to evaluate the impact of testing costs on the cost-effectiveness of implementing entrectinib by excluding the costs of NTRK testing. The second scenario excluded the costs, as well as the health effects resulting from the testing pathway (ie, the probability to die and the QALYs during the waiting time). This scenario reflects a setting in which the patients in our study population would not need any NTRK testing, for example, because RNA-NGS testing is part of standard practice and has already been done at an earlier stage. The third scenario takes the approach that is common in economic evaluations of targeted treatments, which is to only include the patients who carry the targeted genetic marker and to disregard the costs and health effects of testing.⁹ We also performed the base-case analysis from a healthcare perspective.

As mentioned in section "Study Population", a stratified testing protocol (first IHC, then RNA-NGS for patients who test positive) is used for patients in group 3 (tumor types with low NTRK fusion prevalence and no/little wild-type TRK protein expression), which is the largest group. Nevertheless, RNA-NGS has much better test sensitivity and specificity than IHC.²⁹ If costs were not considered, providing RNA-NGS to all patients would therefore likely be preferred. RNA-NGS is a relatively new technology and in many settings is not yet part of standard care. As RNA-NGS becomes more widespread and perhaps further technological improvements are made to achieve efficiency gains, its cost may decrease. We therefore investigated at what price of RNA-NGS the provision of RNA-NGS to all patients would become cost-effective (ie, renders 0 INMB).

Sensitivity Analyses

Parameter uncertainty was tested using univariate sensitivity analysis and probabilistic sensitivity analysis (PSA). In the univariate analysis, model parameters were independently varied over the extremities of the 95% CI and, when this was not available, by a 20% increase/decrease from the parameter value in the base case. In the PSA, all parameters were varied simultaneously according to predefined distributions (Table 1^{6,12-14}). The model was run with 1000 iterations while sampling 1000 patients.

Budget Impact Analysis

A 5-year budget impact was estimated by multiplying the annual incremental healthcare costs per patient tested in the first 5 years with the expected number of patients tested. The number of patients tested each year in The Netherlands was determined by multiplying the number of expected NTRK+ patients (N = 90)^{30,31} by the number of patients that need to be tested to identify 1 NTRK+ patient (as derived from our model results).

Results

Base-Case Analysis

The results of the base-case analysis are presented in Table 2 and Appendix Table 9 in Supplemental Materials found at <https://doi.org/10.1016/j.jval.2022.08.006>. They showed that testing for NTRK fusions and treatment with entrectinib in NTRK+ patients and SoC in NTRK- patients was associated with a QALY gain of 0.0043 at an increased cost of €732 per patient as compared with no testing and SoC for all patients. The incremental cost-effectiveness ratio (ICER) was €169 957/QALY and the INMB was -€388.

Table 2. Base-case analysis and scenario analyses with or without testing costs and consequences

Base-case or scenario analysis	Comparator: base case, scenario 1 and scenario 2	Intervention: base case (including testing costs and consequences)	Intervention: scenario 1 (excluding testing costs)	Intervention: scenario 2 (excluding test costs and consequences)	Comparator: scenario 3	Intervention: scenario 3
Patient population	NTRK+ and NTRK- patients	NTRK+ and NTRK- patients	NTRK+ and NTRK- patients	NTRK+ and NTRK- patients	NTRK+ patients	NTRK+ patients
Treatment	No NTRK testing, SoC for all patients	NTRK testing, entrectinib for NTRK+ patients, SoC for NTRK- patients	NTRK testing, entrectinib for NTRK+ patients, SoC for NTRK- patients	NTRK testing, entrectinib for NTRK+ patients, SoC for NTRK- patients	SoC	Entrectinib
Total costs (in €)	74 071	74 803	74 227	74 244	71 131	129 240
Total effects (in QALYs)	0.985	0.989	0.989	0.989	0.679	2.186
Incremental costs (in €)		732	156	172		58 109
Incremental effects (in QALYs)		0.0043	0.0043	0.0045		1.507
INMB (in €)		-388	188	161		69 945
ICER (in €/QALY)		169 957	36 290	38 658		38 563
Proportion of patients with entrectinib treatment (%)	0	0.28	0.28	0.30	0	100

ICER indicates incremental cost-effectiveness ratio; INMB, incremental net monetary benefit; NTRK-, NTRK negative; NTRK+, NTRK positive; QALY, quality-adjusted life-year; SoC, standard of care.

Scenario Analyses

Table 2 shows the results of scenario analyses with or without testing costs and consequences. Not including testing costs had a large impact, whereas not including mortality and QALYs during waiting time only had a minor impact on the cost-effectiveness results. In the third scenario analysis, in which only NTRK+ patients were considered and the testing phase was disregarded, the ICER of entrectinib versus SoC was €38 563/QALY. The results from the healthcare perspective were similar to the results in the base case (Appendix Table 10 in Supplemental Materials found at <https://doi.org/10.1016/j.jval.2022.08.006>). We estimated that the cost of RNA-NGS would have to be reduced by 90%, to €186, before testing all eligible patients with RNA-NGS would be cost-effective.

Sensitivity Analyses

The results of the univariate analysis are presented in Appendix Figure 6 in Supplemental Materials found at <https://doi.org/10.1016/j.jval.2022.08.006>. The specificity of the IHC test had the largest impact on the INMB, followed by the cost of IHC testing and the HR for the prognostic value of NTRK fusions on OS. Improvements in the specificity of IHC tests (ie, fewer false positives), decreases in the costs for IHC tests, or increases in the HR for the prognostic value of NTRK fusions on OS (ie, worse prognosis of NTRK+) led to increased INMB.

The results of the PSA are presented in the cost-effectiveness plane in Figure 3 and the cost-effectiveness acceptability curve in Appendix Figure 7 in Supplemental Materials found at <https://doi.org/10.1016/j.jval.2022.08.006>. Mean incremental costs of all iterations were €696 and mean incremental effects were 0.0037, resulting in an INMB of –€399 and an ICER of €187 681/QALY. The probability of “Testing + entrectinib/SoC” being cost-effective compared with “No testing + SoC” was 0.2% at a threshold of €80 000 per QALY.

Budget Impact Analysis

The proportion of patients who received entrectinib from all patients that were tested in our model was 0.28%. Combined with the expected number of patients treated with entrectinib (N = 90), this results in 31630 patients being eligible for NTRK testing in

The Netherlands annually. The 5-year incremental budget impact of testing and treatment with entrectinib in NTRK+ patients was €93 million, with testing costs making up 82% of the total budget impact (Table 3).

Discussion

Summary of Results


Our results showed that incorporating the consequences of NTRK testing has a large impact on the cost-effectiveness of implementing the histology-independent treatment entrectinib to the extent that it would alter the reimbursement decision. Excluding the costs and health effects associated with NTRK testing reduced the ICER from €169 957/QALY to €38 563/QALY. The difference is primarily owing to the rarity of NTRK gene fusions, meaning that a large number of patients need to be tested to identify the few patients that are eligible for entrectinib treatment (0.29%). This means that few patients experience QALY gains from NTRK testing (only the NTRK+ patients who experience better health outcomes with entrectinib treatment than SoC), while additional costs (for the tests) are added for all patients.

The estimated five-year budget impact of testing (€93 million) was based on the testing capacity that is needed when all eligible patients are tested. Nevertheless, the testing practice for NTRK fusion-positive cancers in The Netherlands is still in development.¹¹ Hence, fewer tests may be conducted in the earlier years of the implementation of NTRK testing and subsequent treatment, which would make the actual budget impact lower than what we estimated.

Strengths

When the introduction of a new drug requires a new diagnostic test to identify the target population, HTA bodies should be fully informed about the cost-effectiveness of the test-treatment combination.⁹ Therefore, in this study we considered the entire patient population that would be affected (ie, all patients eligible for NTRK testing) and calculated the (downstream) costs and benefits of NTRK testing and subsequent treatment for patients who tested NTRK+ and NTRK–.

Figure 3. Cost-effectiveness plane of “Testing + entrectinib/SoC” versus “No testing + SoC.”



QALY indicates quality-adjusted life-year; SoC, standard of care.

Table 3. Results of budget impact analysis (in €)

Entrectinib	Annual test costs	Annual healthcare costs	Annual total costs
Year 1	15 249 931	810 033 052	825 282 982
Year 2	15 249 931	1 317 036 538	1 332 286 468
Year 3	15 249 931	1 661 785 334	1 677 035 265
Year 4	15 249 931	1 800 859 346	1 816 109 277
Year 5	15 249 931	1 827 257 744	1 842 507 675
SoC	Annual test costs	Annual healthcare costs	Annual total costs
Year 1	-	807 719 086	807 719 086
Year 2	-	1 314 096 035	1 314 096 035
Year 3	-	1 658 446 610	1 658 446 610
Year 4	-	1 796 990 043	1 796 990 043
Year 5	-	1 822 922 722	1 822 922 722
Incremental	Annual test costs	Annual healthcare costs	Annual total costs
Year 1	15 249 931	2 313 966	17 563 896
Year 2	15 249 931	2 940 503	18 190 433
Year 3	15 249 931	3 338 724	18 588 655
Year 4	15 249 931	3 869 303	19 119 234
Year 5	15 249 931	4 335 022	19 584 953

SoC indicates standard of care.

Entrectinib was conditionally approved by the EMA and the Food and Drug Administration based on small single-arm basket trials showing a durable response and longer survival.³² The single-arm nature of the trial data poses a great challenge to HTA bodies making decisions based on comparative effectiveness and cost-effectiveness. Briggs et al,³³ in a study about the estimation of the counterfactual for tumor-agnostic treatments with only single-arm trial data available, described 3 options: (1) historical controls reported in the literature, (2) previous line of therapy in intervention patients of basket trials, or (3) nonresponders in basket trials. As we did not have access to individual patient data from the entrectinib trials, we had to rely on historical data. Nevertheless, unlike Briggs et al,³³ we used genetic data to adjust the historical data for the prognostic value of carrying oncogenic NTRK gene fusions; therewith, likely increasing the accuracy of the estimated comparative effectiveness. Indeed, 2 other studies that estimated an HR for the OS of NTRK+ patients found comparable values (1.44³⁴ and 1.6,³⁵ respectively) to the value we estimated (1.44). With our approach, we illustrated how a database with genomic and clinical data can be used to match patients with and without a specific genetic marker and to estimate HRs (for OS and TTD) for the patients with the marker. A similar approach could be used to estimate HRs for patients with different genetic markers. Although residual confounding cannot be ruled out in our analysis because of the small sample size, lack of matching patients on all relevant covariates (eg, Eastern Cooperative Oncology Group performance status), and other limitations, more robust statistical estimation may be achieved with larger sample sizes (eg, for genetic markers that are less rare) and with data for more (clinical) variables. Nonetheless, RCT data are preferred over historical data for the assessment of comparative effectiveness, and the kind of matching exercise we conducted should only be done when no RCT data are available.

Most health economic models for cancer treatments include the health states “progression-free” and “progression,” which require data on progression status in addition to OS. Because

appropriate data on progression for patients receiving SoC were not available, we instead used a regression model that estimated the impact of proximity to death on the utility of cancer patients. Although this is not a conventional approach, we believe that linking utility to proximity to death is likely more appropriate (ie, closer to the lived experience of cancer patients) than using singular utility values for progression-free and progressed patients. We therefore consider this approach a strength of our study.

Limitations

Although we estimated tumor-specific effectiveness of SoC, we had to assume that the effectiveness of entrectinib was constant across tumor types, as the small number of observations per tumor type in the entrectinib trials precluded the estimation of reliable tumor-specific effectiveness. Nevertheless, Murphy et al⁷ showed heterogeneity in clinical effects across histologies that should be accounted for, for example, by using Bayesian hierarchical models. We present a single ICER in this study, implying an all-or-nothing decision regarding the reimbursement of entrectinib for NTRK+ patients, yet it might be more appropriate to differentiate between tumor types. That is, even when histology-independent therapies receive marketing authorization for all histologies, reimbursement might be warranted only for a subset of histologies because of heterogeneity in the treatment effect. If indeed there is heterogeneity in effectiveness across tumor types, the single ICER estimate we estimated based on the patient population in the entrectinib trials may be biased because the proportional distribution of tumor types in the trial is not fully representative of the distribution of tumor types among eligible patients in clinical practice. Additionally, because of the small number of NTRK+ observations per tumor type in the HMF database, we had to assume that the prognostic value of NTRK fusions was constant across tumor types.

Note that although our analysis does not account for heterogeneity in the treatment effect of entrectinib and the HRs for NTRK

gene fusions across histologies, it does account for differences in the treatment costs and effects of SoC (both TTD and OS) across tumor types and uses tumor-specific estimates of testing costs.

We attempted to construct an appropriate comparator arm by estimating outcomes on a similar target population and adjusting for the prognostic value of NTRK fusions. That is, we only considered patients with locally advanced or metastatic disease who had received at least one previous line of treatment, in line with the patient population in the entrectinib trials. We also only considered patients who had one of the tumor types included in the entrectinib trials. We attempted to create a subgroup of HMF patients with a similar average age and percent of females as those in the entrectinib trials, but the subgroup was too small to enable reliable statistical estimation. Despite our efforts, we cannot be sure that the patient populations in the respective arms are fully comparable because we did not have access to patient-level data from the entrectinib trials. Moreover, there may be unobserved differences between the patient populations.

Implications for Decision Making

Considering the impact of testing costs on the cost-effectiveness of entrectinib, payers should focus on policies supporting a reduction in the costs of testing. Furthermore, the uncertainty around the effectiveness of entrectinib leaves HTA bodies with 2 main choices: wait for more (tumor type-specific) evidence or provide coverage through a managed entry agreement between the pharmaceutical company and the healthcare payer. Considering the rarity of NTRK fusions, stronger evidence will likely not become available soon, meaning that waiting for more evidence would leave patients without access to entrectinib for a long time. Therefore, despite concerns about the feasibility of discontinuing reimbursement for medicines once further evidence does not demonstrate their (cost-)effectiveness,^{36,37} a managed entry agreement, for example, in the form of coverage with evidence development, may be the best option.³⁸ Through coverage with evidence development agreements, treatments are temporarily reimbursed while further evidence is collected. After a specified period, the cost-effectiveness of the treatment is re-evaluated using the additional data. Ideally, data on final outcomes, such as survival and quality of life, are collected. When the follow-up time is limited, surrogate outcomes, such as progression-free survival and tumor response rates, are also used. Although, it is of note that surrogate outcomes are not necessarily predictive of final outcomes.³⁹

In addition, we found that the uncertainty around the HR for NTRK gene fusions on OS has a large effect on cost-effectiveness outcomes. Because of the low prevalence of NTRK fusions, larger genomic databases (paired with clinical information) are needed to gather sufficiently large numbers of NTRK+ patients to obtain statistically significant results on the prognostic value of NTRK fusions. It may therefore be valuable, for patients with other rare genetic markers as well, if decision-makers invest in expanding genomic data collection, possibly including policies that encourage the linking of existing databases. Finally, although our analysis has focused on entrectinib, larotrectinib has a similar target population, mechanism of action, and price-setting in The Netherlands. Our finding that the introduction of TRK inhibitor entrectinib appears to not be cost-effective when considering the consequences of introducing NTRK testing but might be cost-effective if the relevant tests would become standard practice, likely also applies for larotrectinib. A recent study that investigated the cost-effectiveness of larotrectinib in The Netherlands and focused on NTRK+ patients without considering the testing phase, comparable with our scenario 3, found an ICER of €41 424/

QALY, which is similar to the ICER of €38 563/QALY we estimated for entrectinib in scenario 3.⁴⁰

Conclusions

In conclusion, with the currently available evidence, it seems that entrectinib is not cost-effective compared with SoC. Nevertheless, if genetic testing of cancer patients (including RNA-NGS panels that can identify NTRK gene fusions) becomes standard practice, entrectinib may be cost-effective. Nonetheless, our study results are very uncertain because of data limitations.

Supplemental Material

Supplementary data associated with this article can be found in the online version at <https://doi.org/10.1016/j.jval.2022.08.006>.

Article and Author Information

Accepted for Publication: August 12, 2022

Published Online: October 11, 2022

doi: <https://doi.org/10.1016/j.jval.2022.08.006>

Author Affiliations: Institute for Medical Technology Assessment, Erasmus University Rotterdam, Rotterdam, The Netherlands (Huygens, Vellekoop, Versteegh, Santi, Mölken); Syreon Research Institute, Budapest, Hungary (Szilberhorn, Zelei, Nagy); Health Economics Research Centre, University of Oxford, Oxford, England, United Kingdom (Tsiachristas, Koleva-Kolarova, Wordsworth); Erasmus School of Health Policy & Management, Erasmus University Rotterdam, Rotterdam, The Netherlands (Mölken).

Correspondence: Maureen Rutten-van Mölken, Erasmus School of Health Policy & Management, Erasmus University Rotterdam, Burgemeester Oudlaan 50, 3062 PA, Rotterdam, The Netherlands. Email: m.rutten@eshpm.eur.nl

Author Contributions: *Concept and design:* Huygens, Vellekoop, Versteegh, Szilberhorn, Zelei, Nagy, Tsiachristas, Koleva-Kolarova, Wordsworth, Rutten-van Mölken

Acquisition of data: Huygens, Vellekoop, Versteegh, Santi, Rutten-van Mölken
Analysis and interpretation of data: Huygens, Vellekoop, Versteegh, Santi, Tsiachristas, Rutten-van Mölken

Drafting of the manuscript: Huygens, Vellekoop, Tsiachristas, Rutten-van Mölken
Critical revision of the paper for important intellectual content: Huygens, Vellekoop, Versteegh, Santi, Szilberhorn, Zelei, Nagy, Tsiachristas, Koleva-Kolarova, Wordsworth, Rutten-van Mölken

Statistical analysis: Huygens, Vellekoop, Versteegh, Santi

Provision of study materials or patients: Huygens, Vellekoop, Versteegh, Santi, Rutten-van Mölken

Obtaining funding: Versteegh, Nagy, Tsiachristas, Wordsworth, Rutten-van Mölken

Administrative, technical, or logistic support: Huygens, Vellekoop, Versteegh, Rutten-van Mölken

Supervision: Huygens, Versteegh, Rutten-van Mölken

Conflict of Interest Disclosures: Dr Versteegh reported receiving personal fees from EuroQoL research foundation, grants from Boehringer Ingelheim, Merck KgCA, BeiGene, Novartis Netherlands, Biogen Netherlands, Amgen, and AstraZeneca, outside the submitted work. No other conflicts were reported.

Funding/Support: The HECoPerMed project has received funding from the European Union's Horizon 2020 research and innovation program under grant agreement number 824997.

Role of the Funder/Sponsor: The funder had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, and approval of the manuscript; nor the decision to submit the manuscript for publication.

Acknowledgment: The authors express their gratitude to the oncologists and clinical geneticists who provided insights into clinical practice for oncology patients and explained various details about the diagnostic methods for identifying NTRK gene fusions. We would like to acknowledge all other members of the HecoPerMed consortium for their contribution to the project: Ballensiefen W, Walgenbach M and Schartinger D.

REFERENCES

- Assessment report VITRAKVI international non-proprietary name: larotrectinib. European Medicines Agency. https://www.ema.europa.eu/en/documents/assessment-report/vitrakvi-epar-public-assessment-report_en.pdf. Accessed January 20, 2022.
- Rozlytrec (entrectinib). European Medicines Agency. <https://www.ema.europa.eu/en/medicines/human/EPAR/rozlytrec>. Accessed January 20, 2022.
- Rodes Sanchez M, Henderson N, Steuten L. Bridging the gap: pathways for regulatory and health technology assessment of histology independent therapies. <https://www.ohe.org/publications/bridging-gap-pathways-regulatory-and-health-technology-assessment-histology-independent>. Accessed January 20, 2022.
- Drilon A, Laetsch TW, Kummar S, et al. Efficacy of larotrectinib in TRK fusion-positive cancers in adults and children. *N Engl J Med*. 2018;378(8):731–739.
- Hong DS, DuBois SG, Kummar S, et al. Larotrectinib in patients with TRK fusion-positive solid tumors: a pooled analysis of three phase 1/2 clinical trials. *Lancet Oncol*. 2020;21(4):531–540.
- Doebele RC, Drilon A, Paz-Ares L, et al. Entrectinib in patients with advanced or metastatic NTRK fusion-positive solid tumors: integrated analysis of three phase 1–2 trials. *Lancet Oncol*. 2020;21(2):271–282.
- Murphy P, Claxton L, Hodgson R, et al. Exploring heterogeneity in histology-independent technologies and the implications for cost-effectiveness. *Med Decis Mak*. 2021;41(2):165–178.
- van Rosmalen J, Dejardin D, van Norden Y, Löwenberg B, Lesaffre E. Including historical data in the analysis of clinical trials: is it worth the effort? *Stat Methods Med Res*. 2018;27(10):3167–3182.
- Vellekoop H, Huygens S, Versteegh M, et al. Guidance for the harmonisation and improvement of economic evaluations of personalised medicine. *Pharmacoeconomics*. 2021;39(7):771–788.
- Priestley P, Baber J, Lolkema MP, et al. Pan-cancer whole-genome analyses of metastatic solid tumors. *Nature*. 2019;575(7781):210–216.
- Aerts JGJV, Dinjens WNM, Evers MPJ, et al. Consensus diagnose en behandeling van ntrk-genfusie gerelateerde solide tumoren. https://nfk.nl/media/1/Consensus_Rapport_Diagnostiek_en_Behandeling_van_NTRK-Genfusie_Gerelateerde_Solide_Tumoren_14022020.pdf. Accessed January 20, 2022.
- Guideline for Conducting Economic Evaluations in Healthcare [in Dutch: Richtlijn Voor Het Uitvoeren van Economische Evaluaties in de Gezondheidszorg]. Zorginstituut Nederland (ZIN). <https://www.zorginstituutnederland.nl/publicaties/publicatie/2016/02/29/richtlijn-voor-het-uitvoeren-van-economische-evaluaties-in-de-gezondheidszorg>. Accessed January 20, 2022.
- Bins S, Cirkel GA, Gadellaa-Van Hooijdonk CG, et al. Implementation of a multicenter biobanking collaboration for next-generation sequencing-based biomarker discovery based on fresh frozen pretreatment tumor tissue biopsies. *Oncologist*. 2017;22(1):33–40.
- Jackson C, Metcalfe P, Amdahl J, Warkentin MT, Kunzmann K. Package “Flexsurv”: Flexible Parametric Survival and Multi-state Models. CRAN. <https://cran.r-project.org/web/packages/flexsurv/flexsurv.pdf>. Accessed January 20, 2022.
- Williams HL, Walsh K, Diamond A, Oniscu A, Deans ZC. Validation of the OncoPrint™ focus panel for next-generation sequencing of clinical tumor samples. *Virchows Arch*. 2018;473(4):489–503.
- Entrectinib for treating NTRK fusion-positive solid tumors. National Institute for Health and Care Excellence (NICE). <https://www.nice.org.uk/guidance/ta644/evidence/committee-papers-pdf-8831613997>. Accessed January 20, 2022.
- van Riet J, van de Werken HJC, Cuppen E, et al. The genomic landscape of 85 advanced neuroendocrine neoplasms reveals subtype-heterogeneity and potential therapeutic targets. *Nat Commun*. 2021;12(1):4612.
- Thorlund K, Dron L, Park JJH, Mills EJ. Synthetic and external controls in clinical trials – a primer for researchers. *Clin Epidemiol*. 2020;12:457–467.
- Stuart EA. Matching methods for causal inference: a review and a look forward. *Stat Sci*. 2010;25(1):1–21.
- van Baal P, Perry-Duxbury M, Bakx P, Versteegh M, van Doorslaer E, Brouwer W. A cost-effectiveness threshold based on the marginal returns of cardiovascular hospital spending. *Health Econ*. 2019;28(1):87–100.
- Koopmanschap MA, Rutten FF, van Ineveld BM, van Roijen L. The friction cost method for measuring indirect costs of disease. *J Health Econ*. 1995;14(2):171–189.
- Hoogendoorn M, de Groot S, Corroramos I, et al. Deliverable 6.1 Report on Most and Least Cost-Effective SMI's. Appendix 6. – Estimating the Broader Societal Impact of SMIs. Deliverable to the European Commission for COMPAR-EU (Grant Nr: 754936); 2021. To be available at: <https://cordis.europa.eu/project/id/754936>.
- Versteegh M, van der Helm I, Mokri H, Oerlemans S, Blommestein H, van Baal P. Estimating quality of life decrements in oncology using time to death [Published online July 6, 2022]. *Value Health*. <https://doi.org/10.1016/j.jval.2022.06.002>
- Jalal H, Pechlivanoglou P, Krijkamp E, Alarid-Escudero F, Enns E, Hunink MGM. An overview of R in health decision sciences. *Med Decis Mak*. 2017;37(7):735–746.
- Krijkamp EM, Alarid-Escudero F, Enns EA, Jalal HJ, Hunink MGM, Pechlivanoglou P. Microsimulation modeling for health decision sciences using R: a tutorial. *Med Decis Making*. 2018;38(3):400–422.
- Krijkamp EM, Alarid-Escudero F, Enns EA, et al. A multidimensional array representation of state-transition model dynamics. *Med Decis Making*. 2020;40(2):242–248.
- Kosteneffectiviteit in de Praktijk. Zorginstituut Nederland. <https://www.zorginstituutnederland.nl/publicaties/rapport/2015/06/26/kosteneffectiviteit-in-de-praktijk>. Accessed January 20, 2022.
- Versteegh MM, Ramos IC, Buyukkaramikli NC, Ansari-pour A, Reckers-Droog VT, Brouwer WBF. Severity-adjusted probability of being cost effective. *Pharmacoeconomics*. 2019;37(9):1155–1163.
- Koopman B, Kuijpers CCHJ, Groen HJM, et al. Detection of NTRK fusions and TRK expression and performance of pan-TRK immunohistochemistry in routine diagnostics: results from a nationwide community-based cohort. *Diagnostics*. 2022;12(3):1–15.
- Forsythe A, Zhang W, Phillip Strauss U, Fellous M, Korei M, Keating K. A systematic review and meta-analysis of neurotrophic tyrosine receptor kinase gene fusion frequencies in solid tumors. *Ther Adv Med Oncol*. 2020;12:1758835920975613.
- Advies potentiële kandidaat voor voorwaardelijke toelating van larotrectinib (Vitrakvi®) bij solide tumoren met een ntrk-genfusie (Procedure: Weesgeene-middelen, Conditionals En Exceptionals). Zorginstituut Nederland (ZIN). <https://www.zorginstituutnederland.nl/publicaties/adviezen/2021/03/10/advies-over-kandidaat-voorvoorwaardelijke-toelating-larotrectinib-vitrakvi>. Accessed January 20, 2022.
- Ardini E, Siena S. Entrectinib approval by EMA reinforces options for ROS1 and tumor agnostic NTRK targeted cancer therapies. *ESMO Open*. 2020;5(5):e000867.
- Briggs A, Wehler B, Gaultney JG, et al. Comparison of alternative methods to assess the cost-effectiveness of tumor-agnostic therapies: a triangulation approach using larotrectinib as a case study. *Value Health*. 2022;25(6):1002–1009.
- Bazhenova L, Lokker A, Snider J, et al. TRK fusion cancer: patient characteristics and survival analysis in the real-world setting. *Target Oncol*. 2021;16(3):389–399.
- Demetri GD, Peters S, Hibbar DP, et al. Characteristics and outcomes of patients (pts) with NTRK fusion-positive (NTRK+) metastatic/locally advanced (LA) solid tumors receiving non-TRK inhibitor (TRKi) standard of care (SoC), and prognostic value of NTRK fusions in clinical practice. *Ann Oncol*. 2021;32(suppl 5):S399.
- van de Wetering EJ, van Exel J, Brouwer WB. The challenge of conditional reimbursement: stopping reimbursement can be more difficult than not starting in the first place. *Value Health*. 2017;20(1):118–125.
- McCabe C, Chilcott J, Claxton K, et al. Continuing the multiple sclerosis risk sharing scheme is unjustified. *BMJ*. 2010;340:c1786.
- Wenzl M, Chapman S. Performance-Based Managed Entry Agreements for New Medicines in OECD Countries and EU Member States: How They Work and Possible Improvements Going Forward. OECD. <https://www.oecd.org/health/health-systems/pharma-managed-entry-agreements.htm>. Accessed January 20, 2022.
- Gyawali B, Hey SP, Kesselheim AS. Assessment of the clinical benefit of cancer drugs receiving accelerated approval. *JAMA Intern Med*. 2019;179(7):906–913.
- Michels RE, Arteaga CH, Peters ML, Kapiteijn E, Van Herpen CML, Krol M. Economic evaluation of a tumor-agnostic therapy: Dutch economic value of larotrectinib in TRK fusion-positive cancers. *Appl Health Econ Health Policy*. 2022;20(5):717–729.