

Catalytic B–N Dehydrogenation Using Frustrated Lewis Pairs: Evidence for a Chain-Growth Coupling Mechanism

Zhenbo Mo, Arnab Rit, Jesús Campos, Eugene L. Kolychev, and Simon Aldridge*

Inorganic Chemistry Laboratory, Department of Chemistry, University of Oxford, South Parks Road, Oxford OX1 3QR, United Kingdom

Supporting Information

ABSTRACT: The catalytic dehydrogenation of ammonia- and amine-boranes by a dimethylxanthene-derived frustrated Lewis pair is described. Turnover is facilitated on a thermodynamic basis by the ready release of H₂ from the weakly basic PPh₂-containing system. *In situ* NMR studies and the isolation of intermediates from stoichiometric reactions support a mechanism initiated by B–H activation, followed by end-growth BN coupling involving the terminal NH bond of the bound BN fragment and a BH bond of the incoming borane monomer.


Dehydrocoupling represents a powerful synthetic protocol for the construction of both homo- and heteronuclear E–E bonds in molecular and materials science.¹ Among such processes, the formation of B–N bonds, e.g., from ammonia-borane (AB, H₃B·NH₃) or related amine-boranes (H₃B·NH₂R), has been the subject of significant research effort, reflecting potential applications in hydrogen storage technologies and the synthesis of novel polymeric materials.² While B–N coupling can be effected thermally in some cases, reflecting the complementary polarities of B–H and N–H bonds,³ catalytic approaches have also been investigated with a view to enhancing the rate of H₂ release and/or controlling the nature of the dehydrocoupled product(s).⁴ Metal complexes (from all parts of the Periodic Table) feature prominently among the most active catalyst systems reported to date, but other approaches, including the use of ionic liquids and Brønsted and Lewis acids/bases, have also been developed.^{5–9}

Frustrated Lewis pairs (FLPs), although described conceptually as far back as 1942,¹⁰ emerged as a new paradigm for small molecule capture and activation following a landmark report by Stephan et al. in 2006.^{11,12} The ability of such systems (archetypally based on sterically shrouded tertiary phosphine and fluoroarylborane units) to activate strong nonpolar bonds, such as that in H₂, has been exploited in “metal-free” catalysis.¹³ Prominent among such processes is FLP-catalyzed hydrogenation chemistry,¹³ which relies on the ability of the phosphonium hydroborate derived from heterolytic H₂ cleavage to transfer H⁺/H[−] to unsaturated substrates such as imines. By contrast, FLP systems capable of effecting turnover in dehydrogenation chemistry are very rare indeed.^{9,14} In part this is due to the fact that in the absence of a suitable coreagent to act as a hydrogen acceptor (or the application of elevated temperature/vacuum), the spontaneous loss of H₂ in the reverse sense is often thermodynamically unfavorable.^{15,16} As a consequence, the

evolution of H₂ leads to rapid, irreversible sequestering of the FLP and prevents turnover.

In recent work we have developed single component FLPs based on a dimethylxanthene backbone that possess a P···B separation (4.2–4.5 Å) preorganized for the facile uptake of H₂. Moreover, by varying the cumulative Lewis acid/base strength of phosphine and borane components, a system can be designed (1; Scheme 1) which cleaves H₂ to give an equilibrium mixture of free

Scheme 1. Synthesis and H₂ Activation by Dimethylxanthene-Based FLPs 1–3



FLP and zwitterionic 1-(H)₂ in solution at room temperature.¹⁶ With this reversibility in mind, we hypothesized that 1 and related systems might be competent for the catalytic dehydrocoupling of AB and related substrates. As such, a new area of catalytic endeavor might be opened up to mediation by a metal-free FLP approach.⁹

The phosphonium borohydride 1-(H)₂ derived from PMes₂/B(C₆F₅)₂ functionalized 1 has been shown to exist in equilibrium with the “free” FLP and dihydrogen at 295 K (~20:1 1-(H)₂:1 mixture in bromobenzene; Scheme 1).¹⁶ Targeting applications in dehydrogenation chemistry, the related PPh₂ variants 2 and 3 were targeted, reasoning that the incorporation of weaker Lewis acid/base components should lead to more favorable thermodynamics for H₂ loss.¹⁷ These systems can be synthesized from 4,5-dibromo-9,9-dimethylxanthene using a similar approach to that employed for 1,¹⁶ and both 2 and 3 can be shown crystallographically to constitute a FLP (see SI). As expected, the uptake of H₂ in each case is less thermodynamically favorable than that by 1: in the case of 2, only 5% conversion to 2-(H)₂ is observed spectroscopically upon exposure to H₂ at 4 atm pressure at 295 K. We were unable to obtain any evidence for the formation of 3-(H)₂, presumably due to the much weaker Lewis acidity of the –BMe₂ function. For 2, the values of ΔH° and ΔS° obtained by


Received: February 1, 2016

Published: February 26, 2016

monitoring the response of the equilibrium constant to temperature (34 kJ mol^{-1} and $138 \text{ J mol}^{-1} \text{ K}^{-1}$) can be compared to values of 38 kJ mol^{-1} and $102 \text{ J mol}^{-1} \text{ K}^{-1}$ obtained for the corresponding release of H_2 by 1-(H)_2 .

Given the relative ease of H_2 loss from 2-(H)_2 , we set out to investigate the use of **2** as a catalyst in dehydrogenation chemistry. Accordingly, **2** is found to catalyze the dehydrocoupling of methylamine- and ammonia-borane to the corresponding Me- or H-substituted borazine and of dimethylamine-borane to $\text{H}_2\text{B}(\mu\text{-NMe}_2)_2\text{BH}_2$ at 328 K and 1 mol % catalyst loading (Scheme 2).


Scheme 2. Catalytic Dehydrogenation of Amine and Ammonia-Boranes by **2**



While the turnover frequencies for these catalytic processes ($\sim 4 \text{ h}^{-1}$ for $\text{H}_3\text{B}\cdot\text{NH}_3$) are modest in comparison to the most active transition metal catalysts, they do, to our knowledge, represent the first example of methylamine- or ammonia-borane dehydrogenation catalyzed by a main group FLP.^{9,14}

In order to probe potential dehydrocoupling mechanisms, attempts were made to monitor reactivity *in situ* by multinuclear NMR measurements and to examine the products of the stoichiometric reactions of **2** with ammonia- and amine-boranes at 295 K. Revealingly, the 1:1 reactions of **2** with adducts of the general composition $\text{H}_3\text{B}\cdot\text{NMe}_n\text{H}_{3-n}$ ($n = 0\text{--}3$) all proceed via B–H activation, generating a boron-bound hydride and a $\text{P}\text{--}\text{BH}_2\text{NMe}_n\text{H}_{3-n}$ unit (Scheme 3). All four structures were

Scheme 3. Stoichiometric Reactions of **2** with $\text{H}_3\text{B}\cdot\text{NMe}_n\text{H}_{3-n}$ ($n = 0\text{--}3$) Proceeding via B–H Bond Cleavage



confirmed by X-ray crystallography (Figure 1 and SI), and in the case of $2\text{-(H)}(\text{BH}_2\text{NH}_3)$, $2\text{-(H)}(\text{BH}_2\text{NH}_2\text{Me})$, and $2\text{-(H)}(\text{BH}_2\text{NHMe}_2)$, the existence of short $\text{NH}\cdots\text{HB}$ contacts ($\sim 1.9 \text{ \AA}$) involving the hydride of the $[\text{ArB}(\text{C}_6\text{F}_5)_2]^-$ unit and a proton of the coordinated amine could be inferred from the difference Fourier maps.¹⁸

Spectroscopically, each of the B–H activation products derived from **2** is characterized by a ^{31}P resonance in the range $1.7\text{--}6.3 \text{ ppm}$ and by ^{11}B NMR signals at $\delta_{\text{B}} -9.4$ to -20.7 ppm (PBH_2N) and $\sim -21.5 \text{ ppm}$ (for the borohydride unit). With respect to chemistry occurring under catalytic conditions, the ^1H , ^{11}B , and ^{31}P resonances for $2\text{-(H)}(\text{BH}_2\text{NH}_3)$, $2\text{-(H)}(\text{BH}_2\text{NH}_2\text{Me})$, and $2\text{-(H)}(\text{BH}_2\text{NHMe}_2)$ are found to correspond to the major species present in solution at short reaction times. In addition, isolated samples of these adducts are competent in the catalytic





Figure 1. Structures of $2\text{-(H)}(\text{BH}_2\text{NH}_3)\cdot\text{CH}_2\text{Cl}_2$ (left) and $2\text{-(H)}(\text{BH}_2\text{NH}_2\text{Me})$ (right) as determined by X-ray crystallography. Here and elsewhere: Most H atoms and solvate molecule omitted, and selected aryl groups shown in wireframe format for clarity; thermal ellipsoids set at the 40% probability level. Key metrical parameters: (for $2\text{-(H)}(\text{BH}_2\text{NH}_3)\cdot\text{CH}_2\text{Cl}_2$) $\text{P}(1)\cdots\text{B}(1)$ $4.692(5)$ \AA , $\text{P}(1)\text{--}\text{B}(2)$ $1.952(4)$, $\text{N}(1)\text{--}\text{B}(2)$ $1.578(7)$ \AA ; (for $2\text{-(H)}(\text{BH}_2\text{NH}_2\text{Me})$) $\text{P}(1)\cdots\text{B}(1)$ $4.600(3)$, $\text{P}(1)\text{--}\text{B}(2)$ $1.958(3)$, $\text{B}(2)\text{--}\text{N}(1)$ $1.578(4)$ \AA .

dehydrogenation of the corresponding ammonia- or amine-borane, providing comparable performance to **2** itself under matching conditions. As such, it seems likely that these systems are viable intermediates in the respective catalytic dehydrocoupling processes.

Uhl and Sloutweg have previously reported the synthesis of complexes featuring the H_2BNH_2 and H_2BNMe_2 fragments trapped by a phosphorus/aluminum FLP,⁹ and in the case of the dimethylaminoborane adduct showed that such a species is a potential intermediate in the dehydrocoupling of $\text{H}_3\text{B}\cdot\text{NHMe}_2$ to $\text{H}_2\text{B}(\mu\text{-NMe}_2)_2\text{BH}_2$ in a melt. With this in mind, together with the close $\text{NH}\cdots\text{HB}$ contacts measured for $2\text{-(H)}(\text{BH}_2\text{NH}_3)$, $2\text{-(H)}(\text{BH}_2\text{NH}_2\text{Me})$, and $2\text{-(H)}(\text{BH}_2\text{NHMe}_2)$ in the solid state, we hypothesized that the loss of H_2 from these systems might be relevant in a catalytic context. To our surprise, however, solutions of these adducts are inert to thermal loss of H_2 in the temperature regime associated with the catalytic processes. Thus, e.g., $2\text{-(H)}(\text{BH}_2\text{NH}_3)$ remains unchanged in dichloromethane- d_2 after 24 h at 328 K, as judged by *in situ* multinuclear NMR measurements. Dehydrogenation can be effected through the use of a hydrogen acceptor such as $\text{Pr}_2\text{N=BH}_2$ (in a fashion similar to that demonstrated by Manners et al.),¹⁹ and the aminoborane product $2\text{-(BH}_2\text{NH}_2)$ characterized both spectroscopically and crystallographically (Scheme 4). However, it appears that this species is not relevant to the productive catalytic cycle: $2\text{-(BH}_2\text{NH}_2)$ is inert to further reaction with AB under

Scheme 4. Formation of the off-cycle Aminoborane Adduct $2\text{-(BH}_2\text{NH}_2)$ ^a



^aInset: structure of $2\text{-(BH}_2\text{NH}_2)\cdot\text{CH}_2\text{Cl}_2$ as determined by X-ray crystallography. Key metrical parameters: $\text{P}(1)\cdots\text{B}(1)$ $4.268(2)$, $\text{P}(1)\text{--}\text{B}(2)$ $1.968(2)$, $\text{B}(2)\text{--}\text{N}(1)$ $1.612(3)$, $\text{N}(1)\text{--}\text{B}(1)$ $1.582(3)$ \AA .

conditions identical to those used in catalytic runs, remaining unchanged over 24 h at 328 K in dichloromethane- d_2 solution.

The apparent inertness of isolated samples of 2-(H)(BH₂NH₂) and 2-(H)(BH₂NHMe) toward thermal loss of H₂ suggested that their onward reactivity might be dependent on the presence of additional ammonia- or amine-borane. In the case of the 2/H₃B·NH₃ catalytic runs, this hypothesis is also consistent with the observation of a second intermediate species, which can be shown (by comparison of multinuclear NMR data with the independently synthesized complex) to be the oligomeric borane adduct, 2-(H)(BH₂NH₂BH₂NH₃) (SI and Figure 2, left). Although labile in





Figure 2. Structures of 2-(H)(BH₂NH₂BH₂NH₃) (left) and 2-(H)(BH₂NHMeBH₂NH₂Me) (right) as determined by X-ray crystallography. Key metrical parameters: (for 2-(H)(BH₂NH₂BH₂NH₃)) P(1)⋯B(1) 4.690(3), P(1)–B(2) 1.964(3), B(2)–N(1) 1.558(3), N(1)–B(3) 1.575(3), B(3)–N(2) 1.587(3) Å; (for 2-(H)(BH₂NHMeBH₂NH₂Me)) P(1)–B(1) 4.668(4), P(1)–B(2) 1.969(5), B(2)–N(1) 1.562(5), N(1)–B(3) 1.567(5), B(3)–N(2) 1.604(5) Å.

dichloromethane solution at 295 K, this compound can be obtained as a compositionally pure material via the alternative reaction of 2 with H₃B·NH₂BH₂·NH₃ and crystallized from CH₂Cl₂/hexane at 273 K. Structurally, 2-(H)(BH₂NH₂BH₂NH₃) (in common with 2-(H)(BH₂NH₃)) features an P-bound aminoborane chain which is additionally “anchored” at the β-N position through an NH⋯HB interaction with the borohydride group (~1.91 Å).

While the lability of 2-(H)(BH₂NH₂BH₂NH₃) prevents meaningful attempts to study its onward reactivity, the corresponding product derived from H₃B·NHMeBH₂·NH₂Me by terminal B–H activation is stable in dichloromethane solution at 295 K over a period of 24 h and can therefore be used to probe potential mechanistic pathways (Scheme 5 and Figure 2). Moreover, while 2-(H)(BH₂NHMeBH₂NH₂Me) is most conveniently accessed for synthetic purposes from the preformed BNBN oligomer, it is also the product of the reaction of 2-(H)(BH₂NH₂Me) with a second equivalent of H₃B·NHMe₂ (50% conversion over a period of 2 h at 328 K in dichloromethane), thereby providing evidence for a dehydrogenative chain-growth process.²⁰ In addition, the reaction of 2-(H)(BH₂NHMeBH₂NH₂Me) itself with a further equivalent of H₃B·NH₂Me generates the (known) cyclic trimer (H₂BNHMe)₃ as the major product, along with small amount of trimethylborazine. This cyclic trinuclear borazane can also be identified in 2/H₃B·NH₂Me catalytic runs by *in situ* multinuclear NMR spectroscopy (Figure S3). These stoichiometric reactivity studies suggest that a viable reaction pathway under catalytic conditions involves the stepwise assimilation of two BN-containing units to give 2-(H)(BH₂NHMeBH₂NH₂Me), followed by uptake of a third molecule of the amine-borane and ejection of cyclic (H₂BNHMe)₃. The loss of the cyclic borazane via “backbiting” is presumably very rapid, as no spectroscopic evidence is obtained for the preceding P-BNBNBN linear chain. However, consistent with its role as an intermediate in the overall dehydrogenation of

Scheme 5. Stepwise Growth of the Oligomeric Aminoborane Chain by Sequential Uptake of H₃B·NH₂Me Monomer



H₃B·NH₂Me to trimethylborazine, *cyclo*-(H₂BNHMe)₃ can be synthesized independently²¹ and is shown to undergo further dehydrogenation under catalytic conditions (CH₂Cl₂ solution, 328 K, using 1 mol % 2) to generate the final trimethylborazine product over a period of time (24 h) consistent with its competence as a catalytic intermediate.

While the evolution of dihydrogen in both the 2/H₃B·NH₃ and 2/H₃B·NH₂Me catalytic systems can be shown explicitly by ¹H NMR, the unfavorable equilibrium for the capture of H₂ by 2 means that 2-(H)₂ does not build up to sufficient concentrations and that it can be identified *in situ*, even at the end of catalytic runs. Instead, the major FLP-containing species present when all of the substrate has been consumed can be identified as the respective B-bound amine adducts 2-NH₂R (R = H or Me), by comparison with the products obtained from the corresponding Lewis acid/base reactions of 2 with NH₃ or MeNH₂.

With a view to probing in more depth the mechanism by which growth of the BN oligomeric chain occurs, the reaction between 2-(H)(BH₂NHMe₂) and H₃B·NH₂Me was investigated (SI, Scheme S7). Although this reaction is kinetically slow, *in situ* NMR measurements indicate that the incoming methylamine-borane unit is assimilated in a manner consistent with an end-growth dehydrogenative coupling mechanism, rather than via insertion into the P–B bond of 2-(H)(BH₂NHMe₂). The formation of the PBNBN backbone in the product is consistent with the appearance of ¹¹B signals at δ_B = –12.6 and –6.4 (cf. δ_B = –13.8 and –4.6 for the corresponding PBN and NBN units in 2-(H)(BH₂NHMeBH₂NH₂Me)). Most informatively, the ¹H resonance associated with the terminal N-bound methyl group in 2-(H)(BH₂NMe₂BH₂NH₂Me) is a 1:2:1 triplet of intensity three at δ_H = 2.24 (³J_{HH} = 5.6 Hz; cf. δ_H = 2.22 for the NH₂Me unit in 2-(H)(BH₂NHMeBH₂NH₂Me)), consistent with the presence of a capping NH₂Me group (rather than NMe₂H).

In conclusion, we have demonstrated the catalytic dehydrogenation of ammonia- and amine-boranes by a FLP based on a preorganized dimethylxanthene backbone. This chemistry is

initiated by B–H activation, followed by end-growth BN coupling involving the terminal NH bond of the bound BN fragment and a BH bond of the incoming borane monomer. The propensity for competing intramolecular H₂ loss at this stage appears to be minimal: independent synthesis of the P–BH₂NH₂–B amino-borane complex allows us to demonstrate that it is not a viable catalytic intermediate and that it is not formed in detectable amounts as a side-product. The reluctance of the system to lose H₂ intramolecularly potentially reflects the very strong B–H bond to the electron-deficient BAr(C₆F₅)₂ unit. Alternative intermolecular loss of H₂ at this stage to generate a P-bound BBN linear oligomer can be demonstrated explicitly for methylamine-borane. Nonetheless, this is the only step in the stoichiometric formation of the cyclic triborazane (H₂BNHMe)₃ from H₃B–NH₂Me, e.g., which does not occur spontaneously at room temperature (requiring heating to 328 K). As such, it seems plausible that under catalytic conditions this might be the rate-limiting step.²² That the assimilation of the second BN unit appears to be slower than the subsequent uptake of the third equivalent of amine-borane potentially reflects greater steric shrouding of the terminal N–H bond in species such as 2-(H)(BH₂NH₂Me) than in its longer chain analogue 2-(H)(BH₂NHMeBH₂NH₂Me).

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.6b01170.

Experimental details and data (PDF)

Crystallographic data (CIF)

■ AUTHOR INFORMATION

Corresponding Author

*simon.aldridge@chem.ox.ac.uk

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

EPSRC (EP/K503769/1, EP/K014714/1); EU Marie Curie (E.L.K., A.R.: grants PIEF-GA-2013-626441 and PIEF-GA-2013-622806).

■ REFERENCES

- (1) (a) Hansmann, M. M.; Melen, R. L.; Wright, D. W. *Chem. Sci.* **2011**, 2, 1554. (b) Leitao, E. M.; Jurca, T.; Manners, I. *Nat. Chem.* **2013**, 5, 817. (c) Waterman, R. *Chem. Soc. Rev.* **2013**, 42, 5629.
- (2) (a) Staubitz, A.; Robertson, A. P. M.; Sloan, M. E.; Manners, I. *Chem. Rev.* **2010**, 110, 4023. (b) Staubitz, A.; Robertson, A. P. M.; Manners, I. *Chem. Rev.* **2010**, 110, 4079.
- (3) Stephens, F. H.; Pons, V.; Baker, R. T. *Dalton Trans.* **2007**, 2613.
- (4) For an early report see: Jaska, C. A.; Temple, K.; Lough, A. J.; Manners, I. *J. Am. Chem. Soc.* **2003**, 125, 9424.
- (5) (a) Marder, T. B. *Angew. Chem., Int. Ed.* **2007**, 46, 8116. (b) Pons, V.; Baker, R. T. *Angew. Chem., Int. Ed.* **2008**, 47, 9600.
- (6) For an early report of ionic liquids to accelerate AB dehydrogenation see: Bluhm, M. E.; Bradley, M. G.; Butterick, R., III; Kusari, U.; Sneddon, L. G. *J. Am. Chem. Soc.* **2006**, 128, 7748.
- (7) For an early report of acid catalysis of AB dehydrogenation see: (a) Stephens, F. H.; Baker, R. T.; Matus, M. H.; Grant, D. J.; Dixon, D. A. *Angew. Chem., Int. Ed.* **2007**, 46, 746. For a recent example of a similar process effected by a bifunctional borane see: Lu, Z.; Schweighauser, L.; Hausmann, H.; Wegner, A. *Angew. Chem., Int. Ed.* **2015**, 54, 15556.
- (8) Lewis base catalyzed AB dehydrogenation: Himmelberger, D. W.; WonYoon, C.; Bluhm, M. E.; Carroll, P. J.; Sneddon, L. G. *J. Am. Chem. Soc.* **2009**, 131, 14101.
- (9) Uhl, S.; Slootweg, J. C.; Lammertsma, K.; Uhl, W. *Angew. Chem., Int. Ed.* **2013**, 52, 4256. See also: (b) Alexander, J. M.; Bercaw, J. E. *Chem. Commun.* **2010**, 46, 1709. (c) Whittell, G. R.; Balmond, E. I.; Robertson, A. P. M.; Patra, S. K.; Haddow, N.; Manners, I. *Eur. J. Inorg. Chem.* **2010**, 2010, 3967.
- (10) Brown, H. C.; Schlesinger, H. I.; Cardon, S. Z. *J. Am. Chem. Soc.* **1942**, 64, 325.
- (11) Welch, G. C.; Juan, R. R. S.; Masuda, J. D.; Stephan, D. W. *Science* **2006**, 314, 1124.
- (12) For recent reviews see: (a) Stephan, D. W.; Erker, G. *Angew. Chem., Int. Ed.* **2010**, 49, 46. (b) Stephan, D. W.; Greenberg, S.; Graham, T. W.; Chase, P.; Hastie, J. J.; Geier, S. J.; Farrell, J. M.; Brown, C. C.; Heiden, Z. M.; Welch, G. C.; Ullrich, M. *Inorg. Chem.* **2011**, 50, 12338. (c) Erker, G. *Dalton Trans.* **2011**, 40, 7475. (d) Stephan, D. W. *Org. Biomol. Chem.* **2012**, 10, 5740. (e) Stephan, D. W.; Erker, G. *Angew. Chem., Int. Ed.* **2015**, 54, 6400. (f) Stephan, D. W. *Acc. Chem. Res.* **2015**, 48, 306. (g) Stephan, D. W. *J. Am. Chem. Soc.* **2015**, 137, 10018.
- (13) For an early report of FLP-catalysed hydrogenation see: (a) Chase, P. A.; Welch, G. C.; Jurca, T.; Stephan, D. W. *Angew. Chem., Int. Ed.* **2007**, 46, 8050. (b) Spies, P.; Schwendemann, S.; Lange, S.; Kehr, G.; Fröhlich, R.; Erker, G. *Angew. Chem., Int. Ed.* **2008**, 47, 7543. For a review of related hydrosilylation chemistry see: (c) Piers, W. E.; Marwitz, A. J. V.; Mercier, L. G. *Inorg. Chem.* **2011**, 50, 12252.
- (14) FLPs incorporating transition metals also mediate BN dehydrocoupling: Chapman, A. M.; Haddow, M. F.; Wass, D. F. *J. Am. Chem. Soc.* **2011**, 133, 18463.
- (15) For selected examples of FLPs capable of the activation of H₂ and release under vacuum see ref 11 and (a) Wang, H.; Fröhlich, R.; Kehr, G.; Erker, G. *Chem. Commun.* **2008**, 5966. (b) Sumerin, V.; Schulz, F.; Atsumi, M.; Wang, C.; Nieger, M.; Leskelä, M.; Repo, T.; Pyykkö, P.; Rieger, B. *J. Am. Chem. Soc.* **2008**, 130, 14117. (c) Sumerin, V.; Schulz, F.; Nieger, M.; Leskelä, M.; Repo, T.; Rieger, B. *Angew. Chem., Int. Ed.* **2008**, 47, 6001. (d) Ullrich, M.; Lough, A. J.; Stephan, D. W. *J. Am. Chem. Soc.* **2009**, 131, 52. (e) Schwendemann, S.; Fröhlich, R.; Kehr, G.; Erker, G. *Chem. Sci.* **2011**, 2, 1842. (f) Sumerin, V.; Chernichenko, K.; Nieger, M.; Leskelä, M.; Rieger, B.; Repo, T. *Adv. Synth. Catal.* **2011**, 353, 2093. (g) Jiang, C.; Blacque, O.; Fox, T.; Berke, H. *Dalton Trans.* **2011**, 40, 1091. (h) Chernichenko, K.; Nieger, M.; Leskelä, M.; Repo, T. *Dalton Trans.* **2012**, 41, 9029. (i) Binding, S. C.; Zaher, H.; Chadwick, F. M.; O'Hare, D. *Dalton Trans.* **2012**, 41, 9061. Weaker acid/base combinations have also been reported which cleave dihydrogen in solution at low temperatures and/or elevated pressures, which quantitatively evolve H₂ on warming to room temperature: (j) Greb, L.; Oña-Burgos, P.; Shirmer, B.; Grimme, S.; Stephan, D. W.; Paradies, J. *Angew. Chem., Int. Ed.* **2012**, 51, 10164. (k) Greb, L.; Tussing, S.; Schirmer, B.; Oña-Burgos, P.; Kaupmees, K.; Lökov, M.; Leito, I.; Grimme, S.; Paradies, J. *Chem. Sci.* **2013**, 4, 2788. (l) Chernichenko, K.; Kótai, B.; Pápai, I.; Zhivonitko, V.; Nieger, M.; Leskelä, M.; Repo, T. *Angew. Chem., Int. Ed.* **2015**, 54, 1749.
- (16) Mo, Z.; Kolychev, E. L.; Rit, A.; Campos, J.; Niu, H.; Aldridge, S. J. *Am. Chem. Soc.* **2015**, 137, 12227.
- (17) (a) Rokob, T. A.; Hamza, A.; Pápai, I. *J. Am. Chem. Soc.* **2009**, 131, 10701. (b) Neu, R.; Ouyang, E. Y.; Geier, S. J.; Zhao, X.; Ramos, A.; Stephan, D. W. *Dalton Trans.* **2010**, 39, 4285.
- (18) Schulz, F.; Sumerin, V.; Heikkinen, S.; Pedersen, B.; Wang, C.; Atsumi, M.; Leskelä, M.; Repo, T.; Pyykkö, P.; Petry, W.; Rieger, B. *J. Am. Chem. Soc.* **2011**, 133, 20245.
- (19) Leitao, E. M.; Stubbs, N. E.; Robertson, A. P. M.; Helten, H.; Cox, R. J.; Lloyd-Jones, G. C.; Manners, I. *J. Am. Chem. Soc.* **2012**, 134, 16805.
- (20) Johnson, H. C.; Robertson, A. P. M.; Chaplin, A. B.; Sewell, L. J.; Thompson, A. L.; Haddow, M. F.; Manners, I.; Weller, A. S. *J. Am. Chem. Soc.* **2011**, 133, 11076.
- (21) Framery, E.; Vaultier, M. *Heteroat. Chem.* **2000**, 11, 218.
- (22) The reduced reaction rates in catalytic runs using D₃B–NH₂Me ($k_{\text{H}}/k_{\text{D}} = 3.9$) and H₃B–ND₂Me ($k_{\text{H}}/k_{\text{D}} = 2.9$) are consistent with a rate-limiting step involving the breakage of both B–H and N–H bonds (SI).