

Functional single nucleotide polymorphisms within the cyclin-dependent kinase inhibitor 2A/2B region affect pancreatic cancer risk

SUPPLEMENTARY TABLE

Supplementary Table S1: List of SNPs and Hardy Weinberg equilibrium values in the populations in the study

SNP	Alleles (M/m)	Call rate	HWE		Minor allele frequency	
			Caucasians	Japanese	Caucasians	Japanese ^a
rs3731257	C/T	96.62	0.153	0.638	0.28	0.56
rs11515	C/G	98.06	1.000	1.000	0.15	0.01
rs2518719	G/A	98.69	0.136	-	0.13	0.00
rs3731249	C/T	98.28	0.432	-	0.04	0.00
rs3731246	C/G	79.95	0.756	1.000	0.12	0.01
rs2811708	T/G	92.75	0.886	0.796	0.26	0.16
rs3731239	C/T	97.99	0.222	0.771	0.35	0.14
rs3731211	A/T	95.00	0.037	0.215	0.29	0.19
rs2811710	T/C	88.48	0.044	0.143	0.37	0.25
rs3218009	C/G	99.19	0.031	-	0.10	0.00
rs3217992	G/A	98.52	0.607	0.075	0.41	0.57
rs1063192	G/A	96.85	0.574	0.142	0.39	0.20
rs3217986	C/A	98.52	0.198	0.143	0.08	0.05

^a Allele frequency of the Caucasian minor allele in the Japanese population.