

Splitting fields of real irreducible representations of finite groups

Dmitrii V. Pasechnik¹

¹Department of Computer Science, University of Oxford, UK*

2021-07-15

Abstract

We show that any irreducible representation ρ of a finite group G of exponent n , realisable over \mathbb{R} , is realisable over the field $E := \mathbb{Q}(\zeta_n) \cap \mathbb{R}$ of real cyclotomic numbers of order n , and describe an algorithmic procedure transforming a realisation of ρ over $\mathbb{Q}(\zeta_n)$ to one over E .

Introduction

Let G be a finite group of exponent n . A celebrated result by R. Brauer states that any complex irreducible character $\chi \in \text{Irr}(G)$ of G is afforded by an F -representation $\rho_\chi : G \rightarrow \text{GL}_d(F)$, where $F = \mathbb{Q}(\zeta_n)$, the field of cyclotomic numbers of order n (here $\zeta_n := e^{\frac{2\pi i}{n}}$), see [Isa94, (10.3)]. Let $E := \mathbb{Q}(\zeta_n) \cap \mathbb{R} \subset F$ be the maximal real subfield of F . The first result of this note is as follows.

Theorem 1. *Let χ be an irreducible real-valued character of G of degree $d := \chi(1)$ with Frobenius-Schur indicator $\nu_2(\chi) = 1$. Then E is a splitting field of χ , i.e. χ is afforded by an E -representation ρ , and the Schur index $m_E(\chi)$ equals 1.*

Our proof of Theorem 1 invokes Serre's induction theorem for real characters [Ser71], [CR87, Theorem 73.18], and then follows the line of proof of Brauer's theorem [Isa94, (10.3)]. It is surprising that it has not appeared anywhere, at least as far as we know.

Remark. Independently and simultaneously, Robert Guralnick and Gabriel Navarro proved Theorem 1 by a similar method, although not using [Ser71].

Recall that the *Frobenius-Schur indicator* $\nu_2(\chi) := \frac{1}{|G|} \sum_{g \in G} \chi(g^2)$ is an invariant classifying complex representations of G into three different types, see [Isa94, (4.5)]. Namely, $\nu_2(\chi) = 0$ if χ is not real-valued, and $\nu_2(\chi) = -1$ if χ

*email: dima@pasechnik.info

is real-valued, but is not afforded by a real-valued representation; $\nu_2(\chi) = 1$ if and only if χ is afforded by a real-valued representation.

For a number field $K \supseteq \mathbb{Q}$, the *Schur index* $m_K(\chi)$ is an invariant of χ controlling the possibility to realise ρ_χ over K , see e.g. [CR06, Sect. 41] and [Isa94, Chapter 10]. Namely, let $S \supseteq K$ be a splitting field of χ . Then $m_K(\chi) := \min_{\substack{K \subseteq M \subseteq S \\ \rho_\chi \text{ realisable over } M}} [M : K(\chi)]$, where we denoted by $[M : K(\chi)]$

the degree of M as a field extension over $K(\chi)$, the field extension of K generated by the values of χ . In particular, the claim of Theorem 1 amounts to stating that $m_E(\chi) = 1$.

Apart from theoretical significance, the question of finding a splitting field is relevant in group theory algorithms. Standard algorithms such as J. Dixon's algorithm [Dix93] for constructing complex, and real, irreducible representations (one implementation in the computer algebra system GAP [GAP21] of it is described in [DD10]) do induction from 1-dimensional representations of subgroups of G , which are defined over F . One advantage of working over E instead is that the degree of E is half of the degree of F .

In particular for applications, e.g. in extremal combinatorics, in physics, etc. it is often necessary to reduce a representation to a direct sum of real irreducibles, and exact methods for this process benefit from explicit knowledge of the irreducibles, using well known formulas from [Ser77, Sect.2.7], as implemented in our GAP package RepnDecomp [HP20].

Our second result amounts to the algorithmic counterpart of Theorem 1, that is, to a procedure to compute, for a representation $\rho : G \rightarrow \mathrm{GL}_d(F)$ realisable over reals, an explicit matrix $Q \in \mathrm{GL}_d(F)$ such that $Q^{-1}\rho(G)Q \subset \mathrm{GL}_d(E)$, i.e. Q transforms ρ to an E -representation.

Theorem 2. *Let $\rho : G \rightarrow \mathrm{GL}_d(F)$ be a representation of G realisable over \mathbb{R} . Then $P \in \mathrm{SL}_d(F)$ such that $P\rho(g) = \overline{\rho(g)}P$ for any $g \in G$, and $P\overline{P} = I$, can be explicitly computed from the $\rho(G)$ -invariant forms. Let $\xi \in F^*$ s.t. $-\frac{\xi}{\overline{\xi}}$ is not an eigenvalue of P , and $Q := \overline{\xi}P + \xi I$. Then $Q \in \mathrm{GL}_d(F)$ and $Q^{-1}\rho(G)Q \subset \mathrm{GL}_d(E)$.*

The only part of Theorem 2 which uses Theorem 1 is the claim that P can be chosen so that $P\overline{P} = I$. Algorithmically, one computes P s.t. $P\overline{P} = \mu I$ for $0 < \mu \in E$, and then has to solve the *norm equation*

$$x\overline{x} = \mu, \quad \text{for } x \in F. \quad (1)$$

Theorem 1 implies that (1) is always solvable. Several parts of the proof of Theorem 2 are contained in [GH97] and [Fie09], although our approach is more explicit, and for odd d we provide an explicit solution (Lemma 5), not involving solving (1), which is a nontrivial number-theoretic problem.

Proof of Theorem 1

Our main tool is Serre's induction theorem [CR87, (73.18)].

Theorem 3. (Serre) *The character χ of a real representation of G is a \mathbb{Z} -linear combination*

$$\chi = \sum_{\phi} a_{\phi} \text{Ind}_H^G(\phi) \quad (2)$$

of real-valued induced characters $\text{Ind}_H^G(\phi)$, with $H \leq G$, and ϕ a character of H . Further, ϕ is either linear and takes values ± 1 , or $\phi = \lambda + \bar{\lambda}$ for a linear character λ of H , or ϕ is dihedral. \square

A dihedral character ϕ of a group H is a degree 2 irreducible character of H s.t. $H/\ker \phi \cong D_{2m}$, dihedral group of order $2m$.

Note that by [Isa94, (10.2.f)] $m_E(\chi)$ divides $m_{\mathbb{Q}}(\chi) \leq 2$, where the latter inequality holds by the Brauer-Speiser Theorem [Isa94, p.171]. Therefore it suffices to show that $m_E(\chi) = 2$ is not possible in our situation.

Let θ be a character of an E -representation of G . Then by [Isa94, (10.2.c)] $m_E(\chi) \mid [\theta, \chi]$. Here $[\cdot, \cdot]$ is the usual scalar product of characters $[\theta, \chi] = \frac{1}{|G|} \sum_{g \in G} \theta(g) \chi(g)$, cf. [Isa94, (2.16)]. As χ is irreducible, $[\chi, \chi] = 1$, thus (2) implies

$$1 = [\chi, \chi] = \sum_{\phi} a_{\phi} [\text{Ind}_H^G(\phi), \chi]. \quad (3)$$

If every $\text{Ind}_H^G(\phi)$ is an E -representation, then $m_E(\chi) = 2$ is not possible, as otherwise an even integer on the right hand side of (3) equals 1.

It remains to see that every $\text{Ind}_H^G(\phi)$ is an E -representation.

This is trivially the case for linear ϕ , and so we are left with the dihedral case and the case $\phi = \lambda + \bar{\lambda}$. To simplify the rest of the proof, we use [Isa94, (10.9)] which says that if a prime p divides $m_E(\chi)$ then the Sylow p -subgroups of G are not elementary abelian. For $p = 2$ this means that $4 \mid n$, i.e. $i := \sqrt{-1} \in F$.

Lemma 1. *Let $H \leq G$, with G of exponent n , $4 \mid n$, and ϕ a character of H , either $\phi = \lambda + \bar{\lambda}$ with λ linear, or ϕ dihedral. Then ϕ is afforded by an E -representation.*

Proof. Note that $E = \mathbb{Q}(\zeta_n + \zeta_n^{-1})$ and $2 \cos \frac{2\pi}{n} = \zeta_n + \zeta_n^{-1}$. As $4 \mid n$, it can be shown that $\sin \frac{2\pi}{n} \in E$ in this case (in general this is not true).

In the case $\phi = \lambda + \bar{\lambda}$ we have $H/\ker \phi$ a cyclic group C of order m dividing n , $C \cong \langle \zeta_m \rangle$. We have $Z_m := \begin{pmatrix} \cos \frac{2\pi}{m} & -\sin \frac{2\pi}{m} \\ \sin \frac{2\pi}{m} & \cos \frac{2\pi}{m} \end{pmatrix} \in \text{SL}_2(E)$, and

$$\begin{aligned} \rho_{\phi} : C &\rightarrow \text{SL}_2(E) \\ \zeta_m^k &\mapsto Z_m^k, \quad 0 \leq k < m \end{aligned}$$

is the desired E -representation of C with character ϕ .

For dihedral ϕ we have $H/\ker \phi$ a dihedral group $D = \langle a, b \mid 1 = a^m = b^2 = (ab)^2 \rangle$ of order $2m$ dividing n , with normal cyclic subgroup C of order m , so that the restriction $\phi_C = \lambda + \bar{\lambda}$ is as in the previous case, and $\phi_{D-C} = 0$. We have $Z_m \in \mathrm{SL}_2(E)$ as in the previous case, and $R_0 := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \in \mathrm{GL}_2(E)$ satisfying $R_0 Z_m R_0 = Z_m^{-1}$ and

$$\begin{aligned} \rho_\phi : D &\rightarrow \mathrm{GL}_2(E) \\ a^k b^\ell &\mapsto Z_m^k R_0^\ell, \quad 0 \leq k < m, \ 0 \leq \ell \leq 1, \end{aligned}$$

is the desired E -representation of D with character ϕ . □

This completes the proof of Theorem 1. The last step, i.e. the proof of Lemma 1, could also be accomplished in a less explicit way, by invoking the construction of Theorem 2; the matrix P mapping ρ_ϕ to its conjugate can be chosen to be equal to $P = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, satisfying the only condition, $P\bar{P} = I$. In particular this approach allows to prove a more general version of Lemma 1 which does not require $4 \mid n$.

Proof of Theorem 2

The case $n = 2$ is trivial, and we will assume $n \geq 3$ in what follows.

Recall that in general, χ has values in F , while a real-valued character has values in E . Whenever χ is E -valued, the image $\rho(G)$ of G under a representation $\rho := \rho_\chi$ affording χ leaves invariant a unique, up to scalar multiplication, non-zero G -invariant form M . It is a classical result due to Frobenius and Schur that if M is symmetric then χ is afforded by a real representation ρ , and $\nu_2(\chi) = 1$, cf. [Isa94, (4.19)].

Without loss of generality, $\chi(1) > 1$. Indeed, if $\chi(1) = 1$ then ρ is the same as χ , and we are done.

The proof of Frobenius-Schur in [Isa94, (4.19)] starts with the elementary fact that if Q is a transformation making ρ real then $Q^{-1}\rho Q = \overline{Q^{-1}\rho Q}$, thus $\overline{Q}Q^{-1}\rho = \rho\overline{Q}Q^{-1}$, and $P := \overline{Q}Q^{-1}$ transforms ρ to $\bar{\rho}$, i.e. $P^{-1}\bar{\rho}P = \rho$. Such a $P \in \mathrm{GL}_d(\mathbb{C})$ must exist irrespective of the existence of Q , as the characters of ρ and $\bar{\rho}$ are equal, although we can give an explicit construction $P = \Sigma^{-1}M$, with M as above, and Σ the matrix of a positive definite Hermitian $\rho(G)$ -invariant form.

Lemma 2. *Let χ be a real-valued character of G , and $\rho = \rho_\chi$ an F -representation affording χ . Then $P := \Sigma^{-1}M \in \mathrm{GL}_d(F)$ satisfies $P\rho(g) = \rho(g)P$ for all $g \in G$.*

Proof. As χ is real, ρ leaves invariant a nonzero G -invariant bilinear form M , i.e. $g^\top M g = M$ for all $g \in \rho$, cf. e.g. [Isa94, (4.14)]. As M can be found in the trivial sub-representation of the tensor square of ρ , $M \in M_d(F)$. As well, $\det M \neq 0$, as the kernel of M would give rise to a sub-representation of ρ , contradicting irreducibility of ρ .

Let $\Sigma := \sum_{h \in \rho(G)} h^\top \bar{h}$ - note that Σ is a Hermitian positive definite matrix, in particular $\det \Sigma > 0$, and $g^\top \Sigma \bar{g} = \Sigma$ for any $g \in \rho(G)$.

Choose $P := \Sigma^{-1} M$. Let's check that $P^{-1} \bar{P} P = \rho$ (we use $\det M \neq 0$ here). Let $g \in \rho(G)$. Then, as $(\bar{g} \Sigma^{-1} g^\top)^{-1} = (g^\top)^{-1} \Sigma \bar{g}^{-1} = \Sigma$,

$$\Sigma^{-1} M g = \bar{g} \Sigma^{-1} g^\top M g = \bar{g} \Sigma^{-1} M,$$

as required. \square

Now we have the equation

$$PQ = \bar{Q}, \quad \det Q \neq 0 \tag{4}$$

implying $\bar{P} P Q = \bar{P} \bar{Q} = Q$, i.e. $\bar{P} P = I$. The latter is an extra restriction, in the sense that our procedure does not guarantee that P computed as in Lemma 2 satisfies $\bar{P} P = I$. In general, one will need to solve (1) and multiply P by the inverse of a solution. However, (1) will always be solvable by Theorem 1.

Lemma 3. *Let $P \in \text{GL}_d(F)$ such that $Pg = \bar{g}P$ for any $g \in \rho(G)$. Then $P\bar{P} = \mu I$ for some $\mu \in E$.*

Proof. Note that $\bar{P}g = g\bar{P}$. Thus $P\bar{P}\bar{g} = Pg\bar{P} = \bar{g}P\bar{P}$. Thus $P\bar{P}$ lies in the centraliser of an irreducible representation $\bar{\rho}$. Hence, by Schur's Lemma, $P\bar{P} = \mu I$, for some $\mu \in F$.

It remains to show that $\mu \in E$. Using Lemma 2, and recalling that Σ and Σ^{-1} are Hermitian positive definite, i.e. $\Sigma^{-1} = U\bar{U}^\top$, and $M = M^\top$, we have $\mu I = P\bar{P} = \Sigma^{-1} M \bar{\Sigma}^{-1} \bar{M}$, i.e.

$$\mu \Sigma = M \bar{\Sigma}^{-1} \bar{M} = \overline{M U \bar{U}^\top M} = M \bar{U} U^\top \bar{M} = (M \bar{U})(\overline{M \bar{U}})^\top = \bar{\mu} \Sigma^\top = \bar{\mu} \Sigma,$$

implying $\mu = \bar{\mu}$. \square

It remains to solve (4) so that Q has entries in the splitting field of ρ . Note that the solution of (4) in [Isa94, Ch. 4] assumes that ρ is unitary; i.e. $\Sigma = I$; so in this case $P^\top = P$, and an explicit formula for Q is provided - which however does not work for us, as it involves square roots of eigenvalues of P . Fortunately, in [GH97, Prop. 1.3], there is an algorithmic proof of existence of the required solution of (4). In [loc.cit.] it is done for finite fields (and in bigger generality, for a field automorphism σ of finite order, referring to this result as a generalisation of Hilbert's Theorem 90), and in [Fie09] it was noted that it works for number

fields as well. One can also find there an easier observation, that for a randomly chosen $Y \in M_d(F)$ setting $Q = \overline{Y} + \overline{P}Y$ produces a solution to (4) with high probability. Here is an easy to prove variation of this claim.

Lemma 4. *Let $P, Y \in M_d(F)$ and $P\overline{P} = I$. Then $Q := \overline{Y} + \overline{P}Y$ satisfies $PQ = \overline{Q}$. Choosing $Y = \xi P$, with $\xi \neq 0$ and $-\xi/\overline{\xi}$ not being an eigenvalue of \overline{P} we have that $Q \in M_d(F)$ satisfies (4).*

Proof. Note that $PQ = P\overline{Y} + P\overline{P}Y = Y + P\overline{Y} = \overline{Q}$, as claimed. The claimed choice of ξ is possible as F is dense in \mathbb{C} . Further, with $Q = \overline{\xi P} + \xi \overline{P}P = \overline{\xi}(\overline{P} + \frac{\xi}{\overline{\xi}}I)$ we see that $Qv = 0$ holds for a non-zero vector v if and only if $\overline{P}v = -\frac{\xi}{\overline{\xi}}v$, which is not possible by the choice of ξ . \square

To complete the proof of Theorem 2 it suffices to observe that $Q^{-1}\rho(g)Q \in M_d(E)$ for any $g \in G$.

One can solve (1) in the case of odd d without resorting to number-theoretic tools.

Lemma 5. *Let $d = 2k + 1$. Then, (1) for μ in $P\overline{P} = \mu I$ is solved by $x = \mu^{-k} \det P$.*

Proof. Let $\lambda := \det P$. Then $\det(P\overline{P}) = \lambda\overline{\lambda} = \overline{\lambda}\lambda = \det(\mu I) = \mu^{2k+1}$. Thus $\mu = \overline{\mu} = \frac{\lambda}{\mu^k} \frac{\overline{\lambda}}{\mu^k}$. Replacing P with $P' = \frac{\mu^k}{\lambda}P$ we see that $P'\overline{P'} = I$. \square

Related work and remarks

The paper [Fie09] studies a closely related algorithmic question of minimising the degree of the number field needed to write down a complex representation. It is known that such a field need not be cyclotomic. On the other hand, computer algebra systems designed for computing in groups, such as GAP [GAP21] and Magma [BCP97] typically use cyclotomic fields for computation with characteristic zero representations of finite groups. In particular, this work came as an analysis of a question [Ros21] posed on the GAP discussion forum.

Lemma 2 and its proof are essentially a refinement of an argument from the proof of [Ser77, Thm.31]. Lemmata 5 and 4 appear to be novel, as well as Theorem 1.

Acknowledgement.

The author thanks Denis Rosset, Stephen Glasby, Kaashif Hymabaccus, Robert Guralnick, and many participants of MathOverflow, in particular Will Sawin [Saw21], for fruitful discussions.

Special thanks go to Robert Guralnick, who, in his capacity of a moderator for arXiv.org, caught a glaring omission in an earlier version of this text, submitted there.

References

- [BCP97] Wieb Bosma, John Cannon, and Catherine Playoust. The Magma algebra system. I. The user language. *J. Symbolic Comput.*, 24(3-4):235–265, 1997. Computational algebra and number theory (London, 1993).
- [CR87] Charles W. Curtis and Irving Reiner. *Methods of representation theory. Vol. II.* Pure and Applied Mathematics (New York). John Wiley & Sons, Inc., New York, 1987. With applications to finite groups and orders, A Wiley-Interscience Publication.
- [CR06] Charles W. Curtis and Irving Reiner. *Representation theory of finite groups and associative algebras.* AMS Chelsea Publishing, Providence, RI, 2006. Reprint of the 1962 original.
- [DD10] Vahid Dabbaghian and John D. Dixon. Computing matrix representations. *Math. Comp.*, 79(271):1801–1810, 2010.
- [Dix93] John D. Dixon. Constructing representations of finite groups. In *Groups and computation (New Brunswick, NJ, 1991)*, volume 11 of *DIMACS Ser. Discrete Math. Theoret. Comput. Sci.*, pages 105–112. Amer. Math. Soc., Providence, RI, 1993.
- [Fie09] Claus Fieker. Minimizing representations over number fields. II. Computations in the Brauer group. *J. Algebra*, 322(3):752–765, 2009.
- [GAP21] The GAP Group. *GAP – Groups, Algorithms, and Programming, Version 4.11.1*, 2021.
- [GH97] S. P. Glasby and R. B. Howlett. Writing representations over minimal fields. *Comm. Algebra*, 25(6):1703–1711, 1997.
- [HP20] Kaashif Hymabaccus and Dmitrii V. Pasechnik. RepnDecomp: A GAP package for decomposing linear representations of finite groups. *Journal of Open Source Software*, 5(50):1835–1836, 2020.
- [Isa94] I. Martin Isaacs. *Character theory of finite groups.* Dover Publications, Inc., New York, 1994. Corrected reprint of the 1976 original [Academic Press, New York; MR0460423 (57 #417)].
- [Ros21] Denis Rosset. Repsn: constructing representations with real coefficients, 2021. GAP-Forum post, <https://mail.gap-system.org/pipermail/forum/2021/006240.html>.

- [Saw21] Will Sawin. Is every positive real cyclotomic number the norm of a cyclotomic? MathOverflow, 2021. <https://mathoverflow.net/q/391269> (version: 2021-04-27).
- [Ser71] Jean-Pierre Serre. Conducteurs d'Artin des caractères réels. *Invent. Math.*, 14:173–183, 1971.
- [Ser77] Jean-Pierre Serre. *Linear representations of finite groups*. Springer-Verlag, New York, 1977. Translated from the second French edition by Leonard L. Scott, Graduate Texts in Mathematics, Vol. 42.