

Comments on potential health effects of MRI-induced DNA lesions: quality is more important to consider than quantity

M.A. Hill*, P. O'Neill, and W.G. McKenna

CRUK/MRC Oxford Institute for Radiation Oncology, University of Oxford, Gray Laboratories, ORCRB Roosevelt Drive, Oxford OX3 7DQ, UK

Received 15 February 2016; accepted after revision 13 July 2016; online publish-ahead-of-print 22 August 2016

Magnetic resonance imaging (MRI) is increasingly being used in cardiology to detect heart disease and guide therapy. It is mooted to be a safer alternative to imaging techniques, such as computed tomography (CT) or coronary angiographic imaging. However, there has recently been an increased interest in the potential long-term health risks of MRI, especially in the light of the controversy resulting from a small number of research studies reporting an increase in DNA damage following exposure, with calls to limit its use and avoid unnecessary examination, according to the precautionary principle. Overall the published data are somewhat limited and inconsistent; the ability of MRI to produce DNA lesions has yet to be robustly demonstrated and future experiments should be carefully designed to optimize sensitivity and benchmarked to validate and assess reproducibility. The majority of the current studies have focussed on the initial induction of DNA damage, and this has led to comparisons between the reported induction of γ H2AX and implied double-strand break (DSB) yields produced following MRI with induction by imaging techniques using ionizing radiation. However, γ H2AX is not only a marker of classical double-ended DSB, but also a marker of stalled replication forks and in certain circumstances stalled DNA transcription. Additionally, ionizing radiation is efficient at producing complex DNA damage, unique to ionizing radiation, with an associated reduction in repairability. Even if the fields associated with MRI are capable of producing DNA damage, the lesions produced will in general be simple, similar to those produced by endogenous processes. It is therefore inappropriate to try and infer cancer risk by simply comparing the yields of γ H2AX foci or DNA lesions potentially produced by MRI to those produced by a given exposure of ionizing radiation, which will generally be more biologically effective and have a greater probability of leading to long-term health effects. As a result, it is important to concentrate on more relevant downstream end points (e.g. chromosome aberration production), along with potential mechanisms by which MRI may lead to DNA lesions. This could potentially involve a perturbation in homeostasis of oxidative stress, modifying the background rate of endogenous DNA damage induction. In summary, what the field needs at the moment is more research and less fear mongering.

Keywords MRI • DNA damage • ionizing radiation • cancer risk

Introduction

Magnetic resonance imaging (MRI) is a widely used diagnostic technique capable of acquiring both anatomical and functional information on organs within the body. It is increasingly being used in cardiology to detect heart disease and guide therapy. It is mooted to be a safer alternative to x-ray and radioisotope imaging techniques, such as computed tomography (CT) or coronary angiographic imaging, where there is a clear relationship between exposure to ionizing radiation and increased cancer risk, as recently demonstrated by epidemiological studies for paediatric CT exposures.^{1,2} However, with the increase in the clinical use of MRI, there has

also been an increased interest in the potential risks, especially in the light of the controversy resulting from a limited number of studies reporting an increase in DNA damage induction following exposure,^{3–5} with calls to limit its use and avoid unnecessary examination, according to the precautionary principle.^{6–9}

MRI results in exposure of the patient to a mixture of static magnetic fields (SMF), time-varying gradient magnetic fields (GMF) typically in the range 10–100 kHz, and pulsed radiofrequency fields (RF) in the MHz range. While for occupational exposures of staff working with or in close proximity to the scanner, SMF is the most relevant. The known risks of using MRI mainly relate to the interaction of magnetic and radiofrequency fields with

* Corresponding author. Tel: +44 0 1865 617056. E-mail: mark.hill@oncology.ox.ac.uk

© The Author 2016. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

ferromagnetic and paramagnetic objects (limiting its use on patients with metal objects within the body) and that movement of a person through a strong SMF may induce effects such as vertigo and nausea.¹⁰ In addition, RF may cause some heating of tissue, and GMF are capable of stimulating nerves and muscles; however, in practice, the operation limits are set to minimize these effects. With regard to biological and long-term health effects associated with MRI exposure and its various components, a number of comprehensive reviews have been published by national and international committees^{10–16} along with a recent review of genotoxicity associated with MRI exposures.⁴ Although the data are somewhat limited, the current consensus is that no clear link exists between MRI or associated magnetic and pulsed radiofrequency fields and subsequent health risks. The relevant data are discussed below. We conclude that it is essential to avoid over interpretation of the limited data available on potential genetic effects of MRI and making uninformed comparisons of these effects with the kinds of DNA damage produced by ionizing radiation.

Does MRI exposure lead to DNA damage?

The potential of adverse effects associated with exposure to magnetic fields has been studied for a number of years both *in vitro* and *in vivo*. DNA damage, especially DNA double-strand break (DSB) and chromosome aberrations (CA), has typically been a main focus of research, since DSB and CA have the greatest potential to lead to carcinogenesis or hereditary effects. The effects of exposure to individual fields, SMF, GMF, or RF have been extensively reviewed by various national and international committees^{10–16} who typically concluded that there is no clear link between exposure and an increase in genetic damage. However, current MRI exposures are a complex combination of SMF, GMF, and RF. Only a very limited number of investigations of genetic damage using clinical MRI scanners, and the types of sequences routinely used in the clinic, have been published and are briefly summarized in Table 1. Seven of these studies were also recently critically reviewed by Vijayalaxmi *et al.*⁴ What is clear is that the experimental data relating to MRI genotoxicity are limited and inconsistent.

DNA single-strand breaks

Three of the studies investigated the induction of single-strand breaks (SSB) using the alkaline comet assay. The study of Szerencsi *et al.*²¹ showed no enhancement immediately following *in vitro* exposure of human leukocytes. In contrast, statistically significant enhancements were observed by Lee *et al.*¹⁹ following *in vitro* exposure times of 45 min or greater, with the observed level increasing with duration of exposure. For *in vivo* exposures, Yildiz *et al.*²⁰ observed no significant increase in lymphocytes taken after the initial non-contrast-enhanced MRI. However, they did report a significant increase in the lymphocytes taken following the subsequent contrast-enhanced MRI.

DNA double-strand breaks

Five studies investigated DSB induction using γ H2AX assays (a DSB along with several other modifications results in the phosphorylation of surrounding H2AX histones). Three of the studies

showed no enhancement of DSB following either *in vitro* exposure of human cancer cell lines 0, 1, or 24 h post exposure,¹⁸ *in vitro* exposure of human lymphocytes 0, 1, or 20 h post exposure,²³ or in human lymphocytes taken 5 min after *in vivo* exposure of patients undergoing contrast-enhanced cardiac magnetic resonance imaging (CMR).²² In contrast, Fiechter *et al.*⁸ reported a statistically significant enhancement in DSB in human lymphocytes taken from patients immediately following contrast-enhanced CMR. While Lancellotti *et al.*⁷ reported no enhancement in T lymphocytes taken from volunteers 1 and 2 h post exposure to unenhanced CMR, but significant enhancement was reported at day 2 and 1 month post exposure before returning to base line levels after 1 year (a significant increase was observed at 2 h and day 2 for NK cells but not at other time points).

Chromosome aberrations/micronuclei

While Szerencsi *et al.*²¹ showed no increase in the yield of micronuclei (MN) following *in vitro* exposure of human lymphocytes, a significant enhancement of both MN and CA (using Giemsa staining) was reported by Lee *et al.*¹⁹ for *in vitro* exposure times of 45 min and greater, with the observed level increasing with duration of exposure. The most frequent types of aberrations observed were chromatid breaks, with both exchange-type and deletion-type CA being observed for the longest exposure time (89 min). A significant increase in MN frequency was also reported by Simi *et al.*⁹ in lymphocytes at 0 h post *in vitro* exposure, with the frequency increasing with increasing duration of exposure. While for the two shorter exposures no significant increase in MN frequency above controls was observed for cells left for 24 h to repair before performing the assay, a significant increase was still observed for the two longer exposure times. Following *in vivo* exposure, a significant increase was observed for all participants when blood was collected immediately or 24 h post exposure, although no difference was observed at later collection times (72, 94, and 120 h).

Mutations

Single experiment using the Ames test shows no mutagenic or co-mutagenic effect.¹⁷

DNA assays and relevance of DNA lesions to cancer?

Cells have several well-developed DNA-repair pathways and DNA-damage checkpoints to deal with the frequent DNA lesions produced by endogenous or exogenous sources and the vast majority of lesions will be repaired. However, the efficiency and fidelity of repair, and therefore the relevance, will be dependent on the type of lesion produced. Accurate quantification of these lesions is also not trivial, especially at low levels; therefore, great care must be taken to ensure consistency and with appropriately interpreting the results.

The standard alkaline comet assay detects strand breaks along with alkali-labile sites, although the technique can also be modified to visualize the presence of base damage by using lesion-specific endonucleases that convert base damage into SSB.²⁴ However, these lesions are typically repaired quickly with high fidelity and so are much less likely to be relevant to long-term health effects compared with DSB.^{25,26}

Table 1 Investigations of genetic damage using clinically relevant scanners or pulse sequences

	Schreiber et al. ¹⁷	Schwenzer et al. ¹⁸	Simi et al. ⁹	Lee et al. ¹⁹	Yildiz et al. ²⁰	Fiechter et al. ⁸	Szerencsi et al. ²¹	Lancellotti et al. ⁷	Brand et al. ²²	Reddig et al. ²³
Assay	Mutation (Ames test)	DSB (γH2AX: FC and foci)	MN	CA, MN, and SSB (alkaline comet)	SSB (alkaline comet)	DSB (γH2AX: FC and foci)	SSB (alkaline comet); MN	DSB (γH2AX: FC)	DSB (γH2AX: foci)	DSB (γH2AX: FC and foci)
Study	<i>In vitro</i>	<i>In vitro</i>	<i>In vitro</i> and <i>in vivo</i>	<i>In vitro</i>	<i>In vivo</i>	<i>In vivo</i>	<i>In vitro</i>	<i>In vivo</i>	<i>In vivo</i>	<i>In vitro</i>
Flux	1.5 T and 7.2 T	3 T	1.5 T	3 T	1.5 T	1.5 T	3 T	1.5 T	1.5 T	7 T
Scan protocol	Static only; static (1.5 T) + time-varying bipolar GMF; static (1.5 T) + pulsed RF	Static only; Static + turbo spin-echo (TSE); Static + gradient-echo (GE)	Cardiac	Brain: range of pulse sequences	Hypophysial	Cardiac	Brain: range of pulse sequences	Cardiac: range of pulse sequences	Cardiac: range of pulse sequences	Static only; Static with varying GMF and pulsed RF
Scan duration	1.5 sT: 1 h and 24 h 7.2 T: 1 h	TSE: 2 min 20s or 2 h	<i>In vitro</i> : 686, 1186, 1618, 2188s	22, 45, 67, 89 min	~16 min	68 ± 22 min	22, 45, 67, 89 min	35–40 min	30–60 min	1 h
Contrast agent	–	–	No contrast agent	–	With and without gadolinium	Gadolinium	–	No contrast agent	Gadolinium	–
Cells	<i>Salmonella typhimurium</i> bacteria	Human cancer cells (HL-60 and KG-1a)	Human blood lymphocytes	Human blood lymphocytes	Human blood lymphocytes	Human blood lymphocytes	Human blood lymphocytes	Human blood (T lymphocytes and NK cells)	Human blood lymphocytes	Human blood lymphocytes
Expts/Donors	≥ 2 expts	–	<i>In vitro</i> : 8 healthy donors <i>In vivo</i> : 8 patients and volunteers	Single healthy donor	28 patients	20 patients	2 healthy donors/3 repeats	20 healthy donors	45 patients	16 healthy donors
Temp	1.5 T: 32–37°C 7.2 T: 37°C	37°C	<i>In vitro</i> : Room temp <i>In vivo</i> : body temp	25°C	Body temp	Body temp	20°C	Body temp	Body temp	–
Assay time points	Colonies counted after 48 h	0, 1, and 24 h post imaging	PHA stimulation: <i>In vitro</i> at 0 h and 24 h post scan; <i>In vivo</i> at 0, 24, 72, 96, 120 h post scan	PHA stimulation prior to exposure. SSB: 0 h post exposure MN: 0 h CA: 0 h	0 h post non-contrast scan; 0 h post subsequent contrast-enhanced scan	0 h post imaging	SSB: 0 h post exposure; MN: 0 h PHA stimulation (following 20 min transport)	1 h, 2 h, 2 days, 1 month and 1 year post imaging	5 min post imaging	0, 1, 20 h post imaging
Positive control	Chemical mutagens	4 Gy 6MV x-rays	–	SSB: cisplatin MN, CA: bleomycin	–	–	4 Gy γ-ray	–	–	120 kV CT scan and 0.2 Gy γ-rays
Results	No mutagenic or co-mutagenic effect observed	No significant increase observed	<i>In vitro</i> : Significant increase at 0 h, frequency increasing with exposure time. Reduced after 24 h (2 highest exposures still significant) <i>In vivo</i> : Significant increase after 0 h and 24 h but not later times	SSB, MN, CA: significant increase with increasing exposure times of 45 min and above	No significant increase after non-contrast-enhanced MRI; Significant increase following subsequent contrast-enhanced MRI	DSB (FC, foci): Significant increase observed	SSB, MN: No significant increase observed	T lymphocytes: 1 h, 2 h, 1 year—no significant effect; 2 days, 1 month—significant increase; NK cells: a slight increase was observed at 2 h and day 2	No significant increase observed	No significant increase observed

SSB, assessed using the alkaline comet assay; DSB, assessed using γH2AX measured using either flow cytometry (FC) or immunofluorescent microscopy and counting resultant foci (foci); MN assay, following stimulation with phytohaemagglutinin (PHA), cells incubated for 72 h, with cytochalasin-B added after 44 h; CA assay, following stimulation with PHA, cells incubated for 48 h, with colcemid added after 45 h.

DNA DSB and particularly complex DSB (see below) are considered to be particularly important with respect to carcinogenesis as they are intrinsically more difficult to repair than other types of DNA damage with their potential to lead to modification, rearrangement, or loss of chromosomal material.²⁷ An early response to DSB is the phosphorylation of the histone H2AX, which can be detected using a fluorescent antibody in conjunction with immunofluorescent microscopy to visualize discrete nuclear foci at sites of DSB.^{28,29} Radiation-induced foci have been observed following exposure to ionizing radiation with doses as low as 1 mGy in stationary phase human fibroblasts (~ 0.02 – 0.04 DSB per cell), with the yield increasing linearly with dose.³⁰ Studies have also demonstrated that the yield of γ H2AX foci can be a useful quantitative biomarker of human low-level ionizing radiation exposure in blood taken from an exposed individual at doses down to 6 mGy.³¹ This technique has significantly greater sensitivity in detecting DSB than direct methods which require many hundreds of DSB per cell, but it is also associated with several shortcomings. For instance, studies using Pulse Field Gel Electrophoresis (PFGE) demonstrate that the initial DSB produced are quickly repaired with a half time of ~ 20 min, with only a few remaining after 1 h.^{32,33} However, γ H2AX is a marker of the metabolic activities initiated to repair the damage, rather than a direct marker of DSB. Although a few laboratories are able to observe γ H2AX foci within minutes post exposure, their detection is difficult at these early times due to their small size, and they are more reliably scored 15–30 min post exposure. Many laboratories report an initial increase in foci number with time, reaching a maximum ~ 30 – 60 min after exposure, with a reduction being observed at later times. Therefore, the initial yield of DSB may be underestimated due to many of the DSB being repaired during this period; additionally the subsequent rate of DSB repair reported at these later times is often slower than the initial rate of DSB repair.^{33–35} It has been proposed that γ H2AX may, in some cases, continue to mark the site following the initial rejoining of the break and therefore may not always represent a physical break.^{33,36} γ H2AX also detects replication-induced DSB in S-phase cells which will contribute to the persistence or delayed response of the signal. Even non-exposed S-phase cells typically have a higher number of background foci than other phases due to endogenously induced damage.^{26,37} Replication-induced DSB are also very different chemically to double-ended DSB and potentially much more likely to be repaired; it is therefore not clear whether these secondary foci always reflect the presence of DSB.³⁷ γ H2AX may also in some circumstances be generated by DNA transcriptional activity.^{38–40} It is important to note that the sensitivity, reliability, and absolute yields obtained with this technique can vary significantly between laboratories depending on many factors such as staining protocol, cell type, cell cycle, temperature, and critically on the imaging and scoring criteria used. This is illustrated by the large variation in results obtained for the 20 patients by Fiechter *et al.*,⁸ with the median number of γ H2AX per cell varied from 0 to 0.661 pre-exposure and 0 to 1.065 post-exposure; in contrast, the mean number of foci per cell obtained for 45 patients by Brand *et al.*²² is significantly smaller and much more consistent, varying from ~ 0.09 to 0.17 and consistent with background levels of < 0.2 foci per lymphocyte reported for ionizing radiation biodosimeter studies.^{31,41}

Flow cytometry can also be used to analyse γ H2AX with the advantage that thousands of cells can be analysed very quickly, through measuring the total fluorescence intensity rather than detecting individual foci. In general though the sensitivity of this approach is lower than that achieved by scoring individual foci, in part due to background foci intensity having a wider inter-individual variability.^{42,43} As a result, this technique does not maintain the level of sensitivity required to be used as a biodosimeter of ionizing radiation exposure beyond the first hours.⁴¹ The flow cytometry results of Fiechter *et al.*⁸ and Lancellotti *et al.*⁷ both show significant variation between individuals, with some showing an increase while others a decrease in response. The delayed enhancement in γ H2AX expression reported by Lancellotti *et al.*⁷ at day 2 and 1 month post exposure is inconsistent with DSB being produced during imaging as the majority of DSB are reported to be repaired within hours and the limited sensitivity of the flow cytometry assay at extended times following exposure. This could reflect an enhancement in the background level of DSB production (or other events leading to increased γ H2AX expression) post exposure. However, the time course of this increase is still greater than the delayed peak of several hours often observed following exposure of exponentially growing cells to genotoxic agents such as UV and alkylating agents.⁴⁴

When assessing the relevance of the studies investigating the induction of DNA damage such as DSB, it is important to remember that the production of such lesions does not necessarily lead to long-term consequences. In general, the vast majority of DNA damage produced will be transient because of repair processes. As a result, it is the longer term consequences of this damage that are more critical, the unrepaired fraction and the resultant genetic modifications. A more relevant end point to study would be CA production. These have been extensively used (dicentric aberrations in particular) in peripheral blood lymphocytes for assessment of ionizing radiation, both as a biological dosimeter of exposure^{45,46} and as a marker of relative risk to different qualities of radiation [The effectiveness of different qualities/types of radiation (e.g. x-rays, α -particles, and neutrons) at inducing a given biological end point for a given absorbed dose can vary significantly.] exposure.^{47,48} In addition, significant evidence exists for the role of CA in early stages of cancer, with data showing that the frequency of CA in peripheral lymphocytes may be a predictor of cancer incidence in human populations.^{49–52} MN may also be used as measure of CA induction and may result from acentric chromosome/chromatid fragments or whole chromosomes failing to be incorporated into the daughter nuclei. Evidence indicates a link between increased MN and elevated cancer risks.⁵³ In addition to the dicentric chromosome assay, the MN assay has, to a lesser extent, been established as a technique for biodosimetry for ionizing radiation. These have required careful optimization of the sample preparation, data analysis protocols, and harmonization of procedures between laboratories to ensure consistency. However, the sensitivity of these techniques limits their usefulness as a way of retrospectively determining dose in exposures of the order of 100 mGy or above (> 2 DSB per cell). Therefore, investigating the induction of CA or MN in situations where there are fewer DSB per cell becomes increasingly challenging, requiring more cells to be scored and great care to ensure reliability of the data. Dicentric aberrations and MN produced by ionizing radiation are essentially stable and can be detected for several months following irradiation, with their loss

being directly related to the rate of turnover of cells within the blood. It is therefore interesting that although Simi *et al.*⁹ reported an increase in MN in lymphocytes in blood collected immediately after *in vivo* exposure, no difference was observed compared with controls at later collection times (72, 94, and 120 h). Although lymphocytes within the blood are typically in G0 phase, the reported dominance of chromatid aberrations reported by Lee *et al.*¹⁹ suggests that rearrangements occur mainly during S-phase. To investigate CA/MN induction, the cells were stimulated to divide. It is therefore possible that the DNA lesions resulting in CA/MN were produced during DNA replication, potentially by replication-induced breaks/DSB resulting from stalled replication forks. So rather than MRI directly producing DNA lesions, it may perturb the homeostasis within the cells for a period of time resulting in an enhancement of endogenous damage.

Endogenous DNA damage

DNA damage is a constant and a very frequent event in the daily life of all cells. Every cell within the body has a background level of the order of at least 50 000 endogenous DNA lesions per day as a result of reactive oxygen species (ROS) and other reactive metabolites, with the number expected to increase in situations with increasing oxidative stress.^{54,55} ROS include superoxide, hydrogen peroxide, hydroxyl radicals, and singlet oxygen and they can oxidize DNA, which can lead to several types of DNA damage chemically similar to those induced by ionizing radiation, although importantly their spatial and temporal distribution will be very different. These lesions are constantly being repaired with high fidelity by the cell via highly effective repair processes which help maintain genome integrity, minimizing the onset of ageing, and tumourigenesis. However, it has been proposed that these ubiquitous background events may account for an important fraction of oncogenic events in humans.⁵⁶ Identification of these lesions and their production by free radical mechanisms have been the subject of numerous reviews.^{57–60} Although few, if any, DSB are formed by endogenous processes,^{61,62} events such as DNA damage lesions may lead to stalled replications forks during DNA replication (Figure 1) or interfere with DNA transcription, both of which can be detected by the γ H2AX assay as mentioned earlier.^{26,37,39,40}

It is well known that within the body cells do not act in isolation, with intercellular signalling being vital for maintaining the multicellular organization of tissue and the normal functioning of the constituent cells. ROS along with reactive nitrogen species (RNS) are formed as a consequence of both normal cell metabolism and inflammation and as a result of exposure to environmental factors. Free radicals and resulting reactive non-radical species are important for many physiological processes, including immunological defence, intra- and intercellular signalling, their concentration being determined by the balance between the rate of production and clearance by various antioxidants. As a result, there is a delicate balance between the advantageous and detrimental effects of these radicals.⁶³ Environmental factors can lead to the activation of endogenous ROS or RNS-generating systems resulting in either a temporary or persistent dysregulation in signalling cascades and a change in ROS and/or RNS concentrations.⁶³ This perturbation in homeostasis of this signalling and associated changes in oxidative




Figure 1 Schematic illustrating the production of stalled replication fork which can form γ H2AX foci. A replication fork is formed during DNA replication, the two original strands branch forming single strands and serve as templates for synthesis of the complementary strand which is synthesized in the 5' to 3' direction. The presence of a lesion, such as a SSB, can result in the replication fork stalling which can result in γ H2AX foci.

stress can lead to a modulation of the background rate of endogenous DNA damage induction. For example, it is well known that very low doses of ionizing radiation can perturb intercellular signalling, levels of ROS, and other signalling molecules and can result in genotoxic effects such as induction of γ H2AX foci, chromatid aberrations, MN, and apoptosis in unirradiated cells.^{64–67} Interestingly, rather than a continuous increase in effect with increasing exposure, the observed response is often consistent with a switch between two stable states essentially independent of triggering dose (if above a low threshold dose). It has been proposed that MR exposure may be able to increase the rate of production of DNA damage by endogenous processes as a result of an increase in oxidative stress.^{68,69}

Assuming MRI can induce DNA lesions, the kinetics of induction and repair of any such lesions are also unclear from current studies but remain a crucial question that needs to be addressed. Are these lesions formed directly during imaging where some of the lesions may persist to later times, or are they formed through indirect processes that continue after imaging with a constant turnover as lesions are produced and repaired? This information would be helpful in guiding the understanding of the mechanisms, such as changes in homeostasis or persistent damage. A potential approach to address these questions would be to use live cell imaging using fluorescently labelled DNA-repair proteins and cell cycle markers, to identify the kinetics of formation and repair of individual lesions and cell cycle dependence. The relevance of any short-term increase in DNA damage as a result of an enhancement of endogenous processes has to be considered in the context that similar sites of DNA damage will be continuously produced throughout the lifetime of an individual and therefore any lesions, even if induced by MRI exposure, may not significantly add to the burden of risk associated with the background level of endogenous damage.

Unique nature of DNA damage produced by ionizing radiation

The majority of recent experiments in investigating genotoxicity of MRI use the γ H2AX assay as a measure of DSB. It is then tempting to compare induction of γ H2AX and implied DSB yields produced following MRI with induction by imaging techniques using ionizing

radiation.³ However, due to the way ionizing radiation interacts within the body, the resulting DNA damage produced includes damage unique to ionizing radiation, such as clustered DNA damage, and therefore, it is inappropriate to infer long-term health risks based simply on a comparison of yields of DSB or other non-DSB lesions. It is essential to deconvolute the γ H2AX data into different classes of lesions and associated lifetimes, to be able to comment on the risks of MRI to humans.

These assertions are founded on the knowledge that ionizing radiation interacts within the body and deposits energy along the resulting electron tracks producing highly structured tracks of ionization and excitation events that are stochastic in nature. This produces a wide variety of molecular damage, including base damage, abasic sites, SSB, and DNA–protein cross links. However, because these electrons produce multiple energy deposition sites along the resulting radiation track on the nanometre scale (correlated in time and space), it is capable of producing sites with multiple lesions on the DNA within a few base pairs of each other (Figure 2). It is the ability of ionizing radiation to produce clustered DNA damage which make it so biologically effective, with these longer lived clustered lesions dominating the long-term response rather than the much more numerous isolated DNA lesions that are chemically similar to those induced endogenously and are rapidly repaired. In mammalian cells, an exposure of 1 Gy of x-rays or γ -rays will produce ~ 1300 base lesions, 1000 SSB, and 20–40 DSB (see Table 2 for corresponding number for a typical chest CT scan). Although the number of lesions produced by ionizing radiation is significantly less than those produced daily by endogenous processes ($\sim 50\,000$ per cell per day), it is the clustering of multiple lesions at a site of DNA damage^{71,72} and the corresponding reduced ability of their repair that is critical, leading to an increase in genetic modification (clustered lesions produced by endogenous damage are rare and do not match the degree of complexity that can be produced by ionizing radiation^{73,74}). Although the majority of ionizing radiation-induced DSB will be repaired within ~ 60 min, a small fraction may persist for > 24 h.^{75–78} Theoretical analysis of these DSB using Monte Carlo calculations^{79–82} indicates that 20–50% of these DSB are in fact complex DSB (with extra single-strand breaks and/or associated base damage within a distance of 10 base pairs).⁸⁰ In fact, the frequency and complexity of DSB typically increase with ionization density of the radiation and are important factors in accounting for differences in relative biological effectiveness (RBE) with radiation quality.⁸⁰ This is also reflected in the variation in biological efficiency per DSB produced as a function of radiation quality, with even the DSB produced by sparsely ionizing x-rays estimate to be 4–40 times more effective than simple DSB produced by high concentrations of hydrogen peroxide.⁸³ Additionally, as DSB are formed along the path of the resulting electron tracks following exposure, the DSB produced can be correlated in space and time increasing the chance that they may interact and lead to genetic and chromosomal rearrangements.⁸⁴

In addition to DSB and complex DSB, ionizing radiation will also form non-DSB clustered lesions. These non-DSB clustered lesions formed contain two or more lesions within the order of 10 base pairs and may occur either on the same strand (tandem) or opposite strands (bistranded). Complex non-DSB damages have been shown to make up a significant component of the lesions induced by




Figure 2 Schematic showing a typical track produced by the ubiquitous low-energy electron track-ends produced in a cell by ionizing radiation (dark circle = ionization event; light circle = excitation event). Individual DNA lesions are produced either by direct interaction with DNA or indirectly as a result of reactive radicals (most notably hydroxyl radicals) produced in the surrounding water (diffusion distance ~ 6 nm). Clustered DNA damage, comprising combinations of two or more strand breaks and/or base damage, produced within one to two helical turns.

Table 2 Yield of major lesions induced per mammalian cell following a 1 Gy exposure of x-rays (adapted from Cadet et al.⁵⁸ and Lomax et al.⁷⁰) and corresponding number for exposed cell following a typical adult abdominal CT

Radiation-induced DNA lesions	Number/Gy/cell	Adult abdominal CT (10 mGy to stomach)
Pyrimidine lesions	850	8.5
Purine lesions	450	4.5
Single-strand breaks (SSB)	1000	10
Double-strand breaks (DSB)	20–40	0.2–0.4

radiation, being at least 4–8 times greater in number than prompt DSB for x-rays and γ -rays.^{71,72,85–88} While isolated lesions (e.g. base damage or SSB) are repaired quickly with high fidelity, for non-DSB clusters the rate of repair is typically impaired by the presence of additional lesions within the cluster.^{89–91} The extent of the reduction is dependent on the types and number of lesions along with their separation and orientation.⁸⁹ Although processing of certain bistranded lesions may in a few cases lead to repair-induced DSB, the most likely consequence of these non-DSB clustered lesions will be an increase in their lifetime, and therefore, an increased chance of resulting in a stalled replication fork and potentially lead to replication-induced DSB in S-phase (Figure 1). It has been estimated that $\sim 10\%$ of non-DSB clustered lesions will be converted to DSB as a result of processing,³² with the possibility that some of these DSB may be complex by virtue of additional lesions close by.^{72,92} Experiments have shown that these clustered lesions result in an

increase in mutation frequencies in bacteria, yeast, and mammalian cells.^{89,93,94}

Little is known about the mechanisms by which MRI can produce DNA damage, if indeed it does. It is known that the fields associated with MRI are non-ionizing and therefore unable to produce free electrons with sufficient energy to produce clustered damage sites over 1–2 turns of the DNA, similar to that associated with ionizing radiation. Therefore, any lesions that are produced as a result of MRI are much more likely to be repaired with high fidelity than those produced by ionizing radiation.

Conclusions

Although MRI imaging is generally considered to be safe compared with imaging technology using ionizing radiation, there is increased concern about the potential long-term health effects of exposure, especially with the push to higher static fields along with stronger and faster switching fields. Although IARC¹⁵ did not find clear evidence that SMF are carcinogenic, the data are limited, underpowered, or suffer methodological weakness.¹¹ Both IARC and the recent SCENIHR¹⁴ report suggest the need for a large, carefully designed epidemiology study.

The ability of MRI to produce DNA damage and potential mechanisms involved remains unclear. Generally no consistent response has been reported between the very limited number of published studies, with limited or no controls, typically based on data from a small number of subjects or often a single experiment and therefore limited statistical power. It is therefore essential that future studies employ strict standardization in experimental design and scoring criteria, along with quality control measures to ensure reproducibility and consistency. All techniques should be carefully benchmarked against negative and positive controls (potentially using ionizing radiation) to confirm the validity and reproducibility of the results, collected over multiple independent experimental measurements and ideally include inter-laboratory comparisons. There must also be a willingness by the scientists and journals to publish negative results for well-designed experiments. While there is a tendency to focus on the initial induction of DNA damage, it is more relevant to study downstream effects, for example CA production for which there is some evidence to this could be a marker of cancer risk. If a reproducible effect can be observed, then these studies will also need to critically assess the importance of contrast agents, along with potential confounding factors and other potential synergistic effects (such as smoking, medication, health issues relating to reason for the scan, and underlying genetic sensitivity). An additional issue to be addressed would be to determine the critical component of exposure (SMF, GMF, RF) if possible to compare responses for different scans between laboratories.

It is important that great care is taken to ensure that any potential genetic effects associated with MRI exposure, if it exists, are not over interpreted. There is a danger that in the absence of clear evidence, physicians and patients are misinformed by the presentation of preliminary data, affecting their choice of potentially life-saving imaging tests. The ability for MRI to produce DNA lesions has yet to be robustly demonstrated. If lesions are formed, then the relative probability of these lesions to produce long-term health effects has

to be put in the context of the risk associated with the 50 000 lesions produced daily by endogenous processes. As discussed, the fields associated with MRI are non-ionizing, having insufficient energy to directly break chemical bonds and are thus unable to produce free electrons with sufficient energy to produce clustered lesions including complex DSB with an associated reduction in repairability. The complex DSB produced by ionizing radiation have been shown to be more effective than simple DSB at producing a wide range of biological end points, including mutations, with a greater probability of leading to long-term health effects. It is therefore important not to try and infer cancer risk by simply comparing the initial yields of DSB produced by MRI to that produced by a given exposure of ionizing radiation. The γ H2AX assay not only marks classical double-ended DSB, but also stalled replication forks and/or collapse in S-phase or interfere with DNA transcription.^{37,40} It would also be appropriate to investigate how imaging may perturb intercellular signalling and associated levels of oxidative stress which may lead to modulation in endogenous damage levels, epigenetic changes, and genomic stability in cells and their progeny. In summary, what the field needs at the moment is more research and less fear mongering.

Acknowledgments

Funding from Medical Research Council Strategic Partnership Funding (MC-PC-12004) for the CRUK/MRC Oxford Institute for Radiation Oncology is gratefully acknowledged.

Conflict of interest: None declared.

References

1. Pearce MS, Salotti JA, Little MP, McHugh K, Lee C, Kim KP et al. Radiation exposure from CT scans in childhood and subsequent risk of leukaemia and brain tumours: a retrospective cohort study. *Lancet* 2012;**380**:499–505.
2. Brenner DJ, Hall EJ. Cancer risks from CT scans: now we have data, what next? *Radiology* 2012;**265**:330–1.
3. Knuuti J, Saraste A, Kallio M, Minn H. Is cardiac magnetic resonance imaging causing DNA damage? *Eur Heart J* 2013;**34**:2337–9.
4. Vijayalaxmi, Fatahi M, Speck O. Magnetic resonance imaging (MRI): a review of genetic damage investigations. *Mutat Res* 2015;**764**:51–63.
5. Berrington de Gonzalez A, Kleinerman RA, McAreavey D, Rajaraman P. Cardiac MR imaging and the specter of double-strand breaks. *Radiology* 2015;**277**:329–31.
6. Hartwig V, Giovannetti G, Vanello N, Lombardi M, Landini L, Simi S. Biological effects and safety in magnetic resonance imaging: a review. *Int J Environ Res Public Health* 2009;**6**:1778–98.
7. Lancellotti P, Nchimi A, Delierneux C, Hego A, Gosset C, Gothot A et al. Biological effects of cardiac magnetic resonance on human blood cells. *Circ Cardiovasc Imaging* 2015;**8**:e003697.
8. Fiechter M, Stehli J, Fuchs TA, Dougoud S, Gaemperli O, Kaufmann PA. Impact of cardiac magnetic resonance imaging on human lymphocyte DNA integrity. *Eur Heart J* 2013;**34**:2340–5.
9. Simi S, Ballard M, Casella M, De Marchi D, Hartwig V, Giovannetti G et al. Is the genotoxic effect of magnetic resonance negligible? Low persistence of micronucleus frequency in lymphocytes of individuals after cardiac scan. *Mutat Res* 2008;**645**:39–43.
10. HPA. Protection of patients and volunteers undergoing MRI procedures. Health Protection Agency, Chilton, UK; 2008.
11. HPA. Static magnetic fields. Report of an independent Advisory Group on Non-ionising Radiation. Health Protection Agency, Chilton, UK; 2008.
12. HPA. Health effects from radiofrequency electromagnetic fields. Report of the Advisory Group on Non-ionising Radiation. Health Protection Agency, Chilton, UK; 2012.
13. ICNIRP. Exposure to static and low frequency electromagnetic fields, biological effects and health consequences (0–100 kHz). ICNIRP Publication 13/2003, Munich, Germany; 2003.

14. SCENIHR. Potential health effects of exposure to electromagnetic fields (EMF); 2015. http://ec.europa.eu/health/ph_risk/committees/04_scenihir/docs/scenihir_o_024.pdf (26 October 2015, date last accessed).
15. IARC. Non-ionizing radiation, Part 1: static and extremely low-frequency (ELF) Electric and magnetic fields. *IARC Monogr Eval Carcinog Risks Hum* 2002;**80**:1–429.
16. IARC. Non-ionizing radiation, Part 2: radiofrequency electromagnetic fields. *IARC Monogr Eval Carcinog Risks Hum* 2013;**102**:1–460.
17. Schreiber WG, Teichmann EM, Schiffer I, Hast J, Akbari W, Georgi H et al. Lack of mutagenic and co-mutagenic effects of magnetic fields during magnetic resonance imaging. *J Magn Reson Imaging* 2001;**14**:779–88.
18. Schwenzer NF, Bantleon R, Maurer B, Kehlbach R, Schraml C, Claussen CD et al. Detection of DNA double-strand breaks using gammaH2AX after MRI exposure at 3 Tesla: an in vitro study. *J Magn Reson Imaging* 2007;**26**:1308–14.
19. Lee JW, Kim MS, Kim YJ, Choi YJ, Lee Y, Chung HW. Genotoxic effects of 3T magnetic resonance imaging in cultured human lymphocytes. *Bioelectromagnetics* 2011;**32**:535–42.
20. Yildiz S, Cece H, Kaya I, Celik H, Taskin A, Aksoy N et al. Impact of contrast enhanced MRI on lymphocyte DNA damage and serum visfatin level. *Clin Biochem* 2011;**44**:975–9.
21. Szerencsi A, Kubinyi G, Valiczko E, Juhasz P, Rudas G, Mester A et al. DNA integrity of human leukocytes after magnetic resonance imaging. *Int J Radiat Biol* 2013;**89**:870–6.
22. Brand M, Ellmann S, Sommer M, May MS, Eller A, Wuest W et al. Influence of cardiac MR imaging on DNA double-strand breaks in human blood lymphocytes. *Radiology* 2015;**277**:406–12.
23. Reddig A, Fatahi M, Friebe B, Guttek K, Hartig R, Godenschweger F et al. Analysis of DNA double-strand breaks and cytotoxicity after 7 Tesla magnetic resonance imaging of isolated human lymphocytes. *PLoS ONE* 2015;**10**:e0132702.
24. Collins AR, Duthie SJ, Dobson VL. Direct enzymic detection of endogenous oxidative base damage in human lymphocyte DNA. *Carcinogenesis* 1993;**14**:1733–5.
25. Dikomey E, Franzke J. DNA repair kinetics after exposure to X-irradiation and to internal beta-rays in CHO cells. *Radiat Environ Biophys* 1986;**25**:189–94.
26. Harper JV, Anderson JA, O'Neill P. Radiation induced DNA DSBs: contribution from stalled replication forks? *DNA Repair* 2010;**9**:907–13.
27. Khanna KK, Jackson SP. DNA double-strand breaks: signaling, repair and the cancer connection. *Nat Genet* 2001;**27**:247–54.
28. Rogakou EP, Boon C, Redon C, Bonner WM. Megabase chromatin domains involved in DNA double-strand breaks in vivo. *J Cell Biol* 1999;**146**:905–16.
29. Rogakou EP, Pilch DR, Orr AH, Ivanova VS, Bonner WM. DNA double-stranded breaks induce histone H2AX phosphorylation on serine 139. *J Biol Chem* 1998;**273**:5858–68.
30. Rothkamm K, Lobrich M. Evidence for a lack of DNA double-strand break repair in human cells exposed to very low x-ray doses. *Proc Natl Acad Sci USA* 2003;**100**:5057–62.
31. Rothkamm K, Balroop S, Shekhdar J, Fernie P, Goh V. Leukocyte DNA damage after multi-detector row CT: a quantitative biomarker of low-level radiation exposure. *Radiology* 2007;**242**:244–51.
32. Gulston M, de Lara C, Jenner T, Davis E, O'Neill P. Processing of clustered DNA damage generates additional double-strand breaks in mammalian cells post-irradiation. *Nucleic Acids Res* 2004;**32**:1602–9.
33. Kinner A, Wu W, Staudt C, Iliakis G. Gamma-H2AX in recognition and signaling of DNA double-strand breaks in the context of chromatin. *Nucleic Acids Res* 2008;**36**:5678–94.
34. Kato TA, Okayasu R, Bedford JS. Comparison of the induction and disappearance of DNA double strand breaks and gamma-H2AX foci after irradiation of chromosomes in G1-phase or in condensed metaphase cells. *Mutat Res* 2008;**639**:108–12.
35. Pilch DR, Sedelnikova OA, Redon C, Celeste A, Nussenzweig A, Bonner WM. Characteristics of gamma-H2AX foci at DNA double-strand breaks sites. *Biochem Cell Biol* 2003;**81**:123–9.
36. Banath JP, Macphail SH, Olive PL. Radiation sensitivity, H2AX phosphorylation, and kinetics of repair of DNA strand breaks in irradiated cervical cancer cell lines. *Cancer Res* 2004;**64**:7144–9.
37. Rothkamm K, Barnard S, Moquet J, Ellender M, Rana Z, Burdak-Rothkamm S. DNA damage foci: meaning and significance. *Environ Mol Mutagen* 2015;**56**:491–504.
38. Cristini A, Park JH, Capranico G, Legube G, Favre G, Sordet O. DNA-PK triggers histone ubiquitination and signaling in response to DNA double-strand breaks produced during the repair of transcription-blocking topoisomerase I lesions. *Nucleic Acids Res* 2016;**44**:1161–78.
39. Sordet O, Redon CE, Guirouilh-Barbat J, Smith S, Solier S, Douarre C et al. Ataxia telangiectasia mutated activation by transcription- and topoisomerase I-induced DNA double-strand breaks. *EMBO Rep* 2009;**10**:887–93.
40. Dickey JS, Baird BJ, Redon CE, Avdoshina V, Palchik G, Wu J et al. Susceptibility to bystander DNA damage is influenced by replication and transcriptional activity. *Nucleic Acids Res* 2012;**40**:10274–86.
41. Horn S, Barnard S, Rothkamm K. Gamma-H2AX-based dose estimation for whole and partial body radiation exposure. *PLoS One* 2011;**6**:e25113.
42. Andrievski A, Wilkins RC. The response of gamma-H2AX in human lymphocytes and lymphocytes subsets measured in whole blood cultures. *Int J Radiat Biol* 2009;**85**:369–76.
43. Ismail IH, Wadhwa TI, Hammarsten O. An optimized method for detecting gamma-H2AX in blood cells reveals a significant interindividual variation in the gamma-H2AX response among humans. *Nucleic Acids Res* 2007;**35**:e36.
44. Staszewski O, Nikolova T, Kaina B. Kinetics of gamma-H2AX focus formation upon treatment of cells with UV light and alkylating agents. *Environ Mol Mutagen* 2008;**49**:734–40.
45. Bauchinger M. Quantification of low-level radiation exposure by conventional chromosome aberration analysis. *Mutat Res* 1995;**339**:177–89.
46. IAEA. Cytogenetic Dosimetry: Applications in Preparedness for and Response to Radiation Emergencies. Vienna: International Atomic Energy Agency; 2011.
47. ICRP. Relative biological effectiveness (RBE), quality factor (Q), and radiation weighting factor (w(R)). ICRP Publication 92. *Ann ICRP* 2003;**33**:1–117.
48. ICRP. Low-dose extrapolation of radiation-related cancer risk. ICRP Publication 99. *Ann ICRP* 2005;**35**:1–140.
49. Hagmar L, Brogger A, Hansteen IL, Heim S, Hogstedt B, Knudsen L et al. Cancer risk in humans predicted by increased levels of chromosomal-aberrations in lymphocytes - Nordic Study-Group on the Health Risk of Chromosome-Damage. *Cancer Res* 1994;**54**:2919–22.
50. Hagmar L, Bonassi S, Stromberg U, Brogger A, Knudsen LE, Norppa H et al. Chromosomal aberrations in lymphocytes predict human cancer: a report from the European study group on cytogenetic biomarkers and health (ESCH). *Cancer Res* 1998;**58**:4117–21.
51. Bonassi S, Abbondandolo A, Camurri L, Dalpra L, Deferrari M, Degraffi F et al. Are chromosome-aberrations in circulating lymphocytes predictive of future cancer onset in humans - preliminary-results of an Italian Cohort Study. *Cancer Genet Cytogen* 1995;**79**:133–5.
52. Bonassi S, Norppa H, Ceppi M, Stromberg U, Vermeulen R, Znaor A et al. Chromosomal aberration frequency in lymphocytes predicts the risk of cancer: results from a pooled cohort study of 22 358 subjects in 11 countries. *Carcinogenesis* 2008;**29**:1178–83.
53. Bonassi S, El-Zein R, Bolognesi C, Fenech M. Micronuclei frequency in peripheral blood lymphocytes and cancer risk: evidence from human studies. *Mutagenesis* 2011;**26**:93–100.
54. Swenberg JA, Lu K, Moeller BC, Gao L, Upton PB, Nakamura J et al. Endogenous versus exogenous DNA adducts: their role in carcinogenesis, epidemiology, and risk assessment. *Toxicol Sci* 2011;**120**(Suppl. 1):S130–45.
55. De Bont R, van Larebeke N. Endogenous DNA damage in humans: a review of quantitative data. *Mutagenesis* 2004;**19**:169–85.
56. Vilenchik MM, Knudson AG. Endogenous DNA double-strand breaks: production, fidelity of repair, and induction of cancer. *Proc Natl Acad Sci USA* 2003;**100**:12871–6.
57. Pouget JP, Frelon S, Ravanat JL, Testard I, Odin F, Cadet J. Formation of modified DNA bases in cells exposed either to gamma radiation or to high-LET particles. *Radiat Res* 2002;**157**:589–95.
58. Cadet J, Douki T, Ravanat JL. Oxidatively generated damage to the guanine moiety of DNA: mechanistic aspects and formation in cells. *Acc Chem Res* 2008;**41**:1075–83.
59. Dizdaroğlu M, Jaruga P. Mechanisms of free radical-induced damage to DNA. *Free Radic Res* 2012;**46**:382–419.
60. Von Sonntag C. *Free-Radical-Induced DNA Damage and its Repair*. Heidelberg: Springer; 2006.
61. Sutherland BM, Bennett PV, Cintron NS, Guida P, Laval J. Low levels of endogenous oxidative damage cluster levels in unirradiated viral and human DNAs. *Free Radic Biol Med* 2003;**35**:495–503.
62. Bennett PV, Cintron NS, Gros L, Laval J, Sutherland BM. Are endogenous clustered DNA damages induced in human cells? *Free Radic Biol Med* 2004;**37**:488–99.
63. Droge W. Free radicals in the physiological control of cell function. *Physiol Rev* 2002;**82**:47–95.
64. Morgan WF, Sowa MB. Non-targeted bystander effects induced by ionizing radiation. *Mutat Res* 2007;**616**:159–64.
65. UNSCEAR. Effects of ionizing radiation, UNSCEAR 2006 report to the general Assembly, with Scientific Annexes: Volume II, Annex C. New York: United Nations; 2008.
66. Hei TK, Zhou H, Ivanov VN, Hong M, Lieberman HB, Brenner DJ et al. Mechanism of radiation-induced bystander effects: a unifying model. *J Pharm Pharmacol* 2008;**60**:943–50.
67. Kadhim M, Salomaa S, Wright E, Hildebrandt G, Belyakov OV, Prise KM et al. Non-targeted effects of ionising radiation--implications for low dose risk. *Mutat Res* 2013;**752**:84–98.
68. Ghodbane S, Lahbib A, Sakly M, Abdelmelek H. Bioeffects of static magnetic fields: oxidative stress, genotoxic effects, and cancer studies. *Biomed Res Int* 2013;**2013**:602987.

69. Consales C, Merla C, Marino C, Benassi B. Electromagnetic fields, oxidative stress, and neurodegeneration. *Int J Cell Biol* 2012;**2012**:683897.
70. Lomax ME, Folkes LK, O'Neill P. Biological consequences of radiation-induced DNA damage: relevance to radiotherapy. *Clin Oncol (R Coll Radiol)* 2013;**25**:578–85.
71. Sutherland BM, Bennett PV, Sidorkina O, Laval J. Clustered DNA damages induced in isolated DNA and in human cells by low doses of ionizing radiation. *Proc Natl Acad Sci USA* 2000;**97**:103–8.
72. Gulston M, Fulford J, Jenner T, de Lara C, O'Neill P. Clustered DNA damage induced by gamma radiation in human fibroblasts (HF19), hamster (V79-4) cells and plasmid DNA is revealed as Fpg and Nth sensitive sites. *Nucleic Acids Res* 2002;**30**:3464–72.
73. Stewart RD. On the complexity of the DNA damage created by endogenous processes. *Radiat Res* 1999;**152**:101–4.
74. Bennett PV, Cuomo NL, Paul S, Tafrov ST, Sutherland BM. Endogenous DNA damage clusters in human skin, 3-D model, and cultured skin cells. *Free Radic Biol Med* 2005;**39**:832–9.
75. Schmid TE, Dollinger G, Beisker W, Hable V, Greubel C, Auer S et al. Differences in the kinetics of gamma-H2AX fluorescence decay after exposure to low and high LET radiation. *Int J Radiat Biol* 2010;**86**:682–91.
76. Botchway SW, Stevens DL, Hill MA, Jenner TJ, O'Neill P. Induction and rejoining of DNA double-strand breaks in Chinese hamster V79-4 cells irradiated with characteristic aluminum K and copper L ultrasoft X rays. *Radiat Res* 1997;**148**:317–24.
77. Asaithamby A, Uematsu N, Chatterjee A, Story MD, Burma S, Chen DJ. Repair of HZE-particle-induced DNA double-strand breaks in normal human fibroblasts. *Radiat Res* 2008;**169**:437–46.
78. Jenner TJ, deLara CM, O'Neill P, Stevens DL. Induction and rejoining of DNA double-strand breaks in V79-4 mammalian cells following gamma- and alpha-irradiation. *Int J Radiat Biol* 1993;**64**:265–73.
79. Nikjoo H, Goodhead DT, Charlton DE, Paretzke HG. Energy deposition in small cylindrical targets by monoenergetic electrons. *Int J Radiat Biol* 1991;**60**:739–56.
80. Goodhead DT. Energy deposition stochastics and track structure: what about the target? *Radiat Prot Dosimetry* 2006;**122**:3–15.
81. Nikjoo H, Bolton CE, Watanabe R, Terrissol M, O'Neill P, Goodhead DT. Modeling of DNA damage induced by energetic electrons (100 eV to 100 keV). *Radiat Prot Dosimetry* 2002;**99**:77–80.
82. Nikjoo H, O'Neill P, Goodhead DT, Terrissol M. Computational modelling of low-energy electron-induced DNA damage by early physical and chemical events. *Int J Radiat Biol* 1997;**71**:467–83.
83. Prise KM, Folkard M, Newman HC, Michael BD. Effect of radiation quality on lesion complexity in cellular DNA. *Int J Radiat Biol* 1994;**66**:537–42.
84. Hill MA. Fishing for radiation quality: chromosome aberrations and the role of radiation track structure. *Radiat Prot Dosimetry* 2015;**166**:295–301.
85. Sutherland BM, Bennett PV, Sutherland JC, Laval J. Clustered DNA damages induced by x rays in human cells. *Radiat Res* 2002;**157**:611–6.
86. Lomax ME, Gulston MK, O'Neill P. Chemical aspects of clustered DNA damage induction by ionising radiation. *Radiat Prot Dosimetry* 2002;**99**:63–8.
87. Box HC, Dawidzik JB, Budzinski EE. Free radical-induced double lesions in DNA. *Free Radic Biol Med* 2001;**31**:856–68.
88. Douki T, Riviere J, Cadet J. DNA tandem lesions containing 8-oxo-7,8-dihydroguanine and formamido residues arise from intramolecular addition of thymine peroxyl radical to guanine. *Chem Res Toxicol* 2002;**15**:445–54.
89. Eccles LJ, O'Neill P, Lomax ME. Delayed repair of radiation induced clustered DNA damage: friend or foe? *Mutat Res* 2011;**711**:134–41.
90. Magnander K, Hultborn R, Claesson K, Elmroth K. Clustered DNA damage in irradiated human diploid fibroblasts: influence of chromatin organization. *Radiat Res* 2010;**173**:272–82.
91. Tsao D, Kalogerinis P, Tabrizi I, Dingfelder M, Stewart RD, Georgakilas AG. Induction and processing of oxidative clustered DNA lesions in 56Fe-ion-irradiated human monocytes. *Radiat Res* 2007;**168**:87–97.
92. Malyarchuk S, Castore R, Harrison L. Apex1 can cleave complex clustered DNA lesions in cells. *DNA Repair* 2009;**8**:1343–54.
93. Sage E, Harrison L. Clustered DNA lesion repair in eukaryotes: relevance to mutagenesis and cell survival. *Mutat Res* 2011;**711**:123–33.
94. Georgakilas AG, O'Neill P, Stewart RD. Induction and repair of clustered DNA lesions: what do we know so far? *Radiat Res* 2013;**180**:100–9.