

Chalcophile Elements and Sulfides in the Upper Mantle

Ekaterina S. Kiseeva¹, Raúl O. C. Fonseca², Duane J. Smythe¹



1811-5209/17/0013-0111\$2.50 DOI: 10.2113/gselements.13.2.111

Sulfides are among the most important petrogenetic agents in magmatic systems. They are ubiquitous in most upper-mantle rock types, common as inclusions in diamonds and they host significant amounts of geochemically and economically important chalcophile ('sulfur-loving') elements, such as Cu, Ni, Pb, In, Au and the platinum-group elements. Despite their low abundance (<< 1% of the bulk rock), residual sulfides have a disproportionate control over the chalcophile element budget in upper mantle lithologies, as well as that of melts derived from the Earth's mantle.

KEYWORDS: chalcophile, element partitioning, upper mantle, sulfides

INTRODUCTION

Sulfides are minerals that contain the sulfide anion, S^{2-} , as the major structural component. Sulfides are common accessory minerals in the Earth's interior. They are observed in upper-mantle peridotites and oceanic basalts, in mantle xenoliths, and are also the most common mineral inclusion in diamonds. Despite sulfur being a minor component in the mantle – present at concentrations of ~150–300 parts per million (ppm) S in the bulk silicate Earth (e.g. Palme and O'Neill 2013) – sulfides are, nonetheless, very important phases because many elements have a strong affinity for sulfur. These sulfur-loving elements are termed 'chalcophiles' (Fig. 1), and they are concentrated in sulfide minerals. Thus, sulfides control the behaviour of the chalcophile elements during magmatic processes.

Most chalcophile elements (some of which are also siderophile, or 'iron-loving') are economically important metals; ten of these are the most valuable metals in the world [In, Re, Au, Ag and the platinum-group elements (PGEs)]. In Earth's crust, these elements are typically found associated with base-metal sulfides, and sulfide deposits are mined extensively as a result. In addition to their economic importance, residual sulfides exert a strong control on the budget of chalcophile elements during partial melting and crustal differentiation and, hence, have a profound effect on the trace-element concentrations found in erupted magmas.

In this article, we explore the occurrence of sulfides in the Earth's mantle, the role of sulfides in the partitioning of chalcophile elements in magmatic systems, and the abundance of chalcophile elements in mantle-derived silicate melts.

SULFIDES IN THE UPPER MANTLE

Sulfides are ubiquitous accessory phases in all types of mantle xenoliths. The major sulfide phases present in mantle rocks are pentlandite, pyrrhotite and chalcopyrite (Vaughan and Corkhill 2017 this issue). Also present are the 'monosulfide solid solution' (mss) and 'intermediate solid solution' (iss) phases. The observed mineralogy of mantle

sulfides, however, likely represents low temperature (<300°C) re-equilibration of high-temperature mss or, possibly, sulfide melt. Therefore, most recent studies have concentrated on reporting the bulk chemistry of polyphase inclusions (e.g. Bulanova et al. 1996), and this will be the focus of this discussion.

The compositions of sulfides from mantle xenoliths and diamond inclusions are shown in FIGURE 2. Sulfides observed in mantle xenoliths are dominantly Fe-rich with minor amounts of Ni and Cu. Peridotite xenoliths generally contain sulfides with ~20 wt% Ni and 4–5 wt% Cu, on average. Eclogitic sulfides are generally similar with respect to their Cu content compared to peridotite-hosted sulfides, but are consistently lower in Ni, with average concentrations ~3 wt% Ni.

Even though sulfides are accessory phases in mantle xenoliths (<0.1%) they are the most common mineral inclusions found in diamonds, accounting for 35%–46% of all diamond inclusions in the Finsch, Koffiefontein and Premier kimberlite pipes of South Africa (Harris and Gurney 1979). The Ni and Cu concentrations in sulfide inclusions in both peridotite- and eclogite-type diamonds are broadly similar compared to intergranular sulfides in peridotite and eclogite xenoliths, respectively. The marked difference in Ni concentrations between peridotite- and eclogite-hosted sulfides has been used as a paragenetic discriminator (Bulanova et al. 1996). Pearson et al. (1998) also noted that Ni concentrations of diamond-hosted sulfide inclusions correlate broadly with Os, and that Re/Os may be a more robust petrogenetic indicator for differentiating between peridotitic and eclogitic sulfides. Of course, how the concentrations of these elements in sulfides may be used as petrogenetic indicators hinges on our understanding of their partitioning behaviour between sulfide and silicate melt during mantle melting.

¹ Department of Earth Sciences
University of Oxford
Oxford, UK, OX1 3AN
E-mail: kate.kiseeva@earth.ox.ac.uk
duane.smythe@earth.ox.ac.uk

² Steinmann Institut
Universität Bonn
53115 Bonn, Germany
E-mail: raul.fonseca@uni-bonn.de





FIGURE 2 Ternary diagram ($\text{CuS}_{0.5}$, FeS and NiS) showing the major element composition of the mantle sulfides (in weight %). Green (peridotite); red (eclogite and pyroxenite). Data from Efimova et al. (1983), Bulanova et al. (1996), Pearson et al. (1998), Guo et al. (1999), Richardson et al. (2001), Westerlund et al. (2006).

studies (blue symbols in FIG. 3A), where efforts were made to avoid micronugget contamination, agree with the prediction of Equation (2). This contrast is a good illustration of the strides that have been made in investigating the chalcophile nature of trace elements over the last 25 years.

One other complexity that arises from the experimental approach described above is that the partition coefficient for a chalcophile element is related to sulfur and oxygen fugacities, which are easily controlled in laboratory experiments but that can be difficult to measure or estimate for natural systems. To overcome this problem, Kiseeva and Wood (2013, 2015) proposed an alternative second approach.

Following the study by Haughton et al. (1974), who related f_{S_2} to FeS and f_{O_2} to FeO, Kiseeva and Wood (2013) derived an expression that substitutes the $f_{\text{O}_2}/f_{\text{S}_2}$ ratio from Equation (2) with the FeO content of the silicate liquid (provided that the silicate melt is in equilibrium with the FeS sulfide), considering the exchange reaction:



Again, if we use divalent Pb as an example, Equation (3) will become:



As for Equation (1), Equation (4) can be rearranged so that the partition coefficient $D_{\text{Pb}}^{\text{sulf/sil}}$ relates to the equilibrium constant and the activity of FeO in silicate melt (Kiseeva and Wood 2013). This leads to the very simple statement that when a magma is sulfide-saturated (where the activity of FeS will be approximately unity), $D_{\text{Pb}}^{\text{sulf/sil}}$ mainly depends on the FeO content of the silicate liquid:

$$\log D_{\text{Pb}}^{\text{sulf/sil}} \approx A + \frac{x}{2} \log [\text{FeO}] \quad (5)$$

where A is a constant related to the free energy of the Fe–Pb exchange between sulfide and silicate liquids; x is a constant related to the valence of the element; $[\text{FeO}]$ is the

FeO content of the silicate melt in mole fraction or weight %. In theory, $-x$ will be the valence of the chalcophile element being studied (FIG. 3B).

Relationships between partition coefficients and FeO content of the silicate melt, summarised by Equation (5), have been successfully tested on a large number (>15) of chalcophile and moderately chalcophile elements and have proved to be valid for a broad range of temperatures and sulfide and silicate melt compositions. Moreover, the method outlined in Equation (5) allows the partition coefficient of a given chalcophile element to be described in terms of the easily measurable FeO content of silicate magma, precluding the need to account for oxygen and sulfur fugacities.

The two experimental approaches aimed at determining the partitioning relationships between the sulfide and silicate liquids are essentially addressing the same equilibrium, and they provide similar results for chosen T – f_{O_2} conditions, most notably when applied to melting of Earth's mantle.

To illustrate the equivalence of the approaches, we applied the two methods to the same dataset using data from Gaetani and Grove (1997). These authors controlled f_{O_2} and f_{S_2} and they synthesised silicate glasses over a range of FeO concentrations. In FIGURE 4, $D_{\text{Ni}}^{\text{sulf/sil}}$ is plotted as a function of $\log(f_{\text{O}_2}/f_{\text{S}_2})$ and $\log[\text{FeO}]$ and shows an excellent correlation.




FIGURE 3 (A) Compilation of literature sulfide–melt/silicate–melt partition coefficients for Os, given as a function of oxygen fugacity (here given relative to the fayalite–magnetite–quartz redox equilibrium – ΔFMQ) and at different sulfur fugacities (dashed and solid blue lines). Data sources include studies with micronugget contamination that was not taken into account (grey cross symbols). Blue symbols represent experiments where micronugget contamination was avoided. Data from Fonseca et al. (2011) and Mungall and Brenan (2014). (B) Sulfide–melt/silicate–melt partition coefficients for Pb versus $\log[\text{FeO}]$ content (wt %) of the silicate glass. Modified from Kiseeva and Wood (2013).




FIGURE 4 Sulfide–melt/silicate–melt partition coefficients for Ni. **(A)** Values of $\log D_{\text{Ni}}^{\text{sulf/sil}}$ plotted as a function of $\frac{1}{2} \log(f\text{O}_2/f\text{S}_2)$. **(B)** Values of $\log D_{\text{Ni}}^{\text{sulf/sil}}$ plotted as a function of $\log[\text{FeO}]$. The $\log[\text{FeO}]$ values have been corrected for the activity of FeS (see Kiseeva and Wood 2013). It is notable that the model using the relationship shown in Equation 5 (panel B) provides the slope (–1) predicted by the reaction stoichiometry. In contrast, the model using the relationship in Equation 2 (panel A), yields a substantially non-stoichiometric dependency (–0.63 slope). This could potentially be due to the uncertainties in the $f\text{S}_2$ measurements or due to the deviation from ideality in the Fe–Ni–Cu–S system. Additional discussion can be found in Gaetani and Grove (1997). Data from Gaetani and Grove (1997).

The method described in Equation (5) enables us to compile partition coefficients for a number of chalcophile elements in typical magmas (TABLE 1).

CHALCOPHILE ELEMENT BEHAVIOUR DURING MANTLE MELTING

The behaviour of chalcophile elements during mantle melting is summarised in FIGURE 5. Elements with a very strong chalcophile affinity ($D_M^{\text{sulf/sil}}$ of 10^4 – 10^6) (TABLE 1), such as Pt, Ir and Au, are extremely depleted in the silicate liquids due to their retention in the residual mantle sulfide. Only when sulfide is almost exhausted from the mantle source (~16% of aggregate fractional melting, assuming 140 ppm S in the mantle source), will highly chalcophile elements partition into the silicate melt. The concentration of chalcophiles that are incompatible in silicate phases, such as Pt and Au, decrease with continued melting as they become diluted by the silicate melt, while Ir, which is likely retained by olivine (Brenan et al. 2016), remains at constant concentrations.

The moderately chalcophile elements, such as Ni, Cu, In, and Pb have $D_M^{\text{sulf/sil}}$ values of ~10–1,000. Both In and Pb are incompatible elements in mantle silicates, so their concentrations continually decrease with increased melting. Copper, which is only moderately incompatible,

is relatively constant during fractional melting and shows only a small increase in concentration when approaching sulfide exhaustion. Despite being strongly chalcophile, Ni is also compatible in olivine ($D \sim 2$ –6), and this dominates its partitioning. Thus, sulfide exerts little or no control on Ni concentration during the melting of peridotite.

THE ROLE OF SULFIDES IN CONTROLLING THE RE–OS ISOTOPE SYSTEM

Rhenium and osmium were scavenged into the Earth's core during terrestrial accretion, leaving the silicate mantle highly depleted in these elements. Although they share this particular geochemical property, Re and Os have contrasting geochemical behaviours during partial melting of the mantle: Re is thought to be mildly incompatible, whereas Os is thought to be compatible (e.g. Fonseca et al. 2011). This fundamental difference is unique among commonly used isotope chronometers (e.g. K–Ar, Rb–Sr, Sm–Nd, Lu–Hf, and U–Th–Pb–He systems) where both parents and daughters are highly incompatible elements. The fact that Re is mildly incompatible and Os is compatible causes large variations of Re/Os ratios in oceanic basalts. Mid-ocean ridge basalts, for example, have Re/Os ratios




FIGURE 5 Instantaneous concentration of chalcophile elements **(A)** Ni, Cu, In, and Ir. **(B)** Pt, Au, and Pb in basaltic melt during fractional melting of a depleted mantle source, following the method of Mungall and Brenan (2014). The elements are grouped as shown for clarity. Major element chemistry of the melts and phase proportions were modelled using pMELTS (Ghiorso et al. 2002); sulfur solubility was calculated using the model of Smythe et al. (2017). Assuming 140 ppm sulfur in the mantle, sulfide becomes exhausted at approximately 16% melting (denoted as “sulfide out”), which allows highly chalcophile elements to partition into the silicate melt. As the chalcophile base metals have smaller sulfide melt/silicate melt partition coefficients relative to the PGEs (Kiseeva and Wood 2013; Mungall and Brenan 2014) they are less sensitive to the presence of sulfide in the residue. Partition coefficients and depleted mantle concentrations are shown in TABLE 2.

TABLE 1 EXPERIMENTALLY DERIVED PARTITION COEFFICIENTS OF SELECTED CHALCOPHILE ELEMENTS BETWEEN SULFIDE AND SILICATE MELTS

	Andesite	Basalt	Komatiite
	6–9 wt % FeO	8–11 wt % FeO	10–14 wt % FeO
Tl	14–18	13–15	11–13
In	14–24	11–17	8–12
Sb	22–39	17–26	13–20
As	29–36	25–31	22–27
Pb	41–62	33–46	25–37
Co	56–86	45–63	34–50
Cd	65–98	53–73	42–59
Cu	420–560	360–460	290–390
Ag	530–700	460–580	370–490
Se	550–850	450–650	350–500
Ni	770–1,160	630–870	500–690
Bi	1000–1250	900–1,100	800–950
Te	3,000–3,800	2,600–3,200	2,300–2,800
Re	22–22,377 (3,100)		
Au	4,100–11,200 (6,300)		
Pd	$6.70 \times 10^4 - 5.36 \times 10^5 (1.89 \times 10^5)$		
Rh	$5.72 \times 10^4 - 5.91 \times 10^5 (2.05 \times 10^5)$		
Ru	$3.03 \times 10^5 - 4.85 \times 10^5 (4.19 \times 10^5)$		
Ir	$4.80 \times 10^4 - 1.90 \times 10^6 (4.58 \times 10^5)$		
Os	$3.52 \times 10^5 - 1.15 \times 10^6 (7.49 \times 10^5)$		
Pt	$4.38 \times 10^3 - 3.45 \times 10^6 (8.45 \times 10^5)$		

Notes: Partition coefficients for elements Tl through Te (1.5 GPa, 1,400°C, and in equilibrium with FeS) are from Kiseeva and Wood (2015) and Kiseeva (unpubl). Values are derived according to Equation (5), and partition coefficients decrease with increasing FeO content of the silicate liquid. To determine partition coefficients at different temperatures, the reader is referred to the electronic appendix of Kiseeva and Wood (2015). The highest and the lowest partition coefficients for elements Re through Pt are given, with the average value in brackets. Values for Re (0.1–1.5 GPa and 1,200–1,250°C) are from Brenan (2008). Values for Au–Pt (1 atm and 1,200°C) are from by Mungall and Brenan (2014).

ranging between 10 and 2,000 (Shirey and Walker 1998). This variability makes the Re–Os isotopic system a useful geochemical tool for understanding mantle evolution.

The experimental literature has provided some detailed insights into the geochemical behaviour of Re during partial melting of the mantle. Rhenium is the least chalcophile of all the highly siderophile elements (e.g. Mungall and Brenan 2014), and it has been shown to partition into other mantle phases such as clinopyroxene, orthopyroxene, garnet, and spinel (Mallmann and O'Neill 2007), particularly under more reducing conditions. At oxidising conditions, however, Re will become hexavalent and will behave as a highly incompatible element during partial melting of the Earth's mantle. Osmium, on the other hand, is strongly

compatible in sulfide (e.g. Mungall and Brenan 2014): about 97% of the Os budget in source peridotites is controlled by sulfide (Luguet et al. 2007).

Effectively then, residual sulfide is the primary phase in the Earth's mantle that confers on the Re–Os isotope system its unique ability to date episodes of mantle melting. Mantle sulfides efficiently retain the entirety of the Os budget of a peridotite during partial melting, while Re, owing to its high incompatibility, is efficiently removed from the mantle. At high degrees of partial melting (10%–20%), alloys are expected to form from a host sulfide (Luguet et al. 2007; Fonseca et al. 2011), and, because of this, Os will further fractionate from Re.

If melt-depleted peridotites can be presumed to have lost all their Re during partial melting, then their $^{187}\text{Os}/^{188}\text{Os}$ can potentially be used to derive model ages that ostensibly track melt extraction events (reviewed in Lorand and Luguet 2016). This approach can be applied at both the whole-rock and mineral (i.e. sulfide) scale to date ancient melting events in the Earth's mantle (e.g. Wainwright et al. 2015). However, the same thing that makes the Re/Os system so useful in dating ancient melt events is also responsible for its biggest caveat. For example, metasomatic sulfides, which can be introduced to a mantle peridotite through basaltic melt percolation and/or fluid infiltration, will typically introduce Re into a virtually Re-free peridotite. Because of this, these metasomatic sulfides will develop more radiogenic $^{187}\text{Os}/^{188}\text{Os}$ over time, which will mix into the whole-rock signature and obscure the actual age of melt depletion of that peridotite. This difficulty can be partly circumvented by carrying out studies at the mineral scale to identify different generations of mantle sulfide, each with their own $^{187}\text{Os}/^{188}\text{Os}$ (cf. Wainwright et al. 2015). Sulfides are, thus, unique in their ability to fractionate Re from Os in magmatic systems and to control the Os isotope systematics during melting of the Earth's mantle.

CONCLUDING THOUGHTS AND FUTURE DIRECTIONS

The role of sulfides in igneous processes is not limited to the examples listed above. For instance, the Cu/Ag ratios of erupted magmas are fully controlled by the distinctive partitioning of Cu and Ag into solid and liquid sulfides (Wang and Becker 2015). Furthermore, the precipitation of sulfide may be responsible for the decline in Cu concentration of arc lavas during the evolution of volcanic systems at convergent margins (Lee et al. 2012). Sulfide also plays a significant role in the mysterious behaviour of Pb (Hart and Gaetani 2006) during mantle melting, which includes the nearly constant Ce/Pb (~25) and Nd/Pb (~20) ratios of mid-ocean ridge basalts and of ocean island basalts and the 'lead paradox', whereby all major Pb reservoirs appear to be substantially younger than the silicate Earth.

TABLE 2 PARTITION COEFFICIENTS USED FOR MODELLING THE CURVES IN FIGURE 5.

	Olivine	Clinopyroxene	Orthopyroxene	Spinel	Sulfide	Depleted mantle
Ni	5	1	1	0.001	980–1,140	1,960 ppm
Pb	0.003	0.1	0.009	0.001	61–74	23.2 ppb
Cu	0.05	0.043	0.15	0.22	540–635	30 ppm
In	0.04	0.3	0.02	0.001	23–28	12.2 ppb
Pt	0.009	–	–	–	850,000	6.2 ppb
Au	0.01	–	–	–	7,000	1 ppb
Ir	2	–	–	–	520,000	2.9 ppb

The Pd/Cu ratio may be an indicator of sulfide saturation in magmatic systems (Park et al. 2013), a parameter of fundamental importance in volcanic degassing. Ultimately, the precipitation of sulfides that act as a reservoir for the chalcophile and highly siderophile elements, such as Cu, Ag, Pb and the PGEs, is one of the key triggers that eventually leads to the formation of porphyry ore deposits and are key for the formation of magmatic sulfide deposits.

Thus, given that a typical peridotite will have sulfide phases present at less than 600 ppm (or 220 ppm S), such phases could certainly be said to “punch well above their weight”.

ACKNOWLEDGMENTS

The authors thank Frances Jenner, Yuan Li and Jim Mungall for thorough reviews. We also thank Hugh O'Neill and Bernie Wood for comments and discussion. ESK acknowledges support of the NERC grant NE/L010828/1. ■

REFERENCES

- Brenan JM (2008) Re–Os fractionation by sulfide melt–silicate melt partitioning: a new spin. *Chemical Geology* 248: 140–165
- Brenan JM, Bennett NR, Zajacz Z (2016) Experimental results on fractionation of the highly siderophile elements (HSE) at variable pressures and temperatures during planetary and magmatic differentiation. *Reviews in Mineralogy and Geochemistry* 81: 1–87
- Bulanova GP, Griffin WL, Ryan CG, Shestakova OY, Barnes, S-J (1996) Trace elements in sulfide inclusions from Yakutian diamonds. *Contributions to Mineralogy and Petrology* 124: 111–125
- Efimova ES, Sobolev NV, Pospelova LN (1983) Sulphide inclusions in diamonds and peculiarities of their paragenesis (in Russian). *Zapiski Vsesoyuznogo Mineralogicheskogo Obshchestva* 92: 300–309
- Ertel W, Dingwell DB, Sylvester PJ (2008) Siderophile elements in silicate melts — A review of the mechanically assisted equilibration technique and the nanonugget issue. *Chemical Geology* 248: 119–139
- Fonseca ROC, Mallmann G, O'Neill HStC, Campbell IH, Laurenz V (2011) Solubility of Os and Ir in sulfide melt: implications for Re/Os fractionation during mantle melting. *Earth and Planetary Science Letters* 311: 339–350
- Gaetani GA, Grove TL (1997) Partitioning of moderately siderophile elements among olivine, silicate melt, and sulfide melt: constraints on core formation in the Earth and Mars. *Geochimica et Cosmochimica Acta* 61: 1829–1846
- Ghiorso MS, Hirschmann MM, Reiners PW, and Kress VC (III) (2002) The pMELTS: A revision of MELTS for improving calculation of phase relations and major element partitioning related to partial melting of the mantle to 3 GPa. *Geochemistry, Geophysics, Geosystems* 3: 1–35
- Guo J, Griffin WL, O'Reilly SY (1999) Geochemistry and origin of sulphide minerals in mantle xenoliths: Qilin, southeastern China. *Journal of Petrology* 40: 1125–1149
- Harris JW, Gurney JJ (1979) Inclusions in diamond. In: Field JR (ed) *The Properties of Diamond*. Academic Press, London pp 555–591
- Hart SR, Gaetani GA (2006) Mantle Pb paradoxes: the sulfide solution. *Contributions to Mineralogy and Petrology* 152: 295–308
- Houghton DR, Roeder PL, Skinner BJ (1974) Solubility of sulfur in mafic magmas. *Economic Geology* 69: 451–467
- Jugo PJ, Wilke M, Botcharnikov RE (2010) Sulfur K-edge XANES analysis of natural and synthetic basaltic glasses: implications for S speciation and S content as function of oxygen fugacity. *Geochimica et Cosmochimica Acta* 74: 5926–5938
- Kiseeva ES, Wood BJ (2013) A simple model for chalcophile element partitioning between sulphide and silicate liquids with geochemical applications. *Earth and Planetary Science Letters* 383: 68–81
- Kiseeva ES, Wood BJ (2015) The effects of composition and temperature on chalcophile and lithophile element partitioning into magmatic sulphides. *Earth and Planetary Science Letters* 424: 280–294
- Lee C-TA and 8 coauthors (2012) Copper systematics in arc magmas and implications for crust–mantle differentiation. *Science* 336: 64–68
- Li Y, Audétat A (2012) Partitioning of V, Mn, Co, Ni, Cu, Zn, As, Mo, Ag, Sn, Sb, W, Au, Pb, and Bi between sulfide phases and hydrous basanite melt at upper mantle conditions. *Earth and Planetary Science Letters* 355–356: 327–340
- Lorand J-P, Luguet A (2016) Chalcophile and siderophile elements in mantle rocks: trace elements controlled by trace minerals. *Reviews in Mineralogy & Geochemistry* 81: 441–488
- Luguet A, Shirey SB, Lorand J-P, Horan MF, Carlson RW (2007) Residual platinum-group minerals from highly depleted harzburgites of the Lherz massif (France) and their role in HSE fractionation of the mantle. *Geochimica et Cosmochimica Acta* 71: 3082–3097
- Mallmann G, O'Neill HStC (2007) The effect of oxygen fugacity on the partitioning of Re between crystals and silicate melt during mantle melting. *Geochimica et Cosmochimica Acta* 71: 2837–2857
- Mungall JE, Brenan JM (2014) Partitioning of platinum-group elements and Au between sulfide liquid and basalt and the origins of mantle–crust fractionation of the chalcophile elements. *Geochimica et Cosmochimica Acta* 125: 265–289
- Palme H, O'Neill HStC (2013). Cosmochemical estimates of mantle composition. In: Carlson RW (ed) *Treatise on Geochemistry*, Volume 3 (2nd edition), Elsevier, Oxford, pp 1–39
- Pearson DG, Shirey SB, Harris JW, Carlson RW (1998) Sulphide inclusions in diamonds from the Koffiefontein kimberlite, S Africa: constraints on diamond ages and mantle Re–Os systematics. *Earth and Planetary Science Letters* 160: 311–326
- Richardson SH, Shirey SB, Harris JW, Carlson RW (2001) Archean subduction recorded by Re–Os isotopes in eclogitic sulfide inclusions in Kimberley diamonds. *Earth and Planetary Science Letters* 191: 257–266
- Shirey SB, Walker RJ (1998) The Re–Os isotope system in cosmochemistry and high-temperature geochemistry. *Annual Review of Earth and Planetary Sciences* 26: 423–500
- Smythe DJ, Wood BJ, Kiseeva ES (2017) The S content of silicate melts at sulfide saturation: new experiments and a model incorporating the effects of sulfide composition. *American Mineralogist* 102: 795–803
- Vaughan DJ, Corkhill C (2017) Mineralogy of sulphides. *Elements* 13: 83–89
- Wainwright AN, Luguet A, Fonseca ROC, Pearson DG (2015) Investigating metasomatic effects on the ¹⁸⁷Os isotopic signature: a case study on micrometric base metal sulphides in metasomatised peridotite from the Letlhakane kimberlite (Botswana). *Lithos* 232: 35–48
- Wang Z, Becker H (2015). Abundances of Ag and Cu in mantle peridotites and the implications for the behavior of chalcophile elements in the mantle. *Geochimica et Cosmochimica Acta* 160: 209–226
- Westerlund KJ and 5 coauthors (2006) A subduction wedge origin for Paleoproterozoic peridotitic diamonds and harzburgites from the Panda kimberlite, Slave craton: evidence from Re–Os isotope systematics. *Contributions to Mineralogy and Petrology* 152: 275–294 ■