

Title: Hyperpolarized ¹³C MRI: a novel approach for probing cerebral metabolism in health and neurological disease.

Authorship: James T. Grist PhD ^{1,2*}, Jack J. Miller PhD ^{3,4,5*}, Fulvio Zaccagna PhD ^{2*}, Mary A. McLean PhD ^{2,6}, Frank Riemer PhD ², Tomasz Matys PhD ², Damian J. Tyler PhD ^{3,5}, Christoffer Laustsen PhD ⁷, Alasdair J. Coles BM BCh PhD ⁸, Ferdia A. Gallagher PhD FRCR FRCP ².

Author affiliations

1 Institute of Cancer and Genomic Sciences, University of Birmingham, Birmingham.

2 Department of Radiology, University of Cambridge, Cambridge.

3 Department of Physiology, Anatomy, and Genetics, University of Oxford, UK.

4 Department of Physics, Clarendon Laboratory, University of Oxford.

5 Oxford Centre for Clinical Magnetic Resonance Research, John Radcliffe Hospital, Oxford, UK.

6 CRUK Cambridge Institute, Cambridge, United Kingdom.

7 MR Research Centre, Aarhus University, Aarhus.

8 Department of Clinical Neurosciences, University of Cambridge, Cambridge.

*These authors all contributed equally to this work

Corresponding author

Ferdia Gallagher, Department of Radiology, University of Cambridge.

Email: fag1000@cam.ac.uk

Search strategy and selection criteria

References included in this review article were chosen from PubMed using the terms ‘brain’ ‘hyperpolarization’, and ‘MRI’. There were no language restrictions. The final reference list was generated on the basis of relevance to the topics covered in this review.

Characters in title: 94

Running head: Imaging neurological metabolism

Number of words in abstract: 194

Number of words in introduction: N/A

Number of words in body of manuscript: 3290

Number of figures: 5

Number of color figures: 5

Number of tables: 0

Abstract

Cerebral metabolism is tightly regulated and fundamental for healthy neurological function. There is increasing evidence that alterations in this metabolism may be a precursor and early biomarker of later stage disease processes. Proton magnetic resonance spectroscopy (^1H -MRS) is a powerful tool to non-invasively assess tissue metabolites and has many applications for studying the normal and diseased brain. However, the technique has limitations including low spatial and temporal resolution, difficulties in discriminating overlapping peaks, and challenges in assessing metabolic flux rather than steady state concentrations. Hyperpolarized carbon-13 magnetic resonance imaging is an emerging clinical technique that may overcome some of these spatial and temporal limitations, providing novel insights into neurometabolism in both health and in pathological processes such as glioma, stroke and multiple sclerosis. This review will explore the growing body of pre-clinical data that demonstrates a potential role for the technique in assessing metabolism in the central nervous system. There are now a number of clinical studies being undertaken in this area and this review will present the emerging clinical data as well as the potential future applications of hyperpolarized ^{13}C magnetic resonance imaging in the brain, in both clinical and pre-clinical studies.

Key words

MRI, Hyperpolarization, Neurology, Neuro-oncology, Metabolism

Introduction: metabolic studies of the brain using proton and carbon-13 MR spectroscopy

Proton Magnetic Resonance Spectroscopy (^1H -MRS) is a non-invasive technique to probe *in vivo* metabolism that has been applied in a number of neurological diseases such as brain tumors, multiple sclerosis (MS), traumatic brain injury (TBI), stroke, and dementia¹⁻⁵. However, the millimolar concentration of most metabolites in human tissues results in a relatively low signal-to-noise ratio (SNR), therefore limiting the spatial and temporal resolution that can be achieved within a clinically acceptable scan time⁶. Metabolites that are commonly probed with ^1H -MRS include choline, creatine, N-acetyl aspartate, and lactate (markers of cellular proliferation, mitochondrial metabolism, neuronal integrity, and glycolytic metabolism, respectively). However other metabolites such as glutamate, glutamine, myo-inositol, and hypotaurine (which are involved in neurotransmission, amino acid metabolism and as osmolytes; concentrations of these molecules vary depending on cellular composition and can be used to help distinguish tumor subtypes⁷) are more challenging to quantify due to overlapping peaks on the ^1H -MRS spectrum at conventional clinical (1.5-3 T) field strengths⁶. Although the spatial distribution of these metabolites can be imaged with MR spectroscopic imaging techniques, this is usually at low spatial resolution (approximately 3-8 mL single voxel volume⁸) and is time consuming. The metabolite concentrations derived from ^1H -MRS usually reflect steady state tissue concentrations, which although important, do not reflect enzymatic flux or exchange which is complementary additional information.

Carbon-13-MRS (^{13}C -MRS) has also been utilized to non-invasively probe cerebral metabolic pathways such as the tricarboxylic acid (TCA) cycle, in both health and disease in a more detailed manner through the injection of ^{13}C -labelled endogenous tracers such as glucose, whereby incorporation and metabolism of the ^{13}C label can be followed dynamically, allowing metabolic pathways to be deconvolved in ways which complement the detection of steady-state metabolic pool sizes⁹⁻¹³. However, despite the ^{13}C enrichment of the injected molecule and the negligible ^{13}C signal from unlabelled background metabolites, the SNR is still very low, which is partly due to the less favorable magnetic properties of ^{13}C compared to ^1H (i.e. a lower gyromagnetic ratio and typically long nuclear T_1). This also restricts the achievable spatial and temporal resolutions of the technique and limits the technique to the research setting. Metabolic imaging of endogenous ^{13}C metabolites is currently challenging at clinical field strengths, however the increasing number of high field MR systems at 7 T or above offers the potential of using ^{13}C -MRS for thermal experiments in humans^{14,15}.

Dynamic Nuclear Polarisation (DNP) or hyperpolarized carbon-13 MRI

Dynamic Nuclear Polarisation (DNP) has been used to overcome the limitations of ^{13}C -MRS by providing a temporary increase in SNR of between 10,000 and 100,000-fold, allowing the uptake and metabolic conversion of physiologically relevant substrates to be imaged¹⁶⁻²⁰. This advance has opened up the potential of the technique to image tissue metabolism in real time at high resolution (2-4 s temporal resolution and with ~8 mL voxel volume²¹) and also offers the possibility of using it as a clinical tool to stratify patients based on their metabolic phenotype and detect changes in metabolism in response to treatment. Early clinical studies have focused on the use of hyperpolarized pyruvate to probe both glycolysis and TCA metabolism in oncology, renal, and cardiac metabolism²²⁻³¹. This review explores the application of ^{13}C -MRI to cerebral imaging and its future potential in the clinic.

Requirements for clinical hyperpolarized studies

Clinical hyperpolarized ^{13}C -MRI requires a dedicated sterile facility to fill pharmacy kits (a small sterile vial containing pyruvate, a syringe with water to dissolve the pyruvate and buffer to neutralise the solution as shown in Figure 1³²). The filled kit then undergoes hyperpolarization using a commercial clinical system (SPINlab, GE Healthcare, see Figure 2) and is quality checked for parameters such as pH, temperature and polarization level, before intravenous injection over 5-6 s (~ 5 mL/s) of ~ 250 mM hyperpolarized ^{13}C -pyruvate at 0.4 ml/kg body weight. ~ 250 mmol/L has been widely used for the initial clinical studies based on the concentration of neat pyruvic acid and the volume of heated water required to dissolve this rapidly and this concentration has shown a good toxicity profile. This produces an approximate intravascular concentration of ~ 1 -2 mmol/L after injection but a tissue concentration which is an order of magnitude smaller (~ 0.1 mmol/L) and similar to the physiological concentration of endogenous pyruvate³³. Lower doses may be possible in future but produce lower signal, because signal varies linearly with the amount of the injected tracer. Imaging the ^{13}C substrate and subsequent metabolites, is undertaken using dedicated MRI coils and sequences which acquire rapid images of the injected substrate and the metabolites that form from it in real-time^{21,34-40}. Imaging must be performed rapidly, as the increased polarization is transient and decays with the T_1 of the labelled ^{13}C (less than one minute in blood), and is therefore a limitation for *in vivo* measurements of slow enzymatic reactions.

Cellular localisation and quantification of cerebral pyruvate metabolism

Metabolism can be quantified by modelling the dynamics of signal acquired from the substrate (e.g. pyruvate) and its products (e.g. lactate or bicarbonate) to infer enzymatic activity⁴¹. Pyruvate resides at the junction of several important biochemical pathways such as the exchange to lactate in the cytosol, catalyzed by the enzyme lactate dehydrogenase (LDH), or the irreversible conversion to carbon dioxide, catalyzed by the mitochondrial enzyme pyruvate dehydrogenase (PDH). The labelled carbon dioxide subsequently exchanges with the bicarbonate pool, catalysed very rapidly by the enzyme carbonic anhydrase (CA), and this bicarbonate is more easily detectable due to its higher abundance at physiological pH^{42,43}. A confounder for the measurements of PDH flux is an alternative route of CO_2 fixation by pyruvate carboxylase (PC), which if significant, could lead to an overestimation of PDH enzymatic activity using hyperpolarized pyruvate⁴⁴. Therefore, imaging of hyperpolarized pyruvate metabolism can simultaneously probe both cytosolic and mitochondrial metabolism. Many pathological processes can result in alterations in cerebral metabolism. For example, necrosis⁴⁵, inflammatory cell activation⁴⁶, and breakdown of the blood-brain barrier⁴⁷ may result in elevated lactate in the CNS, either due to increased transport of pyruvate (as in the case of BBB disruption) or lactate formation (secondary to increased enzymatic exchange). Discussion of these effects in multiple sclerosis, stroke, traumatic brain injury, and brain tumors is found below.

Commonly, the term k_{PL} (the apparent exchange rate constant for pyruvate conversion to lactate) or the lactate-to-pyruvate ratio (Lac:Pyr or LP) are used as a measure of glycolytic activity in the cytosol⁴¹. In contrast, the term k_{PB} (the apparent rate constant of the irreversible conversion of pyruvate to bicarbonate) or the bicarbonate-to-pyruvate ratio (Bic:Pyr or BP) are used as a measure of mitochondrial TCA cycle activity.

Hyperpolarized ^{13}C -MRI of the healthy and diseased brain

There have been a number of pre-clinical studies assessing the metabolism of hyperpolarized ^{13}C substrates in the healthy and pathological brain, including traumatic brain injury, neuro-oncology, multiple sclerosis, and stroke. Detection of metabolic changes with hyperpolarized MRS is dependent on the magnitude of these alterations present in the pathological state to be assessed and small or transient changes may remain undetectable.

Metabolism in the healthy brain

The normal brain has a high basal metabolic rate and a reliance upon glucose oxidation⁴⁸. Initial hyperpolarized ^{13}C -MRI studies have investigated the metabolism of hyperpolarized $[1-^{13}\text{C}]$ pyruvate and $[2-^{13}\text{C}]$ pyruvate in healthy anaesthetised rodent and macaque brains⁴⁹⁻⁵². These studies have demonstrated the anaerobic metabolism of ^{13}C -pyruvate to ^{13}C -lactate in the brain parenchyma, catalyzed by cytosolic LDH, as well as the formation of ^{13}C -bicarbonate secondary to mitochondrial PDH and oxidative metabolism. The latter has been supported by the alternative approach of using $[2-^{13}\text{C}]$ pyruvate⁵³, which has shown the formation of TCA intermediates. Compared to other metabolically-active organs, the apparent rate of metabolism of hyperpolarized pyruvate in the brain is comparatively low ($\sim 0.003\text{ s}^{-1}$) compared to liver and kidney (~ 0.02 and 0.025 s^{-1} respectively)^{51,54}, which may reflect a limitation of pyruvate transport across the blood-brain barrier (BBB). Studies performed with hyperpolarized $[1-^{13}\text{C}]$ ethyl-pyruvate, a lipophilic analogue of pyruvate, have revealed enhanced diffusion-facilitated transport across the BBB and subsequent hydrolysis to pyruvate followed by exchange to lactate, demonstrating that BBB transport of pyruvate may be rate limiting in some species^{47,55,56}. There is evidence that the detection of $[1-^{13}\text{C}]$ pyruvate metabolism in the anaesthetised porcine brain may to some extent be limited by BBB transport⁴⁷; Figure 3. Furthermore, studies have also assessed the difference in pyruvate to lactate exchange before and after transient opening of the BBB with both mannitol and focused ultrasound, demonstrating elevated exchange after permeabilization^{47,57}.

Further pre-clinical healthy brain studies have demonstrated metabolism with other tracers such as $[1-^{13}\text{C}]$ lactate, to measure MCT1 and LDHA activity, $[1-^{13}\text{C}]$ glutamine, to measure IDH mutation status, and ketoisocaproic acid (KIC) to probe leucine metabolism^{56,58,59}. A further tracer of interest in brain studies could be $[1,4-^{13}\text{C}_2]$ fumarate, which may inform upon cellular necrosis through exchange to malate⁶⁰. Further probes that may be used in future clinical work have been reviewed elsewhere⁶¹.

Transport of ^{13}C pyruvate across the anaesthetised blood-brain barrier

Some degree of controversy exists surrounding the rate of transport and metabolism of pyruvate under differing anaesthetic regimes, with initial work suggesting that the quantities of hyperpolarized ^{13}C -bicarbonate and ^{13}C -lactate detected after the injection of hyperpolarized $[1-^{13}\text{C}]$ pyruvate are to some extent dependent upon the anaesthetic regimen used⁵⁸. In addition to alterations in transport, anaesthesia may also lead to metabolic shifts: in the heart, increasing the concentration of inhaled isoflurane has been shown to shift cardiac metabolism towards glycolysis, increasing the detected lactate-to-pyruvate ratio and decreasing the signal from bicarbonate⁶². As patients are usually imaged awake, this could influence comparisons between pre-clinical and human results: for

example, the low bicarbonate signal detected in the anaesthetised brain may be explained by a shift towards glycolysis driven by anaesthesia⁵⁸.

Imaging the healthy human brain with hyperpolarized ¹³C-MRI and comparison to ¹⁸F-FDG PET

Initial clinical studies of healthy brain metabolism have shown the rapid delivery and metabolism of pyruvate to bicarbonate, as well as exchange to lactate in the normal brain parenchyma^{63,64,65}. A recent study of the healthy brain quantified the metabolism of pyruvate in both white and grey matter, with results revealing significantly increased ¹³C-pyruvate, ¹³C-lactate, and ¹³C-bicarbonate in grey matter compared to white matter, which in part may be due to the higher perfusion in the former⁶³. However, this regional variation is also likely to relate to real alterations in tissue metabolism as there is increased ¹⁸F-FDG uptake on PET in grey matter compared to white matter, which is less dependent on speed of perfusion⁶⁶.

In comparison to hyperpolarized ¹³C-MRI measurements of pyruvate metabolism, ¹⁸F-FDG PET provides a highly sensitive measure of glucose metabolism, particularly the transport and phosphorylation of the FDG glucose analogue by GLUT-1 and hexokinase respectively. However, as PET measures the accumulation of radiolabel, it is unable to discriminate between the injected radiolabelled tracer and downstream metabolic products or their distribution in the cellular or extracellular spaces. Despite the significant increase in SNR afforded by hyperpolarization, it remains significantly less sensitive to the detection of metabolites than PET and in general this means a lower spatial resolution. However, the major strength of HP MRS over PET is the ability to discriminate an injected substrate from its metabolic product or products allowing a quantitative measure of enzymatic activity. Moreover, although the spatial resolution of HP MRS is insufficient to discriminate between intracellular and extracellular metabolites, as the formation of hyperpolarized lactate is intracellular, its presence indicates that intracellular metabolism has occurred even if that lactate has been exported to the extracellular space. Another important distinction between the two techniques is the timescale of metabolism that is being assessed as HP MRS detects pyruvate metabolism over seconds to minutes due to the short half-life, whereas PET assesses FDG accumulation over minutes to hours; therefore, the two methods are measuring metabolism on very different timescales. Therefore the two techniques are highly complementary.

Quantitative modelling has shown that there are regional variations in ¹³C-pyruvate metabolism across the brain with lower k_{PL} in deep white matter compared to the brainstem and basal ganglia. Interestingly, steady state lactate is not commonly detected in the resting healthy adult brain using ¹H-MRS and the difference between this finding and the significant ¹³C-lactate signal demonstrated with hyperpolarized ¹³C-MRI may be partly explained by the fact that the latter probes dynamic changes in lactate labelling following the injection of a supraphysiological bolus of pyruvate. Indeed, it may be that the static pool of lactate measured by ¹H-MRS is relatively small due to the rapid transport and metabolism of pyruvate in neurons. A further study focusing on the use of [2-¹³C]pyruvate in the healthy brain has been recently undertaken, with results showing the formation of TCA intermediates dynamically in the parenchyma, particularly glutamate⁶⁷. Example hyperpolarized ¹³C images from healthy volunteers are seen in Figures 4 and 5.

Further studies with larger cohorts of healthy volunteers at higher spatial resolutions may elucidate other regional differences in the metabolism of the healthy brain, allowing for a better understanding of normal cerebral physiology, as well as providing a baseline for comparison to pathological tissue.

Imaging the diseased brain with hyperpolarized ^{13}C -MRI

There are a number of neurological conditions which could benefit from the application of hyperpolarized ^{13}C -MRI to understand alterations in neurological metabolism due to cerebral insult and therapeutic response. This review will discuss neuro-oncology, multiple sclerosis (MS), traumatic brain injury (TBI) and stroke. Elevation in tissue lactate is a common feature in all of these conditions which raises the potential for the use of hyperpolarized ^{13}C -MRI as a tool to probe metabolism in many neurological conditions.

Brain tumors

Several early studies using hyperpolarized $[1-^{13}\text{C}]$ pyruvate as a probe for pre-clinical models of brain tumors have demonstrated increased lactate in implanted cerebral tumors, both orthotopic xenograft models in immunocompromised animals and implanted rodent-derived cell cultures in immunocompetent animals⁶⁸⁻⁷⁰. Lactate labelling has been shown to correlate with the histological aggressiveness of the tumor and a reduction in lactate labelling has been shown to correspond to therapy response, as chemotherapeutic agents such as temozolomide may have a metabolic effect before a reduction in tumor volume^{56,71}. Several studies assessing HP ^{13}C -MRS to detect therapeutic response in neuro-oncology have been undertaken, with reductions in the lactate signal observed following successful therapies, such as anti-VEGF⁷², temozolomide⁷³, dichloroacetate⁷⁴, and radiotherapy⁷⁵. Therefore, the use of hyperpolarized ^{13}C -MRI may be a valuable tool in the monitoring of neuro-oncological disease before and after chemotherapy and radiotherapy. The isocitrate dehydrogenase-1 (IDH-1) mutant status of glioma has been shown to correlate with the metabolism of hyperpolarized $[1-^{13}\text{C}]$ alpha-ketoglutarate and its subsequent conversion through the branched-chain amino acid transaminase BCAT1 to $[1-^{13}\text{C}]$ glutamate with the latter being significantly lower in mutant IDH1 cells and tumors⁷⁶. Such reactions form a specific readout of the tumor metabolome and are more challenging to assess with conventional *in vivo* non-invasive methods, such as ^1H proton spectroscopy⁷⁷. Therefore, this offers a promising target for non-invasively phenotyping tumors with hyperpolarized ^{13}C -MRI.

Initial human studies have assessed the feasibility of using hyperpolarized $[1-^{13}\text{C}]$ pyruvate to probe brain tumor metabolism^{21,78}. In keeping with the many pre-clinical studies in this field, this work has shown that there is significant conversion of pyruvate to lactate in a diverse range of tumors such as high grade glioma⁷⁵, oligodendroglioma⁷⁹ and melanoma metastases⁷⁹. However, there is a high degree of variation in lactate labelling between tumors, reflecting the complex heterogeneity of human tumors in comparison to animal models^{21,78}. This preliminary evidence provides some initial evidence for the potential role for the technique in studying brain tumors and future work will address the significance of the technique.

Multiple Sclerosis

Multiple Sclerosis (MS) is characterised by inflammation of the central nervous system, leading to inflammatory demyelination of neurons and oligodendrocyte death, and subsequently elevated metabolic demands on neurons

caused by a toxic elevation of extracellular ions such as calcium and sodium⁸⁰. Preclinically, hyperpolarized ¹³C-MRS has been used to characterise a cuprizone model of MS, which reproducibly develops inflammatory lesions that attract proinflammatory mononuclear phagocytes, particularly in the corpus callosum, similar to the human disorder⁸¹. An elevated lactate signal was demonstrated within MS lesions that were histologically confirmed to be rich in mononuclear phagocytes. Moreover, these activated proinflammatory mononuclear phagocytes had elevated pyruvate dehydrogenase kinase 1 (PDK1), responsible for inhibition of the oxidation of pyruvate by mitochondrial PDH, leading to increased metabolism to lactate in the cytosol. As well as furthering our understanding of lesional metabolism, hyperpolarized ¹³C-MRI could provide key insights into the viability of normal appearing tissue in the MS brain, through the assessment of lactate exchange and bicarbonate metabolism in normal appearing white and grey matter.

Traumatic brain injury

Acute TBI produces characteristic metabolic changes in the brain, both locally and remote to the site of injury, with elevated lactate detected with ¹H-MRS^{82,83}. As TBI is known to actively inhibit PDH activity and initiate mitochondrial dysfunction within four hours after injury⁸⁴, hyperpolarized pyruvate may potentially form a useful metric for understanding and quantifying the degree of injury sustained⁸⁵. TBI is a major cause of death and long-term disability, characterised by cognitive and memory impairment, mood disorders and neurodegenerative diseases.

Non-invasively detecting the spatial distribution of this metabolic dysregulation is challenging with conventional MRS and consequently invasive measures have been used to characterise this heterogeneity⁸⁶. Initial preclinical results with hyperpolarized [1-¹³C]pyruvate have shown regional alterations in lactate and bicarbonate formation after TBI, reporting a ~24% decrease in oxidative metabolism at the injury site and a concomitant increase in lactate production⁸⁵. Interestingly, there is some evidence that pyruvate or ethyl-pyruvate administration alone may be neuroprotective following TBI in the rat, albeit at higher doses than are typically used for hyperpolarized imaging experiments⁸⁷.

Stroke

Hyperpolarized ¹³C-MRI may have a significant role in understanding the development and response of the ischemic penumbra in stroke. A pre-clinical endothelin-1-induced ischaemic stroke model has been used to assess the metabolic alterations following acute ischaemic stroke in rats⁴⁹. The results showed an increase in total lactate production in the ischemic penumbra compared to the contralateral brain, which can partly be attributed to increased pyruvate supply and partly to LDH-mediated lactate formation. It is known that ischemia is a gradual process characterised by an intermediate stage in which the metabolic activity of the penumbral tissue is impaired but the damage is reversible and there is therefore the potential for complete functional recovery if blood flow is restored⁴. Currently, the ischemic penumbra is identified by the mismatch between perfusion and diffusion on MRI; however, as the penumbra has a high rate of glucose extraction characterised by anaerobic glycolysis and subsequent elevated lactate production this could be detected through the altered metabolism of hyperpolarized ¹³C-pyruvate⁸⁸. Indeed, invasive studies have shown that a measurement of metabolic activity in the penumbra may be more sensitive than conventional imaging approaches when estimating penumbral viability⁸⁹. Therefore,

since hyperpolarized ^{13}C -MRI can identify and quantify the dynamic production of lactate, it may be used to better visualise and characterise the penumbra, in a non-invasive fashion. However, studies focusing on longitudinal measurements of penumbral metabolism are required to understand the true potential of hyperpolarized ^{13}C -MRI to detect early response of the penumbra to treatment.

Concluding remarks

In conclusion, hyperpolarized ^{13}C -MRI provides a highly sensitive technique to directly and non-invasively probe the dynamics of tissue metabolism *in vivo* and offers the possibility to undertake novel *in vivo* biological explorations of neurological disease. The temporal and spatial resolution of conventional proton MR spectroscopy has been limited by low signal-to-noise, but the very significant increase in signal afforded by hyperpolarized ^{13}C -MRI has allowed the cerebral metabolism of ^{13}C -pyruvate to be probed in real-time. The technique has applications in a wide spectrum of diseases where metabolic status is impaired, reprogrammed or jeopardised. This method has revealed new insights into cerebral metabolism and offers a novel approach to study neurological disease, in particular the interplay between aerobic and anaerobic metabolism as seen in diseases as diverse as stroke, multiple sclerosis and brain tumors.

Author contributions:

Writing manuscript: JG, JJM, FZ, FAG

Reviewing Manuscript: JG, JJM, FZ, MAM, FR, TM, DJT, CL, AJC, FAG

Conflict of interest statement: The authors report no conflicts of interest.

Funding sources: The authors acknowledge grant support from the Medical Research Council, Little Princess Trust, Aarhus University, Evelyn Trust UK and research support from the British Heart Foundation, Cancer Research UK (CRUK), CRUK Cambridge Centre, MS Society, Prostate Cancer UK, National Institute of Health Research Cambridge Biomedical Research Centre, CRUK and the Engineering and Physical Sciences Research Council Imaging Centre in Cambridge and Manchester.

Ethics approval: N/A

References

- 1 Orphanidou-Vlachou E, Auer D, Brundler M a., Davies NP, Jaspan T, MacPherson L *et al.* ¹H magnetic resonance spectroscopy in the diagnosis of paediatric low grade brain tumours. *Eur J Radiol* 2013; **82**: e295–e301.
- 2 Tomiyasu M, Aida N, Shibasaki J, Tachibana Y, Endo M, Nozawa K *et al.* Normal lactate concentration range in the neonatal brain. *Magn Reson Imaging* 2016; **34**: 1269–1273.
- 3 Zhu X, Cao L, Hu X, Dong Y, Wang H, Liu F *et al.* Brain metabolism assessed via proton magnetic resonance spectroscopy in patients with amnesic or vascular mild cognitive impairment. *Clin Neurol Neurosurg* 2014.
- 4 Bivard A, Krishnamurthy V, Stanwell P, Yassi N, Spratt NJ, Nilsson M *et al.* Spectroscopy of reperfused tissue after stroke reveals heightened metabolism in patients with good clinical outcomes. *J Cereb Blood Flow Metab* 2014; **34**: 1944–1950.
- 5 Coles JP, Cunningham AS, Salvador R, Chatfield DA, Carpenter A, Pickard JD *et al.* Early metabolic characteristics of lesion and nonlesion tissue after head injury. *J Cereb Blood Flow Metab* 2009; **29**: 965–975.
- 6 Faghihi R, Zeinali-Rafsanjani B, Mosleh-Shirazi MA, Saeedi-Moghadam M, Lotfi M, Jalli R *et al.* Magnetic Resonance Spectroscopy and its Clinical Applications: A Review. *J Med Imaging Radiat Sci* 2017; **48**: 233–253.
- 7 Verma A, Kumar I, Verma N, Aggarwal P, Ojha R. Magnetic resonance spectroscopy - Revisiting the biochemical and molecular milieu of brain tumors. *BBA Clin* 2016; **5**: 170–178.
- 8 Wilson M, Andronesi O, Barker PB, Bartha R, Bizzi A, Bolan PJ *et al.* A methodological consensus on clinical proton MR spectroscopy of the brain: review and recommendations. 2019; : 1–39.
- 9 Sailasuta N, Robertson LW, Harris KC, Gropman AL, Allen PS, Ross BD. Clinical NOE ¹³C MRS for neuropsychiatric disorders of the frontal lobe. *J Magn Reson* 2008; **195**: 219–25.
- 10 Gruetter R, Novotny J, Boulware SD, Mason GF, Rothman DL, Shulman G *et al.* Localized ¹³C NMR Spectroscopy in the Human Brain of Amino Acid Labeling from D- [1- ¹³C] Glucose. 1994.
- 11 Wijnen JP, Van der Graaf M, Scheenen TWJ, Klomp DWJ, de Galan BE, Idema AJS *et al.* In vivo ¹³C magnetic resonance spectroscopy of a human brain tumor after application of ¹³C-1-enriched glucose. *Magn Reson Imaging* 2010; **28**: 690–697.
- 12 Sonnay S, Poirot J, Just N, Clerc AC, Gruetter R, Rainer G *et al.* Astrocytic and neuronal oxidative metabolism are coupled to the rate of glutamate–glutamine cycle in the tree shrew visual cortex. *Glia* 2018; **66**: 477–491.
- 13 Girault FM, Sonnay S, Gruetter R, Duarte JMN. Alterations of Brain Energy Metabolism in Type 2 Diabetic Goto-Kakizaki Rats Measured In Vivo by ¹³C Magnetic Resonance Spectroscopy. *Neurotox Res* 2019; **36**: 268–278.
- 14 Cheshkov S, Dimitrov IE, Jakkamsetti V, Good L, Kelly D, Rajasekaran K *et al.* Oxidation of [U-¹³C]glucose in the human brain at 7T under steady state conditions. *Magn Reson Med* 2017; **78**: 2065–2071.
- 15 De Feyter HM, Herzog RI, Steensma BR, Klomp DWJ, Brown PB, Mason GF *et al.* Selective proton-

- observed, carbon-edited (selPOCE) MRS method for measurement of glutamate and glutamine ^{13}C -labeling in the human frontal cortex. *Magn Reson Med* 2018; **80**: 11–20.
- 16 Ardenkjaer-Larsen JH, Fridlund B, Gram A, Hansson G, Hansson L, Lerche MH *et al.* Increase in signal-to-noise ratio of > 10,000 times in liquid-state NMR. *Proc Natl Acad Sci U S A* 2003; **100**: 10158–63.
- 17 Gallagher FA, Bohndiek SE, Kettunen MI, Lewis DY, Soloviev D, Brindle KM. Hyperpolarized ^{13}C MRI and PET: In Vivo Tumor Biochemistry. *J Nucl Med* 2011; **52**: 1333–1336.
- 18 Golman K, Olsson LE, Axelsson O, Månsson S, Karlsson M, Petersson JS. Molecular imaging using hyperpolarized ^{13}C . *Br J Radiol* 2003; **76 Spec No**: S118–S127.
- 19 Keshari KR, Wilson DM. *Chemistry and biochemistry of ^{13}C hyperpolarized magnetic resonance using dynamic nuclear polarization*. 2014.
- 20 Månsson S, Johansson E, Magnusson P, Chai C-M, Hansson G, Petersson JS *et al.* ^{13}C imaging—a new diagnostic platform. *Eur Radiol* 2006; **16**: 57–67.
- 21 Park I, Larson PEZ, Gordon JW, Carvajal L, Chen HY, Bok R *et al.* Development of methods and feasibility of using hyperpolarized carbon-13 imaging data for evaluating brain metabolism in patient studies. *Magn. Reson. Med.* 2018; **00**.
- 22 Nelson SJ, Kurhanewicz J, Vigneron DB, Larson PEZ, Harzstark AL, Ferrone M *et al.* Metabolic Imaging of Patients with Prostate Cancer Using Hyperpolarized [^{13}C]Pyruvate. *Sci Transl Med* 2013; **487**: 109–113.
- 23 Aggarwal R, Vigneron DB, Kurhanewicz J. Hyperpolarized 1- ^{13}C -Pyruvate Magnetic Resonance Imaging Detects an Early Metabolic Response to Androgen Ablation Therapy in Prostate Cancer. *Eur Urol* 2017; : 2–3.
- 24 Cunningham CH, Lau JYC, Chen AP, Geraghty BJ, Perks WJ, Roifman I *et al.* Hyperpolarized ^{13}C Metabolic MRI of the Human Heart: Initial Experience. *Circ Res* 2016; **119**: 1177–1182.
- 25 von Morze C, Bok RA, Sands JM, Kurhanewicz J, Vigneron DB. Monitoring urea transport in rat kidney in vivo using hyperpolarized ^{13}C magnetic resonance imaging. *Am J Physiol - Ren Physiol* 2012; **302**: 1658–1662.
- 26 Qi H, Mariager CØ, Nielsen PM, Schroeder M, Lindhardt J, Nørregaard R *et al.* Glucagon infusion alters the hyperpolarized ^{13}C -urea renal hemodynamic signature. *NMR Biomed* 2018; : 1–8.
- 27 Qi H, Nørtinger TS, Nielsen PM, Bertelsen LB, Mikkelsen E, Xu Y *et al.* Early diabetic kidney maintains the corticomedullary urea and sodium gradient. *Physiol Rep* 2016; **4**: 1–6.
- 28 Reed GD, Morze C Von, Verkman AS, Koelsch BL, Chaumeil MM, Lustig M *et al.* Imaging Renal Urea Handling in Rats at Millimeter Resolution Using Hyperpolarized Magnetic Resonance Relaxometry. *Tomography* 2016; **2**: 125–137.
- 29 Mariager CØ, Nielsen PM, Qi H, Ringgaard S, Laustsen C. Hyperpolarized ^{13}C , $^{15}\text{N}_2$ -urea T_2 relaxation changes in acute kidney injury. *Magn Reson Med* 2017; **702**: 696–702.
- 30 Nielsen PM, Eldirdiri A, Bertelsen LB, Jørgensen HS, Ardenkjaer-Larsen JH, Laustsen C. Fumarase activity: an in vivo and in vitro biomarker for acute kidney injury. *Sci Rep* 2017; **7**: 40812.
- 31 Grist JT, Mariager CØ, Qi H, Nielsen PM, Laustsen C. Detection of acute kidney injury with hyperpolarized [^{13}C , ^{15}N]Urea and multiexponential relaxation modeling. *Magn Reson Med* 2019; : 1–7.

- 32 Zaccagna F, Grist JT, Deen SS, Woitek R, Lechermann LMT, McLean MA *et al.* Hyperpolarized carbon-13 magnetic resonance spectroscopic imaging: A clinical tool for studying tumour metabolism. *Br J Radiol* 2018; **91**: 1–11.
- 33 Wang J, Wright AJ, Hesketh RL, Hu D, Brindle KM. A referenceless Nyquist ghost correction workflow for echo planar imaging of hyperpolarized [1-¹³C]pyruvate and [1-¹³C]lactate. *NMR Biomed* 2017; : e3866.
- 34 Le Page LM, Ball DR, Ball V, Dodd MS, Miller JJ, Heather LC *et al.* Simultaneous in vivo assessment of cardiac and hepatic metabolism in the diabetic rat using hyperpolarized MRS. *NMR Biomed* 2016; **29**: 1759–1767.
- 35 Gordon JW, Hansen RB, Shin PJ, Feng Y, Vigneron DB, Larson PEZ. 3D hyperpolarized C-13 EPI with calibrationless parallel imaging. *J Magn Reson* 2018; **289**: 92–99.
- 36 Lau AZ, Chen AP, Ghugre NR, Ramanan V, Lam WW, Connelly K a *et al.* Rapid multislice imaging of hyperpolarized ¹³C pyruvate and bicarbonate in the heart. *Magn Reson Med* 2010; **64**: 1323–31.
- 37 Vinding MS, Laustsen C, Maximov II, Søgaard LV, Ardenkjær-Larsen JH, Nielsen NC. Dynamic nuclear polarization and optimal control spatial-selective ¹³C MRI and MRS. *J Magn Reson* 2013; **227**: 57–61.
- 38 Larson PEZ, Kerr AB, Chen AP, Lustig MS, Zierhut ML, Hu S *et al.* Multiband excitation pulses for hyperpolarized ¹³C dynamic chemical-shift imaging. *J Magn Reson* 2008; **194**: 121–127.
- 39 Schmidt R, Laustsen C, Dumez J-N, Kettunen MI, Serrao EM, Marco-Rius I *et al.* In vivo single-shot ¹³C spectroscopic imaging of hyperpolarized metabolites by spatiotemporal encoding. *J Magn Reson* 2014; **240**: 8–15.
- 40 Durst M, Koellisch U, Frank A, Rancan G, Gringeri C V., Karas V *et al.* Comparison of acquisition schemes for hyperpolarised ¹³C imaging. *NMR Biomed* 2015; **28**: 715–725.
- 41 Daniels CJ, Mclean MA, Schulte RF, Robb FJ, Gill AB, Mcglashan N *et al.* A comparison of quantitative methods for clinical imaging with hyperpolarized ¹³C-pyruvate. *NMR Biomed* 2016; **29**.
- 42 Gallagher FA, Kettunen MI, Day SE, Hu D-E, Ardenkjaer-Larsen JH, Zandt R in 'T *et al.* Magnetic resonance imaging of pH in vivo using hyperpolarized ¹³C-labelled bicarbonate. *Nature* 2008; **453**: 940–3.
- 43 Gallagher FA, Sladen H, Kettunen MI, Serrao EM, Rodrigues TB, Wright A *et al.* Carbonic Anhydrase Activity Monitored in Vivo by Hyperpolarized¹³C-Magnetic Resonance Spectroscopy Demonstrates Its Importance for pH Regulation in Tumors. *Cancer Res* 2015; **75**: 4109–4118.
- 44 Siddiqui S, Kadlecsek S, Pourfathi M, Xin Y, Mannherz W, Hamedani H *et al.* The Use of Hyperpolarized Carbon-13 Magnetic Resonance for Molecular Imaging. *Adv Drug Deliv Rev* 2017; **113**: 3–23.
- 45 Bohndiek SE, Kettunen MI, Hu D, Witney TH, Kennedy BWC, Gallagher FA *et al.* Detection of tumor response to a vascular disrupting agent by hyperpolarized ¹³C magnetic resonance spectroscopy. *Mol Cancer Ther* 2010; **9**: 3278–3288.
- 46 Grist JT, Jarvis LB, Georgieva Z, Thompson S, Sandhu HK, Burling K *et al.* Extracellular lactate: A novel measure of t cell proliferation. *J Immunol* 2018; **200**.
- 47 Miller JJ, Grist JT, Serres S, Larkin JR, Lau AZ, Ray K *et al.* ¹³C Pyruvate Transport Across the Blood-

- Brain Barrier in Preclinical Hyperpolarised MRI. *Sci Rep* 2018; **8**: 15082.
- 48 Kaphingst KA, Persky S, Lachance C. Brain: Normal Variations and Benign Findings in FDG PET/CT imaging. *PET clin* 2010; **14**: 384–399.
- 49 Xu Y, Ringgaard S, Østergaard Mariager C, Bonde Bertelsen L, Schroeder M, Qi H *et al.* Hyperpolarized ¹³C Magnetic Resonance Imaging Can Detect Metabolic Changes Characteristic of Penumbra in Ischemic Stroke. *Tomography* 2017; **3**: 67–73.
- 50 Chen Y, Kim H, Bok R, Sukumar S, Mu X, Sheldon RA *et al.* Pyruvate to Lactate Metabolic Changes during Neurodevelopment Measured Dynamically Using Hyperpolarized ¹³C Imaging in Juvenile Murine Brain. *Dev Neurosci* 2016; **38**: 34–40.
- 51 Park I, Larson PEZ, Tropp JL, Carvajal L, Reed G, Bok R *et al.* Dynamic hyperpolarized carbon-13 MR metabolic imaging of nonhuman primate brain. *Magn Reson Med* 2014; **71**: 19–25.
- 52 Mayer D, Yen Y-F, Takahashi A, Josan S, Tropp J, Rutt BK *et al.* Dynamic and high-resolution metabolic imaging of hyperpolarized [1-¹³C]-pyruvate in the rat brain using a high-performance gradient insert. *Magn Reson Med* 2011; **65**: 1228–33.
- 53 Park JM, Josan S, Grafendorfer T, Yen YF, Hurd RE, Spielman DM *et al.* Measuring mitochondrial metabolism in rat brain in vivo using MR Spectroscopy of hyperpolarized [2-¹³C]pyruvate. *NMR Biomed* 2013; **26**: 1197–1203.
- 54 Bahrami N, Swisher CL, Von Morze C, Vigneron DB, Larson PEZ. Kinetic and perfusion modeling of hyperpolarized ¹³C pyruvate and urea in cancer with arbitrary RF flip angles. *Quant Imaging Med Surg* 2014; **4**: 24–32.
- 55 Hurd RE, Yen Y, Mayer D, Chen A, Wilson D, Kohler S *et al.* Metabolic imaging in the anesthetized rat brain using hyperpolarized [1-¹³C] pyruvate and [1-¹³C] ethyl pyruvate. *Magn Reson Med* 2011; **63**: 1137–1143.
- 56 Takado Y, Cheng T, Bastiaansen JAM, Yoshihara H, Lanz B, Mishkovsky M *et al.* Hyperpolarized ¹³C MRS reveals the rate-limiting role of the blood-brain barrier on the cerebral uptake and metabolism of L-lactate in vivo. *ACS Chem Neurosci* 2018; **9**: 2554–2562.
- 57 Mishkovsky M, Lengacher S, Comment A. Hyperpolarized ¹³C Magnetic Resonance Spectroscopy Reveals the Rate-Limiting Role of the Blood–Brain Barrier in the Cerebral Uptake and Metabolism of L-Lactate in Vivo. *ACS Chem Neurosci* 2018; **9**: 2554–2562.
- 58 Josan S, Hurd R, Billingsley K, Senadheera L, Park JM, Yen YF *et al.* Effects of isoflurane anesthesia on hyperpolarized ¹³C metabolic measurements in rat brain. *Magn Reson Med* 2013; **70**: 1117–1124.
- 59 Chaumeil MM, Larson PEZ, Woods SM, Cai L, Eriksson P, Robinson AE *et al.* Hyperpolarized [1-¹³C] glutamate: A metabolic imaging biomarker of IDH1 mutational status in glioma. *Cancer Res* 2014; **74**: 4247–4257.
- 60 Gallagher FA, Kettunen MI, Hu D-E, Jensen PR, Zandt R In 'T, Karlsson M *et al.* Production of hyperpolarized [1,4-¹³C₂]malate from [1,4-¹³C₂]fumarate is a marker of cell necrosis and treatment response in tumors. *Proc Natl Acad Sci U S A* 2009; **106**: 19801–19806.
- 61 Chaumeil MM, Najac C, Ronen SM. Studies of Metabolism Using ¹³C MRS of Hyperpolarized Probes. *Methods Enzymol* 2015; **561**: 1–71.
- 62 Steinhäuser J, Wespi P, Kwiatkowski G, Kozerke S. Assessing the influence of isoflurane anesthesia on

- cardiac metabolism using hyperpolarized [1-¹³C]pyruvate. *NMR Biomed* 2017; : e3856.
- 63 Grist JT, McLean MA, Riemer F, Schulte RF, Deen SS, Zaccagna F *et al.* Quantifying normal human brain metabolism using hyperpolarized [1-¹³C]pyruvate and magnetic resonance imaging. *Neuroimage* 2019; **189**: 171–179.
- 64 Tang S, Milshteyn E, Reed G, Gordon J, Bok R, Zhu X *et al.* A regional bolus tracking and real-time B1 calibration method for hyperpolarized ¹³C MRI. *Magn Reson Med* 2018; : 839–851.
- 65 Lee CY, Soliman H, Geraghty BJ, Chen AP, Connelly KA, Endre R *et al.* Lactate topography of the human brain using hyperpolarized ¹³C-MRI. *Neuroimage* 2019; **204**: 116202.
- 66 Arakawa S, Wright PM, Koga M, Phan TG, Reutens DC, Lim I *et al.* Ischemic thresholds for gray and white matter: A diffusion and perfusion magnetic resonance study. *Stroke* 2006; **37**: 1211–1216.
- 67 Chung BT, Chen HY, Gordon J, Mammoli D, Sriram R, Autry AW *et al.* First hyperpolarized [2-¹³C]pyruvate MR studies of human brain metabolism. *J Magn Reson* 2019; **309**: 106617.
- 68 Park I, Larson PEZ, Zierhut ML, Hu S, Bok R, Ozawa T *et al.* Hyperpolarized ¹³C magnetic resonance metabolic imaging: Application to brain tumors. *Neuro Oncol* 2010; **12**: 133–144.
- 69 Park I, Hu S, Bok R, Ozawa T, Ito M, Mukherjee J *et al.* Evaluation of heterogeneous metabolic profile in an orthotopic human glioblastoma xenograft model using compressed sensing hyperpolarized 3D ¹³C magnetic resonance spectroscopic imaging. *Magn Reson Med* 2013; **70**: 33–39.
- 70 Harrison C, Yang C, Jindal A, DeBerardinis RJ, Hooshyar MA, Merritt M *et al.* Comparison of kinetic models for analysis of pyruvate-to-lactate exchange by hyperpolarized ¹³C NMR. *NMR Biomed* 2012; **25**: 1286–1294.
- 71 Park I, Bok R, Ozawa T, Phillips JJ, James CD, Vigneron DB *et al.* Detection of early response to temozolomide treatment in brain tumors using hyperpolarized ¹³C MR metabolic imaging. *J Magn Reson Imaging* 2011; **33**: 1284–1290.
- 72 Ruterjng J, Ilmer M, Recio A, Coleman M, Vykoukal J, Alt E *et al.* Hyperpolarized ¹³C lactate-to-bicarbonate ratio as a biomarker for monitoring acute response of anti-VEGF treatment. *Nat Rev Drug Discov* 2016; **5**: 1–8.
- 73 Hyochol Ahn, PhD, Michael Weaver, PhD, Debra Lyon, PhD, Eunyoung Choi, RN, and Roger B. Fillingim P. Detection of Early Response to Temozolomide Treatment in Brain Tumors Using Hyperpolarized ¹³C MR Metabolic Imaging. *J Magn Reson Imaging* 2011; **33**: 1284–1290.
- 74 Park JM, Recht LD, Josan S, Merchant M, Jang T, Yen Y *et al.* Metabolic response of glioma to dichloroacetate measured in vivo by spectroscopic imaging. 2013; **15**: 433–441.
- 75 Day SE, Kettunen MI, Cherukuri MK, Mitchell JB, Lizak MJ, Morris HD *et al.* Detecting response of rat C6 glioma tumors to radiotherapy using hyperpolarized [1-¹³C]pyruvate and ¹³C magnetic resonance spectroscopic imaging. *Magn Reson Med* 2011; **65**: 557–563.
- 76 Izquierdo-Garcia JL, Cai LM, Chaumeil MM, Eriksson P, Robinson AE, Pieper RO *et al.* Glioma cells with the IDH1 mutation modulate metabolic fractional flux through pyruvate carboxylase. *PLoS One* 2014; **9**: 1–10.
- 77 Natsumeda M, Igarashi H, Motohashi K, Suzuki Y, Ohkubo M, Okamoto K. Advances and Challenges in Assessing 2-Hydroxyglutarate in Gliomas by Magnetic Resonance Spectroscopy : A Short Review. 2018; **8**: 1831–1838.

- 78 Miloushev VZ, Granlund KL, Boltyanskiy R, Lyashchenko SK, DeAngelis LM, Sosa E *et al.* Metabolic imaging of the human brain with hyperpolarized ^{13}C pyruvate demonstrates ^{13}C lactate production in brain tumor patients. *Cancer Res* 2018.
- 79 Miloushev VZ, Granlund KL, Boltyanskiy R, Lyashchenko SK, DeAngelis LM, Mellinghoff IK *et al.* Metabolic imaging of the human brain with hyperpolarized ^{13}C Pyruvate demonstrates ^{13}C lactate production in brain tumor patients. *Cancer Res* 2018; **78**: 3755–3760.
- 80 Mahad D, Lassmann H, Turnbull D. Review: Mitochondria and disease progression in multiple sclerosis. *Neuropathol Appl Neurobiol* 2008; **34**: 577–589.
- 81 Guglielmetti C, Najac C, Van Der Linden A, Ronen SM, Chaumeil MM. Hyperpolarized ^{13}C MR metabolic imaging can detect neuroinflammation in vivo in a preclinical model of Multiple Sclerosis. *PNAS* 2017; **18**: E6982–E6991.
- 82 Dhillon HS, Dose JM, Scheff SW, Prasad MR. Time course of changes in lactate and free fatty acids after experimental brain injury and relationship to morphologic damage. *Exp Neurol* 1997; **146**: 240–249.
- 83 Guglielmetti C, Chou A, Krukowski K, Najac C, Feng X, Riparip L-K *et al.* In vivo metabolic imaging of Traumatic Brain Injury. *Sci Rep* 2017; **7**: 17525.
- 84 Robertson CL, Saraswati M, Fiskum G. Mitochondrial dysfunction early after traumatic brain injury in immature rats. *J Neurochem* 2007; **101**: 1248–1257.
- 85 DeVience SJ, Lu X, Proctor J, Rangghran P, Melhem ER, Gullapalli R *et al.* Metabolic imaging of energy metabolism in traumatic brain injury using hyperpolarized $[1-^{13}\text{C}]$ pyruvate. *Sci Rep* 2017; **7**: 1907.
- 86 Carpenter KLH, Jalloh I, Gallagher CN, Grice P, Howe DJ, Mason A *et al.* ^{13}C -labelled microdialysis studies of cerebral metabolism in TBI patients. *Eur J Pharm Sci* 2014; **57**: 87–97.
- 87 Moro N, Sutton RL. Beneficial effects of sodium or ethyl pyruvate after traumatic brain injury in the rat. *Exp Neurol* 2010; **225**: 391–401.
- 88 Dani K a, Warach S. Metabolic Imaging of Ischemic Stroke: The Present and Future. *AJNR Am J Neuroradiol* 2014; **35**: S37–S43.
- 89 Pinczolit A, Zdunczyk A, Dengler NF, Hecht N, Kowoll CM, Dohmen C *et al.* Standard-sampling microdialysis and spreading depolarizations in patients with malignant hemispheric stroke. *J Cereb Blood Flow Metab* 2017; **37**: 1896–1905.

Figure Legends




Figure 1. A sterile fluid path for clinical studies.

The fluid path contains pyruvate (in sample vial), water for injection (dissolution syringe), and a neutralization medium (held in the receiver vessel). The filter on the receiver vessel removes electron paramagnetic agent (EPA) prior to quality control (QC). Figure adapted from ²¹.

A



B



Figure 2. The clinical 'SPINlab' hyperpolariser system exterior (A) and interior (B).

The Quality Control (QC) unit is shown on the right of each image, and the hyperpolarizer on the left. The system is sited next to a clinical scanner with a hatch in the wall for the delivery of the hyperpolarized sample; this will be placed into a syringe driver and injected into the patient.




Figure 3. Metabolic imaging of the porcine brain.

Imaging of the naïve porcine brain following injection of hyperpolarized ^{13}C -pyruvate. (A) ^{13}C -pyruvate signal is demonstrated in the vasculature. However, no ^{13}C -lactate signal (B) is seen in the brain parenchyma and therefore no significant metabolism is demonstrated on the kinetic rate constant map (k_{PL} in s^{-1} ; calculated only in the brain region). After the introduction of mannitol, both ^{13}C -pyruvate (D) and ^{13}C -lactate (E) are seen in the brain parenchyma as demonstrated on the calculated kinetic rate constant map (F). Figure adapted from reference ⁴⁷.




Figure 4. ^{13}C imaging demonstrating metabolite distribution in the healthy human brain.

Example summed images from the brain of a healthy volunteer demonstrating ^{13}C -pyruvate, ^{13}C -lactate, and ^{13}C -bicarbonate signal from three axial slices: superior, central and inferior. The T_1 -weighted images have also been shown, as have the quantitative maps of the exchange of pyruvate to lactate (k_{PL} in s^{-1})⁶³.




Figure 5. Resolution of ^{13}C imaging of the normal human brain.

IDEAL spiral Chemical Shift Images (CSI) are shown at the acquired spatial resolution to demonstrate the difference in SNR between: (B) pyruvate; (C) lactate; and (D) bicarbonate. ^{13}C images are normalized to the metabolite peak signal in the slice. Imaging parameters: field of view = 240 mm, matrix size = 40 x 40, slice thickness = 30 mm. Adapted from reference⁶³.