

Human Innate Lymphoid Cell Subsets Possess Tissue-Type Based Heterogeneity in Phenotype and Frequency

Yannick Simoni,^{*} Michael Fehlings, Henrik N. Kløverpris, Naomi McGovern, Si-Lin Koo, Chiew Yee Loh, Shawn Lim, Ayako Kurioka, Joannah R. Fergusson, Choong-Leong Tang, Ming Hian Kam, Koh Dennis, Tony Kiat Hon Lim, Alexander Chung Yaw Fui, Chan Weng Hoong, Jerry Kok Yen Chan, Maria Curotto de Lafaille, Sriram Narayanan, Sonia Baig, Muhammad Shabeer, Sue-Anne Ee Shiow Toh, Henry Kun Kiaang Tan, Rosslyn Anicete, Eng-Huat Tan, Angela Takano, Paul Klenerman, Alasdair Leslie, Daniel S.W. Tan, Iain Beehuat Tan, Florent Ginhoux, and Evan W. Newell^{*}

^{*}Correspondence: yannick_simoni@immunol.a-star.edu.sg (Y.S.), evan_newell@immunol.a-star.edu.sg (E.W.N.)

<https://doi.org/10.1016/j.immuni.2018.04.028>

(Immunity 46, 148–161, January 17, 2017)

In the originally published version of this article, the Supplemental Experimental Procedures were omitted from the Supplemental PDF. This has now been corrected online, and the journal apologizes for this error.

