

Discovery of the First Selective, Non-Peptidic Orexin 2 Receptor Agonists

Alexander Heifetz^{1}, Mike J. Bodkin¹ and Philip C. Biggin²*

¹Evotec (UK) Ltd., 114 Innovation Drive, Milton Park, Abingdon, Oxfordshire OX14 4RZ, United Kingdom;

²Department of Biochemistry, University of Oxford, South Parks Road Oxford OX1 3QU, United Kingdom;

*Corresponding author. Tel.: +44 (0)1235 83 89 25; Fax: +44 (0)1235 86 31 39

E-Mail address: Alexander.Heifetz@Evotec.com

ABSTRACT

In this issue, Nagase and colleagues report the discovery of the first selective non-peptidic Orexin 2 Receptor (OX₂R) agonists. The discovery of these OX₂R selective agonists opens up new avenues for therapies related to the activation of the orexin system, especially with respect to the treatment of sleep disorders such as narcolepsy.

VIEWPOINT

Through an extensive synthesis and screening program, Nagahara et al., 2015¹ report the discovery of the first selective non-peptidic Orexin 2 Receptor (OX₂R) agonists culminating in compound **26** (OX₂R EC₅₀ = 0.023 μM, E_{max} = 98%; OX₁R EC₅₀ = 1.616 μM, E_{max} = 100%). The design of small-molecule agonists (activators) rather than antagonists (inhibitors) of such receptors, and especially of peptide-activated G-protein coupled receptors (GPCRs), is considered as one of the current challenges in drug discovery. Such ligands hold high therapeutic potential especially for neuropeptide-activated GPCRs like the orexin receptors, where natural peptides are frequently non-selective and are often inefficient for *in-vivo* studies due to lack of ability to penetrate the blood-brain barrier (BBB). The selectivity profile of the agonists reported here is also attractive, and as such, they will serve as good tool compounds for exploration of OX₂R function that was not previously feasible with non-selective peptide agonists. Furthermore, given the recent appearance of a crystal structure of an antagonist (Suvorexant) in complex with OX₂R, it would seem likely that X-ray crystallography can now also be used to guide further drug-design efforts for this important target for several different conditions.

OX₁R and OX₂R, which are class A GPCRS, are located predominantly in the brain and are linked to a range of different physiological functions, including the control of feeding, energy metabolism, modulation of neuro-endocrine function, and the regulation of the sleep-wake cycle. The natural agonists for OX₁R and OX₂R are two non-selective neuropeptides, Orexin-A (OxA) and Orexin-B (OxB), which have dual activity, at both receptors. This last phenomenon has limited their use as probe compounds to dissect out precise contributions, but has enabled several conditions to be related to OXR activity. Years of research have suggested that OXR agonists could be useful for the treatment of sleep disorders, narcolepsy, cataplexy, obesity, hypophagia, as well as attention deficit hyperactivity, depression and related bipolar disorders. Furthermore, the discovery of OX₂R agonists will provide an excellent start point for the design of agonists for the related orexin-1 receptor (OX₁R). OX₁R has been shown to drive apoptosis in human colon cancer cells and treatment with orexins dramatically slowed the growth, and even reversed the development of established tumours. OX₁R agonists are therefore also prime candidates for colon cancer therapy. OXR agonists could also be useful for the treatment of Parkinson's disease, which is characterised by massive loss of hypocretin neurons. The design of selective OXR agonists has been a challenging problem, despite extensive mutagenesis² and modelling work³. This new chemical screening information along with the recently solved OX₂R crystal structure⁴ (PDB entry 4SOV) should create a step change in the development of drugs against this important family.

It is known that for effective antagonism, it is sufficient for small-molecule ligands to occupy a relevant receptor site in order to inhibit the binding of agonists. However for the discovery of the agonist, there is the additional complication and requirement that the ligand is able to activate the receptor. Peptide-activated GPCRs are considered especially challenging in this respect because of the potential for large number of specific and non-specific interactions that could be involved in binding and activation. The OXRs are particularly difficult because the natural agonist peptides, orexin A and orexin B, are quite

large at 33 and 28 amino acids respectively. Directly trying to mimic the effects of these large peptides with small molecules has been viewed as extremely challenging.

Although peptides or small molecule agonists activate the vast majority of GPCRs a spontaneous self-activation can also often be observed. Full agonists are traditionally defined as ligands that maximally activate the receptor, partial agonists induce submaximal activation, inverse agonists inhibit basal activity and neutral antagonists have no effect on basal activity, but competitively block access of other ligands.




Figure 1. Scheme describing the agonist promoted activation process from inactive ground state to fully activated G-protein bound state established crystallographically. The existence of such activation processes was demonstrated by GPCR crystallography for $\beta 1$ - and $\beta 2$ -Adrenergic and adenosine A_{2a} receptors and others.

Ligand binding is translated into conformational changes in the receptor that result in activation of intracellular G-proteins and/or β -arrestins, which in turn modulate the activity of downstream effectors inside the cell. GPCRs are in a dynamic structural equilibrium between multiple distinct conformations that can include more than one active or inactive conformation⁵ (see Figure 1). The mechanism and structural changes associated with the activation of

GPCRs remains a challenge. It was observed that agonists are able to break and mediate key interactions between trans-membrane helices (TMs) and by doing so enable them to move closer to each other, push them further apart, or rotate one relative to the other. These structural changes are known as “molecular switches”.^{5b} In recent years, it has become apparent that these molecular switches can induce an ensemble of activated conformations which trigger different activation pathways.

So, key a question that arises now is: Do we know how these new OX₂R agonists, and compound **26** in particular, work as agonists? The answer is not yet, but we are in a position to generate testable hypotheses. By combining these new agonists with the antagonist-bound crystal structure we can begin to postulate likely binding modes. Two such example poses are shown in Fig. 2. In the first pose (Fig. 2A) residues T111^{2.61}, Q134^{3.32}, T135^{3.33}, C210^{ECL2}, E212^{ECL2}, H224^{5.39}, F227^{5.42}, Y317^{6.48}, N324^{6.55}, K327^{6.58}, H350^{7.39} and two water molecules are involved in binding of compound **26** (superscript represents residue indices according to the Ballesteros and Weinstein⁶ numbering scheme). Furthermore, these modelling observations are directly supported by the published SDM data²⁻³. In the SDM studies the alanine mutations of T111^{2.61}, Q134^{3.32}, D211^{45.51}, W214^{5.54}, Y223^{5.38}, F227^{5.42}, F346^{7.35} and H350^{7.39} caused a large (> 50) fold decrease in the potency of OxA without affecting the efficacy compared to WT.




Figure 2. Two binding modes of compound **26** with OX₂R produced by hierarchical GPCR modelling protocol (HGMP)^{3a} that included post-docking OX₂R-**26** complex optimisation. **(A)** Literature-like "L" shape docking pose as reported by Nagahara et al., 2015¹. The carbon atoms of compound **26** are shown in cyan and for the receptor in blue. Nitrogen atoms are shown in blue, oxygen in red, sulfur in yellow and chlorine in light green. The key interactions are shown as yellow dashed line. **(B)** Superposition of OX₂R-Suvorexant crystal structure (PDB entry 4S0V) (backbone and Suvorexant coloured green) with the OX₂R-**26** complex (backbone coloured blue). The key differences in the TM conformations are shown as red arrows. **(C)** Alternative "U" shape docking pose. The carbon atoms of the **26** are shown in light pink and for the receptor are orange. **(D)** Superposition of OX₂R-Suvorexant crystal structure (PDB entry 4S0V) (backbone and Suvorexant coloured green) with the OX₂R-**26** complex (backbone coloured orange) in "U" docking pose. The key differences in the TM conformations are shown as red arrows.

The mutations Y232A^{5.47} and Y317A^{6.48} resulted in a reduction of both EC₅₀ (by 28.4 and 17.7 fold, respectively) and E_{max} of 44.9% and 49.6%, respectively of OxA. These mutations caused a moderate decrease in potency of OxA (by 22.3 fold) without affecting its efficacy. These SDM data suggest that there is no clear correlation between the importance of residues for potency and for efficacy. However, Y232^{5.47} and Y317^{6.48} are involved in OX₂R activation. The direct interaction of **26** with these key residues might explain to some extent its agonist activity. Recently solved crystal structures propose that the movement of TM5 and TM6 is made possible through rearrangement of the TM3-5-6 interface and it is potentially the most commonly conserved switch among Class A GPCRs.^{5b} This agonist-bound switch was recently proposed to be part of a larger “transmission switch” that accounts for the relocation of conserved residues W^{6.48} (Y317^{6.48} in OX₂R) and F^{6.44} towards P^{5.50}.^{5b} On the other hand the role of F/Y^{5.47} (Y232^{5.47} in OX₂R) is not clear but it is frequently engaged in interaction with agonists.^{5b} This typical inward movement of TM5 and TM6 with respect to the inactive state is also observed in compound **26** binding to the OX₂R (see Figure 2B).

It is also possible to model compound **26** in a Suvorexant-like flipped ‘U’ shape (see Figure 2C). According to this pose the residues T111^{2.61} (Ser in OX₁R), T135^{3.33}, E212^{ECL2}, H224^{5.39}, F227^{5.42}, Y317^{6.48}, N324^{6.55}, H350^{7.39} and two water molecules are involved in **26** binding (see Figure 2D). The interactions with the toggle switch residue Y317^{6.48} and with the aromatic cluster residue F227^{5.42}, support the activation switch mechanism that allows compound **26** to have OX₂R agonism function. For this pose, outward movements of TM5 and inward movement of TM6 (with respect to the antagonist-bound conformation) were observed to occur as part of the docking protocol (see Figure 2D). A potential explanation for OX₁R selectivity arises from potential interactions with the non-conserved residues T111^{2.61} (S102^{2.61} in OX₁R) and T135^{3.33} (A135^{3.33} in OX₁R). Further structural and structure activity relationship (SAR) exploration will be required to explore these hypotheses.

The excellent work of Nagese and colleagues suggests that with further application of X-ray crystallography, mutagenesis and modelling approaches we can look forward to the future development of agonists against these important targets for the treatment of multiple different conditions.

ABBREVIATIONS

OXR, Orexin receptors 1 and 2; BBB, blood brain barrier; HGMP, hierarchical GPCR modelling protocol; TM, transmembrane helix; ECL extracellular loop

References

1. Nagahara, T.; Saitoh, T.; Kutsumura, N.; Irukayama-Tomobe, Y.; Ogawa, Y.; Kuroda, D.; Gouda, H.; Kumagai, H.; Fujii, H.; Yanagisawa, M.; Nagase, H., Design and Synthesis of Non-Peptide, Selective Orexin Receptor 2 Agonists. *Journal of medicinal chemistry* **2015**.
2. Malherbe, P.; Roche, O.; Marcuz, A.; Kratzeisen, C.; Wettstein, J. G.; Bissantz, C., Mapping the binding pocket of dual antagonist almorexant to human orexin 1 and orexin 2 receptors: comparison with the selective OX1 antagonist SB-674042 and the selective OX2 antagonist N-ethyl-2-[(6-methoxy-pyridin-3-yl)-(toluene-2-sulfonyl)-amino]-N-pyridin-3-ylmethyl-acetamide (EMPA). *Molecular pharmacology* **2010**, 78 (1), 81-93.
3. (a) Heifetz, A.; Barker, O.; Morris, G. B.; Law, R. J.; Slack, M.; Biggin, P. C., Toward an understanding of agonist binding to human Orexin-1 and Orexin-2 receptors with G-protein-coupled receptor modeling and site-directed mutagenesis. *Biochemistry* **2013**, 52 (46), 8246-60; (b) Karhu, L.; Turku, A.; Xhaard, H., Modeling of the OX1R-orexin-A complex suggests two alternative binding modes. *BMC structural biology* **2015**, 15, 9.
4. Yin, J.; Mobarec, J. C.; Kolb, P.; Rosenbaum, D. M., Crystal structure of the human OX2 orexin receptor bound to the insomnia drug suvorexant. *Nature* **2015**, 519 (7542), 247-50.
5. (a) Manglik, A.; Kobilka, B., The role of protein dynamics in GPCR function: insights from the beta2AR and rhodopsin. *Current opinion in cell biology* **2014**, 27, 136-43; (b) Trzaskowski, B.; Latek, D.; Yuan, S.; Ghoshdastider, U.; Debinski, A.; Filipek, S., Action of molecular switches in GPCRs-- theoretical and experimental studies. *Current medicinal chemistry* **2012**, 19 (8), 1090-109.
6. Ballesteros, J. A.; Weinstein, H., Integrated methods for construction three dimensional models and computational probing of structure-function relations in G protein-coupled receptors. *Methods. Neurosci.* **1995**, 25, 366-428.