

TITLE PAGE

Article title: IndOSS-Assam: Investigating the feasibility of introducing a simple maternal morbidity surveillance and research system in Assam, India

Authors

1. Manisha Nair
National Perinatal Epidemiology Unit (NPEU), Nuffield Department of Population Health, University of Oxford, Oxford, United Kingdom
Email: manisha.nair@npeu.ox.ac.uk
2. Manoj K Choudhury*
Srimanta Sankaradeva University of Health Sciences, Guwahati, Assam, India
E-mail: drmanojchoudhury@yahoo.com
3. Saswati S Choudhury*
F.A.A. Medical College, Barpeta, Assam, India
E-mail: saswatisc@rediffmail.com
4. Swapna D. Kakoty
F.A.A. Medical College, Barpeta, Assam, India
E-mail: swapna_kakoty@rediffmail.com
5. Umesh C Sarma
Srimanta Sankaradeva University of Health Sciences, Guwahati, Assam, India
Email: ssuhs_assam@yahoo.in
6. Premila Webster
Oxford School of Public Health, Oxford Deanery, Oxford, United Kingdom
Email: premila.webster@dph.ox.ac.uk
7. Marian Knight
NPEU, Nuffield Department of Population Health, University of Oxford, Oxford, United Kingdom
Email: marian.knight@npeu.ox.ac.uk

On behalf of the IndOSS-Assam steering committee

*Authors contributed equally

Corresponding author

Manisha Nair
National Perinatal Epidemiology Unit (NPEU), Nuffield Department of Population Health, University of Oxford, Richard Doll Building, Old Road Campus, Headington, Oxford, OX3 7LF, United Kingdom
Phone: 01865-617820; Email: manisha.nair@npeu.ox.ac.uk

Key words: Maternal mortality, maternal morbidity, surveillance, India

Word count: 3234

List of abbreviations

| | |
|--------------|---|
| CI | Confidence intervals |
| e-MDR | Electronic maternal death reporting system |
| FAAMCH | Fakhruddin Ali Ahmed Medical College and Hospital |
| GMCH | Guahati Medical College and Hospital |
| IndOSS-Assam | Indian Obstetric Surveillance System - Assam |
| MMR | Maternal Mortality Ratio |
| PPH | Postpartum haemorrhage |
| UK | United Kingdom |
| UKOSS | UK Obstetric Surveillance System |

ABSTRACT

Objective: To assess the feasibility of establishing a simple maternal morbidity surveillance system in Assam (Indian Obstetric Surveillance System-Assam (IndOSS-Assam)) to investigate the incidence and trends in severe maternal complications. This study presents the surveillance platform of IndOSS-Assam.

Design: Four tasks were undertaken: (i) setting up a steering committee; (ii) establishing priorities for the region; (iii) mapping of surveillance sites; (iv) piloting case-notification system in selected centres.

Setting: Two government tertiary hospitals in the state.

Study population: Pregnant women delivering in the hospitals between March and August 2015.

Main outcome measures: Incidence and case fatality rates with 95% confidence intervals.

Results: Local stakeholder ownership and a simple uncomplicated anonymous system for case notification were the key strengths of this project. Cases and deaths were reported for six conditions: eclampsia, postpartum haemorrhage, puerperal sepsis, septic abortion, uterine rupture and anaemic heart failure. Among 10,475 women delivering over six months, 402 had one of these conditions and 66 died (case fatality 16%). The incidence of eclampsia was 17 per 1000 deliveries (95%CI=14 to 19), postpartum haemorrhage 11 per 1000 deliveries (95%CI=10 to 13) and anaemic heart failure was 3 per 1000 deliveries (95%CI=2 to 5). For each of the other three conditions, puerperal sepsis, septic abortion and uterine rupture, the incidence rate was 2 per 1000 deliveries.

Conclusion: IndOSS-Assam was shown to be a feasible and simple system for ongoing surveillance of maternal morbidity which can be used to monitor the trends in the incidence of specific severe life-threatening conditions during pregnancy.

SUMMARY BOX

What is already known on this subject?

- In low and middle income countries there is a continued focus on increasing the proportion of facility based deliveries, thereby increasing the need to monitor the quality of maternity care in hospitals.
- Case reviews of severe life-threatening maternal conditions have been initiated in a few countries such as South Africa, Benin, Ghana, Ivory Coast, Malaysia and Papua New Guinea, but these studies do not describe use of a hospital-based surveillance system in low and middle income countries to measure incidence and case-fatality of severe maternal morbidity, on an ongoing basis.

What are the new findings?

- This study showed the feasibility of establishing a simple anonymous hospital-based obstetric surveillance system (the Indian Obstetric Surveillance System-Assam (IndOSS-Assam)) as a first step towards estimating the incidence and case-fatality of severe life-threatening complications during pregnancy and childbirth in the Indian state of Assam.
- The surveillance platform of IndOSS-Assam is already established in the two medical colleges that participated in this pilot study. Involvement of local clinicians, policy makers, researchers, public health specialists, and academics from the inception of the project to testing the system showed that there is willingness and commitment to establish and scale up IndOSS-Assam to other hospitals in the state.

What are the recommendations for policy and practice?

- This pilot study showed a high incidence of eclampsia, postpartum haemorrhage, puerperal sepsis, septic abortion, uterine rupture and heart failure due to severe anaemia in pregnancy in the state along with a high case fatality rate. This suggests the need to focus research and policies to improve management of these conditions to reduce the burden of maternal mortality and morbidity in the state.

INTRODUCTION

India has the highest number of maternal deaths worldwide (50,000 maternal deaths in 2013),¹ and within India, the north-eastern state of Assam has a high maternal mortality ratio (MMR = 301/100,000 live births in 2012-13).² higher than the national average (MMR=178/100,000 live births in 2010-12).³ It has been hypothesised from the existing demographic and health surveys that anaemia and teenage pregnancy are important contributors to this high burden of maternal deaths in Assam, but there are limited data to investigate this.^{4,5} There are no up-to-date epidemiological data to refine existing programmes and policies, or develop new ones. Routine surveillance systems would provide the much needed data to investigate the risk factors for maternal morbidity and mortality in the state, as well as to assess the impact of ongoing maternal health programmes such as the 'Mamoni Scheme' and Janai Suraksha Yojana. These cash incentive schemes encourage pregnant women to undergo at least three antenatal check-ups and to deliver in a health institution.^{6,7} Currently, the electronic maternal death reporting system (e-MDR) of the national Health Mission in the state collates information on the number of maternal deaths per district per month.⁸ However, deaths are only the tip of the iceberg and a large number of women develop severe morbidity due to these conditions.

Establishing a state-wide hospital-based obstetric surveillance and research system is a first step towards generating evidence about the major risk and prognostic factors for maternal morbidity and mortality in the state of Assam. Audits are not generally acceptable due to concerns about attribution of blame,⁹ and thus an anonymous data collection system provides an ideal route to obtaining the appropriate information to conduct much needed research and surveillance. The proportion of childbirths in a health institution in Assam has been reported to have increased from 58% in 2010-11¹⁰ to 66% in 2012-13 (rural 63% and urban 84%)¹¹, thereby raising the need for a meaningful and straightforward means of monitoring the quality of maternity care in the hospitals. Once established, a surveillance system could provide a platform to conduct epidemiological studies, to generate evidence to prevent severe maternal morbidity (defined as life-threatening complications during pregnancy and childbirth) and to guide management of severe obstetric complications to reduce mortality.

However, existing approaches, such as the WHO near-miss approach, which has 25 markers for severe maternal morbidity¹² are complex and difficult to implement in practice, and not all criteria are reported to be relevant to the Indian context.^{13, 14} The United Kingdom has successfully implemented a surveillance and research system, the UK Obstetric Surveillance System (UKOSS),¹⁵ which has enabled the collection of anonymous information on a range of uncommon but severe pregnancy complications nationally. Since its inception in 2005, UKOSS has successfully generated evidence to guide prevention and management of major obstetric complications, inform policies, service planning and address patient safety issues.¹⁶ This has encouraged Australia, New Zealand and several countries in Europe to establish similar systems.¹⁷ The aim of this project was to assess the feasibility of establishing a simple obstetric surveillance and research system in Assam (the Indian Obstetric Surveillance System - Assam (IndOSS-Assam)) using the UKOSS model in order to describe the changing epidemiology of risk factors, management and outcomes associated with maternal deaths and specific morbidities in the state. IndOSS-Assam has two components: surveillance and research. This paper presents the results of the pilot study that assessed the feasibility of setting up the surveillance platform of IndOSS-Assam.

METHODS

The objective of this pilot study was to adapt the UKOSS methodology to the context of Assam in order to develop a simple maternal morbidity surveillance system for the state (IndOSS-Assam). Details of the UKOSS methodology are described elsewhere.^{15, 16} Briefly, UKOSS is a surveillance and research platform that enables collection of information on specific pregnancy complications through case notification cards sent to nominated reporting doctors

and midwives in all consultant-led obstetric units in the UK every month with an approach of 'nil-reporting'.¹⁵ For every case reported, details are collected on a data collection form by the clinician responsible for managing the case. Rigorous follow-up of non-responders ensures completeness of data.¹⁵ A similar methodology was used for IndOSS-Assam.

Four main tasks were undertaken in the development and piloting of the surveillance platform of the IndOSS-Assam: (i) setting up a steering committee; (ii) establishing priorities for the region; (iii) mapping of reporting hospitals; and (iv) piloting the system of case notification cards in selected centres. The fifth task was to assess the feasibility of using routine medical records for conducting epidemiological studies through the IndOSS-Assam 'research platform' in order to describe the risk factors, management and outcomes associated with specific life-threatening complications during pregnancy and child birth in the state of Assam. The details are presented in a separate paper.¹⁸ The activities undertaken to examine the feasibility of establishing IndOSS-Assam are summarised in Figure-1.

The conditions to be included in surveillance during the pilot study were identified by stakeholder consultation with members of the steering committee and others, based on their main concerns about severe maternal mortalities specific to Assam and informed by an initial analysis of the causes of death reported through the electronic maternal death reporting system (e-MDR). Currently, the e-MDR of the National Health Mission in Assam¹⁹ collates information on the number of maternal deaths per district per month and also has information about the causes and place of death.

Two tertiary hospitals, Fakhruddin Ali Ahmed Medical College and Hospital (FAAMCH), Barpeta and Guahati Medical College and Hospital (GMCH), Guahati were purposively selected by the steering committee as centres for pilot testing the case notification system of IndOSS-Assam, to include one hospital in the state capital and one more peripheral hospital. A case notification card (Figure-2) was designed to collect information on the number of cases and deaths due to the six selected priority conditions associated with increased risk of maternal mortality in the state. The IndOSS-Assam monthly case notification system was launched in March 2015 in the two medical colleges under the supervision of the steering committee.

Statistical analyses

We compiled the information from the case notification cards and calculated the proportion of cases as a percentage of the total deliveries with 95% confidence intervals (CI) and the case fatality rate with 95% CI for each condition in each hospital.

RESULTS

Setting up a steering committee

A multi-disciplinary steering committee was successfully constituted in the state including senior representatives from obstetrics, medical education, public health, medical research and university organisations, as well as individual representatives from professional specialties. A total of six steering committee meetings were held during a period of 18 months, either in person or by teleconference, including in consultation with UKOSS researchers in order to adapt the UKOSS methodology to the context of Assam.

Establishing priorities for the region

There were 3,753 maternal deaths reported through the e-MDR between April 2011 and August 2014, which is approximately half of the estimated 6,333 deaths based on the MMR of 301/100,000 live births. The main causes of death were reported to be anaemia, eclampsia, sepsis, haemorrhage, obstructed labour, and septic abortion (Figure-3). For almost half of the cases, the cause was not known or reported as 'others'. Almost 64% of the deaths occurred

in hospitals (Figure-4), which further confirmed that a hospital-based surveillance system would be able to capture a majority of the cases of severe maternal morbidity and mortality.

Based on the analyses of the causes of death and stakeholder consultation, six conditions were included in the pilot phase (Phase-1) of IndOSS-Assam: five causes of direct maternal mortality – eclampsia, postpartum haemorrhage, puerperal sepsis, septic abortion and uterine rupture, and one suspected cause of indirect maternal death in the state – anaemic heart failure (heart failure during pregnancy and delivery as a consequence of severe anaemia²⁰).

Mapping of reporting hospitals

The major tertiary and community government and private hospitals were identified and manually mapped to assess their geographical distribution. All 14 major tertiary government hospitals were identified (Figure 5).

Piloting the surveillance system in two selected centres

The pilot system was successfully established, with 100% of monthly reports returned to the steering committee at the Srimanta Sankaradeva University of Health Sciences for the two medical colleges during the pilot phase between March and August 2015. Paper-based case notification cards were received in a timely manner, nil-reporting ensured completeness and there were no identified problems in reporting. The reporting was overseen by either a senior obstetrician or the Head of the Department of Obstetrics and Gynaecology in each medical college.

There were a total of 10,475 deliveries in the two medical colleges participating in IndOSS-Assam in the six months of the pilot study; 402 cases and 66 deaths as a result of one of the six conditions under surveillance were reported (Table-1). The incidence of eclampsia was 17 per 1000 deliveries (95% CI 14 to 19), postpartum haemorrhage 11 per 1000 deliveries (95% CI 10 to 13) and anaemic heart failure was 3 per 1000 deliveries (95% CI 2 to 5). For each of the other three conditions, puerperal sepsis, septic abortion and uterine rupture, the incidence rate was 2 per 1000 deliveries. Case fatality due to these conditions was high, 16% (95% CI 13 to 20%) of women who were diagnosed with these conditions died, the highest case fatality rate being for puerperal sepsis and anaemic heart failure (Table-1).

Table-1: Total cases notified in 6 months of pilot study through the IndOSS-Assam

| Conditions under surveillance | Cases | Deaths | % cases (95% CI) | % fatality (95% CI) | Total deliveries |
|-------------------------------|------------|-----------|-------------------------|---------------------------|------------------|
| FAAMCH, Barpeta | | | | | |
| Eclampsia | 92 | 5 | 2.9 (2.4 to 3.6) | 5.4 (1.8 to 12.2) | 3145 |
| PPH | 44 | 5 | 1.4 (1.0 to 1.9) | 11.4 (3.8 to 24.6) | |
| Puerperal Sepsis | 6 | 1 | 0.2 (0.1 to 0.4) | 16.7 (0.4 to 64.1) | |
| Septic abortion | 10 | 1 | 0.3 (0.2 to 0.6) | 10.0 (0.4 to 45.0) | |
| Uterine rupture | 9 | 1 | 0.3 (0.1 to 0.5) | 11.1 (0.3 to 48.2) | |
| Anaemic heart failure | 17 | 5 | 0.5 (0.3 to 0.9) | 29.4 (10.3 to 55.9) | |
| Total | 178 | 18 | 5.7 (4.9 to 6.5) | 10.1 (6.1 to 15.5) | |
| GMCH, Guwahati | | | | | |
| Eclampsia | 82 | 16 | 1.1 (0.9 to 1.4) | 19.5 (11.6 to 29.7) | 7330 |
| PPH | 74 | 5 | 1.0 (0.8 to 1.3) | 6.8 (2.2 to 15.1) | |
| Puerperal Sepsis | 17 | 9 | 0.2 (0.1 to 0.4) | 52.9 (27.8 to 77.0) | |
| Septic abortion | 16 | 7 | 0.2 (0.1 to 0.4) | 43.8 (19.8 to 70.1) | |
| Uterine rupture | 17 | 2 | 0.2 (0.1 to 0.4) | 11.8 (1.5 to 36.4) | |

| | | | | | |
|-----------------------|--|----|------------------|---------------------|-------|
| Anaemic heart failure | 18 | 9 | 0.2 (0.1 to 0.4) | 50.0 (26 to 74) | |
| Total | 224 | 48 | 3.1 (3.0 to 3.5) | 21.4 (16.2 to 27.4) | |
| Both sites | Total in two tertiary hospitals in 6 months of pilot study | | | | |
| Eclampsia | 174 | 21 | 1.7 (1.4 to 1.9) | 12.1 (7.6 to 17.9) | 10475 |
| PPH | 118 | 10 | 1.1 (1.0 to 1.3) | 8.5 (4.1 to 15.0) | |
| Puerperal Sepsis | 23 | 10 | 0.2 (0.1 to 0.3) | 43.5 (23.2 to 65.5) | |
| Septic abortion | 26 | 8 | 0.2 (0.2 to 0.4) | 30.8 (14.3 to 51.8) | |
| Uterine rupture | 26 | 3 | 0.2 (0.2 to 0.4) | 11.5 (2.4 to 30.2) | |
| Anaemic heart failure | 35 | 14 | 0.3 (0.2 to 0.5) | 40.0 (23.9 to 57.9) | |
| Total | 402 | 66 | 3.8 (3.5 to 4.2) | 16.4 (12.9 to 20.4) | |

PPH=Postpartum haemorrhage

DISCUSSION

This study showed that setting up a simple surveillance system is feasible, and there is also willingness and enthusiasm among the clinicians in the state to establish IndOSS-Assam. Data collected during the pilot study on six conditions, eclampsia, postpartum haemorrhage, puerperal sepsis, septic abortion, uterine rupture and anaemic heart failure, showed a high incidence (4% of all deliveries) and fatality (16% of all cases) associated with these conditions among pregnant women who delivered in the two pilot hospitals.

A number of factors were felt to have contributed to the success of the pilot project, the most important being local stakeholders including clinicians, public health specialists, and academics from the medical colleges, hospitals and the university taking ownership and driving the project, with support from UKOSS researchers. The other key factor was the simple straight forward design of the IndOSS-Assam system. Regular meetings with the steering committee to identify hurdles in implementing the system and anonymous reporting of cases and deaths without attribution of blame were other strengths identified by the steering committee members and the reporting clinicians.

Strengths and limitations of this study

Despite the fact that Assam has a high maternal mortality ratio in India, little work has been undertaken to date to generate good quality evidence about the incidence of the severe life-threatening complications during pregnancy and childbirth, with the aim of monitoring the patterns and trends of these conditions in the state. The surveillance platform of the IndOSS-Assam can be used to generate evidence about trends in the incidence and case-fatality of important life-threatening conditions on an ongoing basis. This will help direct research and policies to improve the management of conditions with high incidence and case-fatality in order to reduce the burden of maternal mortality and morbidity in the state. This study demonstrated the feasibility of establishing IndOSS-Assam as a hospital-based surveillance system in the state. IndOSS-Assam being an anonymised data collection system avoids the attribution of blame while generating robust incidence estimates to improve maternal health in the state.

Although we were able to estimate hospital-based incidence rates for six major causes of maternal mortality and morbidity in Assam, these do not necessarily reflect the population-level incidence of these conditions. The estimated incidence and fatality could be lower than the true population incidence, as not all pregnant women deliver in a health facility. It is also possible that our estimates are higher than the true population rate, since pregnant women with serious complications are more likely to be referred to the tertiary hospitals. High fatality could be due to late referral of cases, but it could also be associated with inadequate management and care during pregnancy and delivery. Nevertheless, this pilot study showed that 1 in 25 women who delivered in two tertiary hospitals (FAAMCH and GMCH) suffered

from one of the six major causes of maternal deaths in the state and 1 in 130 women died as a result of these conditions. This further justifies the need for establishing IndOSS-Assam as a platform for ongoing surveillance and research in the state. Expanding the network of reporting hospitals for IndOSS-Assam to all tertiary government hospitals and major private maternity hospitals in subsequent phases of the project would ensure that information on the large majority of cases with severe complications are captured.

Findings in relation to other studies

An earlier study in the FAAMCH tertiary hospital identified 66 cases of severe maternal morbidity in 1,729 deliveries between June and October 2014 of which 39% were eclampsia and 24% postpartum haemorrhage²¹ leading to an estimated incidence rate of 1.5% for eclampsia and about 1% for postpartum haemorrhage among the total deliveries. These rates are a half to two thirds lower than that estimated by the IndOSS-Assam active surveillance in the same hospital between March and August 2015. The maternal death reviews in the state show eclampsia, postpartum haemorrhage and anaemia to be the major causes of maternal mortality (Figure-3), however, maternal mortality rates are not calculable as denominator information about the number of women delivering is not available. The added value of IndOSS is not in estimating the number of deaths due to these known causes of maternal death in Assam, but to monitor the incidence and case-fatality of these important conditions. Allied studies aim to improve their in-hospital management, as the number of facility-based deliveries increase in the state, in order to reduce maternal morbidity and mortality. IndOSS-Assam showed a high incidence of these conditions among pregnant women who delivered in the two pilot centres as well as high fatality rates associated with anaemic heart failure and sepsis.

The million death study estimated that a quarter of the maternal deaths in India were due to postpartum haemorrhage.²² 'Dead women talking', a civil society report on maternal deaths in India identified anaemia to be the second largest cause of maternal death in India after postpartum haemorrhage²³. Of the 22 women (out of 124) who were reported to die from anaemia, 11 had cardiac failure during labour or delivery, and in at least eight of the 29 deaths from postpartum haemorrhage, anaemia was a contributing factor²³. Another study in India reported a prevalence of 9.4% cardiac failure among women with severe anaemia.²⁴ This is the first system in Assam to initiate data collection to estimate the incidence of maternal morbidity on an ongoing basis, including cardiac failure due to anaemia during pregnancy. Prior to this study the incidence and case-fatality rates due to anaemic heart failure among pregnant women in Assam were not known. Through the research platform, we were also able to conduct a retrospective cohort study on the adverse maternal and fetal outcomes of anaemia during pregnancy. As mentioned above, the findings are presented separately.¹⁸

The incidence (2%) and case-fatality (12%) rates of eclampsia estimated by the IndOSS-Assam pilot study in the two medical colleges are higher than the rates reported by hospital-based studies in other states in India. A prospective study conducted in a hospital in Nagpur between January 2008 and December 2010 estimated the incidence of eclampsia to be 0.9% and case-fatality 5.5%.²⁵ Similar rates were reported by a teaching hospital in Karnataka (incidence 1% and case-fatality 4%).²⁶ A study in a tertiary hospital in Kolkata identified PPH to be a major cause of maternal morbidity with an estimated case-fatality rate of 7.5%.²⁷ While puerperal sepsis is a known cause of maternal mortality in India, we found only one study that provided estimated incidence of puerperal sepsis in six hospitals in the state of Gujarat.²⁸ The study reported an incidence of 4% which is higher than the 0.2% estimated through IndOSS-Assam.²⁸ A study that reviewed the records of all deliveries in a tertiary hospital in Chandigarh over a period of 15 years (January 1988 to December 2002) found 315 cases of septic abortion among 46,417 deliveries, giving an incidence of 0.7% which is comparable to that estimated through IndOSS-Assam in the two pilot hospitals, although the case-fatality rate in the Assam hospitals (31%) was higher than that estimated in the Chandigarh hospital (25%).²⁹ A hospital in central India estimated an incidence of 0.06% for uterine rupture over a period of 12 years (1989 to 2000) with a case-fatality rate of 6%.³⁰ It therefore appears that the incidence rates

of the six conditions estimated through the IndOSS-Assam are in general comparable with the incidence of these conditions estimated by hospital-based studies in others parts of India, but the reported case-fatality rates in the two hospitals in Assam are much higher, further justifying the need for establishing IndOSS-Assam.

Implications and future research

IndOSS-Assam has been shown to be a feasible and simple system to initiate surveillance of the major causes of maternal death in the state to estimate the incidence and case-fatality rates on an ongoing basis. An examination of the research platform alongside the feasibility study for setting up the surveillance system showed that through IndOSS-Assam detailed studies on these conditions can be conducted to identify the specific preventive and management factors that need to be improved to reduce deaths and severe complications associated with the conditions.¹⁸ For example, the study showed that iron deficiency anaemia during pregnancy is associated with increased odds of postpartum haemorrhage and the risk increased by 17-fold among women with moderate-severe anaemia who underwent induction of labour and by 19-fold among women who had infection and moderate-severe anaemia.¹⁸ These findings raise important questions about how pregnant women with moderate-severe anaemia should be managed during labour and delivery. At present, there are no clear guidelines and lessons learnt from further studies can be used to inform local and global guidance about management of labour and delivery among pregnant women with moderate-severe anaemia. The implementation and outcomes of the resultant guidelines/ actions can be monitored through the research and the surveillance platform of IndOSS-Assam. In addition to generating new hypotheses, the research platform will be used to investigate the gaps in management of women with specific life-threatening conditions on a rolling basis, as well as to identify good practices. The data collected through both platforms, being anonymous, avoids attribution of blame to any particular hospital or healthcare provider.

Involvement of local clinicians, policy makers, researchers, public health specialists, and academics from the inception of the project to testing the system shows that there is willingness and commitment to establish and scale up IndOSS-Assam to other public and private maternity hospitals in the state. We will explore the possibility of integrating the surveillance component of IndOSS-Assam with the existing health management information system of the National Health Mission in Assam. Integration would minimise duplication of effort, enable sustainable scaling up of the surveillance platform to all government health centres in the state, streamline data flow, and improve government ownership of the data facilitating timely action. In the longer term, the IndOSS-Assam model could be used in other states in India to encourage the establishment of a national system.

Competing interest: We have read and understood the BMJ policy on declaration of interests and declare that we have no competing interests.

Ethical approvals: The IndOSS-Assam feasibility study was approved by the institutional ethics committee of Srimanta Sankaradeva University of Health Sciences (No SSUHS/Ethics/2014/1, dated 27 June 2014); the Health Ministry's Screening Committee, Indian Council of Medical Research (No 5/7/12 12/14-RCH, dated 25.09.2014) and the University of Oxford's Oxford Tropical Research Ethics Committee (OxTRTEC Reference 57-14).

Funding statement: This study was funded by the John Fell Oxford University Press (OUP) Research Fund, PRAC/JF/131/124; and by the Srimanta Sankaradeva University of Health Sciences, Assam. Marian Knight is funded by a National Institute for Health Research (NIHR) Research Professorship.

The funding sources had no role in the study, and the researchers were independent from the funders. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Authorship and contributions: MN designed the study, trained the field staff, conducted literature searches, planned and carried out the data analysis, interpreted the data, and wrote the first draft of the article. MKC designed the study, trained the field staff, supervised data collection and data entry, and edited the manuscript. SSC designed the study, trained the field staff, supervised data collection and data entry, and edited the manuscript. SDK designed the study, trained the field staff, and edited the manuscript. UCS designed the study, trained the field staff, supervised data collection, and edited the manuscript. PW designed the study, and edited the manuscript. MK designed the study, contributed to the data analysis plan, data interpretation, and to writing the article.

Data sharing statement: This was a pilot project and there is no additional data from the study.

Transparency declaration: MN is the guarantor and affirms that the manuscript is an honest, accurate, and transparent account of the study being reported and no important aspects of the study have been omitted.


Exclusive licence statement: “I, Manisha Nair, the corresponding author of this article contained within the original manuscript which includes any diagrams & photographs within and any related or stand-alone film submitted (the Contribution”) has the right to grant on behalf of all authors and does grant on behalf of all authors, a licence to the BMJ Publishing Group Ltd and its licences, to permit this Contribution (if accepted) to be published in the BMJ and any other BMJ Group products and to exploit all subsidiary rights, as set out in our licence set out at: <http://www.bmj.com/about-bmj/resources-authors/forms-policies-and-checklists/copyright-openaccess-and-permission-reuse>.”

REFERENCE LIST

1. WHO, Unicef, UNFPA, et al. Trends in maternal mortality: 1990 to 2013. Estimates by WHO, UNICEF, UNFPA, The World Bank and the United Nations Population Division. Geneva: World Health Organisation, 2014.
2. Office of Registrar General India. Special bulletin on maternal mortality in India 2010-12: Sample registration system. New Delhi: Office of Registrar General, India - Vital Statistics Division, 2013.
3. National Health Mission. State wise information: Assam. Secondary State wise information: Assam. <http://nrhm.gov.in/nrhm-in-state/state-wise-information/assam.html>.
4. International Institute for Population Sciences (IIPS) and Macro International. National Family Health Survey (NFHS-3), 2005-06, India: Key Findings Mumbai: IIPS, 2007.
5. Health Administrator. Assam. Health Administrator 2004; XVI(1). <http://medind.nic.in/haa/t04/i1/haat04i1c.shtml> (accessed 24 March 2013).
6. National Health Mission. Janani Suraksha Yojana. Secondary Janani Suraksha Yojana. <http://nrhm.gov.in/nrhm-components/rmnch-a/maternal-health/janani-suraksha-yojana/background.html>.
7. Planning and Development Department; Government of Assam. Chief Minister's Vision for Women & Children 2016. http://planassam.info/Chief_Minister_Vision_for_Women_Children_2016.pdf (accessed 13 October 2015).
8. National Health Mission A. Maternal death reporting system (eMDR). Secondary Maternal death reporting system (eMDR). <http://nrhmassam.in/index.php>.
9. World Health Organisation. *Beyond the numbers : reviewing maternal deaths and complications to make pregnancy safer*. Geneva World Health Organisation, 2004.


10. Office of Registrar General and Census Commissioner India. Annual Health Survey 2010-11: Fact sheet. New Delhi: Ministry of Home Affairs, Government of India, 2012.
11. Office of Registrar General and Census Commissioner India. Annual Health Survey 2012-13: Fact sheet. New Delhi: Ministry of Home Affairs, Government of India, 2014.
12. Souza JP, Cecatti JG, Haddad SM, et al. The WHO Maternal Near-Miss Approach and the Maternal Severity Index Model (MSI): Tools for Assessing the Management of Severe Maternal Morbidity. *PloS one* 2012;**7**(8):e44129.
13. Halder A, Jose R, Vijayselvi R. Maternal mortality and derivations from the WHO near-miss tool: An institutional experience over a decade in Southern India. *Journal of the Turkish German Gynecological Association* 2014;**15**(4):222-7.
14. Sangeeta G, Leena W, Taru G, et al. Evaluation of severe maternal outcomes to assess quality of maternal health care at a tertiary center. *Journal of obstetrics and gynaecology of India* 2015;**65**(1):23-7.
15. Knight M, Kurinczuk JJ, Tuffnell D, et al. The UK Obstetric Surveillance System for rare disorders of pregnancy. *BJOG: An International Journal of Obstetrics & Gynaecology* 2005;**112**(3):263-65.
16. Knight M, Lindquist A. The UK Obstetric Surveillance System: Impact on patient safety. *Best Practice & Research Clinical Obstetrics & Gynaecology* 2013;**27**(4):621-30.
17. Knight M, Inoss. The International Network of Obstetric Survey Systems (INOSS): benefits of multi-country studies of severe and uncommon maternal morbidities. *Acta Obstetrica et Gynecologica Scandinavica* 2014;**93**(2):127-31.
18. Nair M, Choudhury MK, Choudhury SS, et al. The association between maternal anaemia and pregnancy outcomes: a cohort study in Assam, India.**[Under review]**.
19. National Health Mission A. Integrated MIS-GIS system. Secondary Integrated MIS-GIS system. <http://www.nrhmassam.info/login.php>.
20. Grewal A. Anaemia and pregnancy: Anaesthetic implications. *Indian Journal of Anaesthesia* 2010;**54**(5):380-86.
21. Sarma HKD, Sarma HK, Kalita AK. A prospective study of maternal near-miss and maternal mortality cases in FAAMCH, Barpeta; with special reference to its aetiology and management: First 4 months report. *J Obstet Gynaecol Barpeta* 2015;**1**:100-6.
22. Montgomery AL, Ram U, Kumar R, et al. Maternal Mortality in India: Causes and Healthcare Service Use Based on a Nationally Representative Survey. *PloS one* 2014;**9**(1):e83331.
23. Sri BS, Khanna R. Dead women talking: a civil society report on maternal deaths in India: Commonhealth and Jan Swasthya Abhiyan, 2014.
24. Rohilla M, Raveendran A, Dhaliwal L, et al. Severe anaemia in pregnancy: a tertiary hospital experience from northern India. *Journal of Obstetrics & Gynaecology* 2010;**30**(7):694-96.
25. Bhalerao A, Kulkarni S, Ghike S, et al. Eclampsia: Maternal and fetal outcome. *JSAFOG* 2013;**5**(1):19-21.
26. Sunita T, Desai RM, Hon N, et al. Eclampsia in a teaching hospital: incidence, clinical profile and response to magnesium sulphate by Zuspan's regimen. *IOSR J Dent Med Sci* 2013;**4**(2):01-05.
27. Shirazee HH, Saha SK, Das I, et al. Postpartum haemorrhage: a cause of maternal morbidity. *J Indian Med Assoc* 2010;**108**(10):663-6.
28. Hussein J, Ramani KV, Kanguru L, et al. The effect of surveillance and appreciative inquiry on puerperal infections: a longitudinal cohort study in India. *PloS one* 2014;**9**(1):e87378.
29. Jain V, Saha SC, Bagga R, et al. Unsafe abortion: A neglected tragedy. Review from a tertiary care hospital in India. *Journal of Obstetrics and Gynaecology Research* 2004;**30**(3):197-201.
30. Chhabra S, Bhagwat N, Chakravorty A. Reduction in the occurrence of uterine rupture in Central India. *Journal of Obstetrics and Gynaecology* 2002;**22**(1):39-42.

Figure-1: Flow-chart showing activities undertaken to examine the feasibility of establishing IndOSS-Assam



*electronic maternal death reporting system (e-MDR) of the national Health Mission in the state of Assam

Figure-2: IndOSS-Assam case notification card



Indian Obstetric Surveillance System

Assam

MONTHLY CASE NOTIFICATION CARD

(Please submit the card on the 4th of every month)


Date

Total deliveries in the month

| Sl. No | Conditions | Number of cases* | Number died* |
|--------|------------------------|--|--|
| 1 | Eclampsia | <input style="width: 100%; height: 30px; border: 1px solid black;" type="text"/> | <input style="width: 100%; height: 30px; border: 1px solid black;" type="text"/> |
| 2 | Postpartum haemorrhage | <input style="width: 100%; height: 30px; border: 1px solid black;" type="text"/> | <input style="width: 100%; height: 30px; border: 1px solid black;" type="text"/> |
| 3 | Puerperal Sepsis | <input style="width: 100%; height: 30px; border: 1px solid black;" type="text"/> | <input style="width: 100%; height: 30px; border: 1px solid black;" type="text"/> |
| 4 | Septic abortion | <input style="width: 100%; height: 30px; border: 1px solid black;" type="text"/> | <input style="width: 100%; height: 30px; border: 1px solid black;" type="text"/> |
| 5 | Uterine rupture | <input style="width: 100%; height: 30px; border: 1px solid black;" type="text"/> | <input style="width: 100%; height: 30px; border: 1px solid black;" type="text"/> |
| 6 | Anaemic heart failure | <input style="width: 100%; height: 30px; border: 1px solid black;" type="text"/> | <input style="width: 100%; height: 30px; border: 1px solid black;" type="text"/> |

**Please enter '0' (zero) if no cases / no death*


Figure-3: Causes of maternal deaths in Assam April 2012 to August 2014*



Source of data: Electronic maternal deaths reporting system (eMDR), National Health Mission, Assam

*causes of most deaths reported in 2011-12 were either unknown or not classified hence not included in this graph, n=2610.

Figure-4: Maternal deaths in Assam from April 2011 to August 2014: PLACE OF DEATH



Source of data: Electronic maternal deaths reporting system (eMDR), National Health Mission, Assam

Figure-5: Geographical mapping of future reporting hospitals for IndOSS-Assam and reporting hospitals taking part in the pilot study

