

Title:

Intrusive thoughts and images of intentional harm to infants in the context of maternal postnatal depression, anxiety and obsessive-compulsive disorder.

Authors:

Peter J Lawrence, Research Training Fellow and Clinical Psychologist, Anxiety and Depression in Young People Clinical Research Unit, School of Psychology & Clinical Language Sciences, University of Reading, UK; Honorary Clinical Psychologist, Department of Psychiatry, University of Oxford, UK.

Michelle G Craske, Distinguished Professor and Vice Chair, Department of Psychology, University of California, Los Angeles (UCLA), Los Angeles, CA, USA

Claire Kempton, Senior Psychological Therapist, Department of Psychiatry, University of Oxford, UK

Anne Stewart, Consultant Child and Adolescent Psychiatrist, Boundary Brook House, Oxford Health NHS Foundation Trust; Senior Clinical Lecturer, Department of Psychiatry, University of Oxford, UK.

Alan Stein, Professor of Child and Adolescent Psychiatry, Department of Psychiatry, University of Oxford, UK; Honorary Consultant Child and Adolescent Psychiatrist, Oxford Health NHS Foundation Trust.

Corresponding Author: Pete Lawrence; AnDY Clinical Research Unit, PCLS, Whiteknights Campus, Reading RG6 6AL e: p.j.p.lawrence@reading.ac.uk

Introduction

Intrusive thoughts are key features of depression, anxiety and obsessive-compulsive disorder (OCD) [1]. Such thoughts are also common in the general population, where their content is the same as found in those with a psychiatric disorder. Intrusive thoughts comprise unwanted negative thoughts and images that frequently intrude, are difficult to dismiss and, when dismissed, recur. Furthermore they lead to a narrowed focus of attention which, in turn, can impair a person's ability to respond to the external world. They can play an important role in maintaining the disorders in which they occur [2].

One form of intrusive thoughts that is particularly distressing in the postnatal period is thoughts of intentionally harming one's infant (see Box One). Such thoughts and images have been reported to occur in very nearly half of parents of infants in the general population [3]. While such intrusions are not in themselves indicative of risk, they are likely to be of particular significance when they occur as part of depression, anxiety disorders or OCD where they are often associated with great distress and shame. While much of the research in this area has been with mothers, these intrusive thoughts also frequently occur in fathers [4].

Should I enquire about intrusive thoughts?

A common concern of clinicians is that by enquiring about thoughts related to sensitive and undesirable behaviors (for example, suicidal thoughts), they might in some way contribute to the likelihood of a patient acting on such thoughts. However, as with suicidal thoughts, there is no evidence that clinicians increase this likelihood of acting on thoughts of harming an infant by enquiring about such thoughts. Rather, patients often feel relieved at being able to talk about these thoughts.

Parents very rarely spontaneously volunteer having intrusive thoughts of harming their baby, even when answering questions concerning their depression, anxiety disorder or OCD. This can be because of a feeling of shame and a fear that the baby may be removed from them by statutory services. However, mothers can be helped to understand that such intrusions are common and by themselves signify nothing sinister.

Insert Box One about here

How do I assess them?

These intrusive thoughts need to be carefully assessed, by a General Practitioner, Health Visitor or mental health professional, to distinguish them from those that should, in fact, trigger child protection and safeguarding proceedings.

The assessment is likely to reveal that the thoughts are intrusive, ego-dystonic (that is, they are inconsistent with the mother's view of herself), inconsistent with the mother's behaviour, and cause the mother distress and dismay and there is no intention to act on them or a history of harming the baby; and as such they pose no risk of harm to their infant.

Insert Box Two about here

The content of intrusions is such that patients are often unwilling to disclose them. Disclosure of intrusive thoughts is more likely within a trusting relationship, possibly one developed via continuity of care. Even so, on a first meeting, we have found that enquiry can be fruitful. We have found that, directly before enquiring about intrusive thoughts of harm, it can be help build a parent's trust by explaining to them that negative intrusive thoughts of all kinds - for example violent thoughts - are common and that we, as mental health professionals, also experience them. Following such explanation and disclosure, patients have been able to reveal, often with great relief, that they *have*, indeed, experienced just such thoughts. Assessing features of mothers' emotions and behaviours associated with these thoughts (see Box Two) has shown us that mothers almost invariably find the intrusions to be repugnant and distressing, that they have done nothing consistent with the intrusions and have no intention to do anything consistent with the intrusions.

Insert Box Three about here

Is further guidance available?

There is an increasing recognition of the importance of identifying and treating postnatal depression and anxiety as well as OCD (for example, NICE Guidelines for perinatal mental health).[5] The issue of screening for intrusive thoughts of harm to the baby or managing them, however, is not mentioned in these guidelines or any other standard information, and neither is such enquiry included in the screening instruments. The only easily available information of which we are aware that mentions this issue is a leaflet produced by the Royal College of Psychiatrists for families (<http://www.rcpsych.ac.uk/healthadvice/problemsdisorders/postnataldepression.aspx>) and the OCD-UK website (<http://www.ocduk.org/prenatal-postnatal-ocd>). Box Three contains important advice that can be given, confidently, to parents reporting intrusive thoughts of intentionally harming their infants.

Conclusion

Postnatal depression occurs in the UK in approximately 10-12% of mothers, and anxiety and OCD are also common in the postpartum period. Intrusive thoughts of harming one's baby are common amongst clinical and non-clinical samples alike. However, they may become more distressing and harder to cope with in mothers who are suffering from mental health problems. Such thoughts can be elicited through careful and sensitive questioning (especially emphasizing to parents that these are common and do not mean that they will act on them). Once parents are able to acknowledge and talk about these thoughts and appreciate that experiencing them is a very common experience of parents, they commonly diminish in significance and frequency. Where they persist, treatment for intrusive thoughts is available using standard cognitive behavioural techniques. [5]

Acknowledgements

We are grateful to Charlotte Granger, Peter Cooper, Lynne Murray and Louise Dalton for their help with this work and to the Wellcome Trust (grant no 090139) for funding our work.

References

1. Harvey A, Watkins E, Mansell W, Shafran R. Cognitive Behavioural Processes Across Psychological Disorders: A Transdiagnostic Approach to Research and Treatment. OUP, 2004, 376 p.
2. Clark DA, de Silva P. The nature of depressive and anxious, intrusive thoughts: Distinct or uniform phenomena? Behav Res Ther 1985; **23(4)**: 383-393.
3. Fairbrother N, Woody SR. New mothers' thoughts of harm related to the newborn. Arch Women's Ment Health 2008; **11(3)**: 221-9.
4. Abramowitz JS, Khandker M, Nelson CA, Deacon BJ, Rygwall R. The role of cognitive factors in the pathogenesis of obsessive-compulsive symptoms: a prospective study. Behav Res Ther; 2006. **44(9)**:1361-74.
5. NICE. (2015). Antenatal and Postnatal Mental Health: Clinical Management and Service Guidance. CG192. Retrieved from United Kingdom: <http://www.nice.org.uk/guidance/cg192/evidence>

Box One

Specific content of intrusive thoughts of intentionally harming one's infant.

- Shaking the baby
- Hitting the baby too hard during winding.
- Throwing the baby to the ground or against a wall.
- Puncturing the infant's fontanelle.
- Drowning the baby in the bath.
- Smothering the baby e.g., with a baby's milk bottle or pillow.
- Releasing the baby in a pram from the top of a hill / into traffic.

Accepted

Box Two

Features to assess

- Is the mother indifferent to the intrusive thoughts?
 - Is the mother emotionally unaffected by the thoughts?
 - Does the mother try to trigger the thoughts?
 - Does the mother decline help to manage the thoughts?
 - Does the mother intend to act out the thoughts?
 - Has the mother ever intentionally harmed her baby?
- Negative answers to these questions are entirely consistent with a *very low risk* of a mother deliberately harming their infant. Conversely, an affirmative answer to the first four points does not necessarily indicate that a mother *will* harm their infant. Rather, **the overall clinical picture** from answers to all of these questions is the more informative indicator. However, if the mother acknowledges intention to act out the thoughts or has a history of previously harming her baby, the risk needs to be taken seriously.
- Unlike in puerperal psychosis, intrusive thoughts of intentional harm to the infant are *ego dystonic*. That is, they are experienced as unacceptable and inconsistent with a person's sense of self.
 - The content of intrusions is important only so far as secondary risks might arise. For example, if a mother fears smothering the baby using the baby's milk bottle, they might refuse to feed the baby.

Box Three

Normalizing intrusive thoughts for concerned parents

- Intrusive thoughts or images of causing harm to one's infant are common in the general population.
- Experiencing the intrusive thoughts makes them no more likely to harm their infant intentionally than any other parent is to harm their own infant intentionally.
- There is no need to avoid triggers or situations that give rise to the intrusive thoughts or images. Avoiding them actually tends to increase the frequency of the thoughts and or images.