

Third-wave cognitive behaviour therapies for weight management: A systematic review and network meta-analysis

Emma R. Lawlor¹  | Nazrul Islam¹  | Sarah Bates²  | Simon J. Griffin^{1,3} |
Andrew J. Hill⁴ | Carly A. Hughes^{5,6} | Stephen J. Sharp¹ | Amy L. Ahern¹ 

¹MRC Epidemiology Unit, University of Cambridge, Cambridge, UK

²School of Health and Related Research, Faculty of Medicine, Dentistry and Health, University of Sheffield, Sheffield, UK

³Primary Care Unit, Institute of Public Health, University of Cambridge, Cambridge, UK

⁴Division of Psychological and Social Medicine, School of Medicine, University of Leeds, Leeds, UK

⁵Fakenham Medical Practice, Norfolk, UK

⁶Norwich Medical School, University of East Anglia, Norwich, UK

Correspondence

Emma R. Lawlor, MRC Epidemiology Unit, School of Clinical Medicine, University of Cambridge, Box 285 Institute of Metabolic Science, Cambridge Biomedical Campus, Cambridge CB2 0QQ, UK.

Email: emma.lawlor@mrc-epid.cam.ac.uk

Funding information

National Institute of Health Research (NIHR) Programme Grants for Applied Research, Grant/Award Number: RP-PG-0216-20010; Medical Research Council, Grant/Award Number: MC_UU_12015/4

Summary

This systematic review and network meta-analysis synthesized evidence on the effects of third-wave cognitive behaviour therapies (3wCBT) on body weight, and psychological and physical health outcomes in adults with overweight or obesity. Studies that included a 3wCBT for the purposes of weight management and measured weight or body mass index (BMI) pre-intervention and ≥ 3 months post-baseline were identified through database searches (MEDLINE, CINAHL, Embase, Cochrane database [CENTRAL], PsycINFO, AMED, ASSIA, and Web of Science). Thirty-seven studies were eligible; 21 were randomized controlled trials (RCT) and included in the network meta-analyses. Risk of bias was assessed using RoB2, and evidence quality was assessed using GRADE. Random-effects pairwise meta-analysis found moderate- to high-quality evidence suggesting that 3wCBT had greater weight loss than standard behavioural treatment (SBT) at post-intervention (standardized mean difference [SMD]: -0.09 , 95% confidence interval [CI]: -0.22 , 0.04 ; $N = 19$; $I^2 = 32\%$), 12 months (SMD: -0.17 , 95% CI: -0.36 , 0.02 ; $N = 5$; $I^2 = 33\%$), and 24 months (SMD: -0.21 , 95% CI: -0.42 , 0.00 ; $N = 2$; $I^2 = 0\%$). Network meta-analysis compared the relative effectiveness of different types of 3wCBT that were not tested in head-to-head trials up to 18 months. Acceptance and commitment therapy (ACT)-based interventions had the most consistent evidence of effectiveness. Only ACT had RCT evidence of effectiveness beyond 18 months. Meta-regression did not identify any specific intervention characteristics (dose, duration, delivery) that were associated with greater weight loss. Evidence supports the use of 3wCBT for weight management, specifically ACT. Larger trials with long-term follow-up are needed to identify who these interventions work for, their most effective components, and the most cost-effective method of delivery.

Abbreviations: 3wCBT, third-wave cognitive behaviour therapy; ACT, acceptance and commitment therapy; BMI, body mass index; CBT, cognitive behaviour therapy; CFT, compassion focused therapy; CI, confidence intervals; CONSORT, Consolidated Standards of Reporting Trials; DBT, dialectical behaviour therapy; GRADE, Grading of Recommendations, Assessment, Development and Evaluations; MBCT, mindfulness-based cognitive behavioural therapy; RCT, randomized controlled trial; ROBINS-I, Risk Of Bias In Non-randomised Studies of Interventions; RoB2, Risk of Bias 2; SBT, standard behavioural treatment; SD, standard deviation; SMD, standardized mean difference; TIDieR, Template for Intervention Description and Replication.

Emma R. Lawlor and Nazrul Islam, joint first authors.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Obesity Reviews* published by John Wiley & Sons Ltd on behalf of World Obesity Federation

KEYWORDS

network meta-analysis, obesity, third-wave behavioural therapy, weight loss

1 | BACKGROUND

Although behavioural interventions are effective at helping people to lose weight, many people struggle to sustain effective weight management behaviours over extended periods due to a combination of biological, psychological, social, and environmental factors that drive weight gain.^{1,2} Standard behavioural programmes can be effective in the short term, but less so in the longer term.^{3–6} These usually combine diet and physical activity advice with core behavioural change techniques including goal setting, self-monitoring, problem solving, and planned social support.⁷ It has been proposed that third-wave cognitive behaviour therapies (3wCBT), including acceptance and commitment therapy (ACT), dialectical behaviour therapy (DBT), mindfulness-based cognitive behavioural therapy (MBCT), and compassion-focused therapy (CFT),^{8–10} may have better short- and long-term outcomes.²

The theoretical case for 3wCBT for weight management has been well articulated.² In brief, these therapies encourage people to accept aversive internal experiences (eg, food cravings, physical discomfort) rather than avoid them. Increased present-moment, non-judgemental awareness and psychological flexibility may assist an individual in recognizing internal and external cues to overeat and alter behavioural responses to be more in line with their values. Fostering a compassionate attitude towards the self could also help prevent discouragement following minor lapses.^{2,8,9}

However, the evidence of their superior effectiveness is less clear. Previous systematic reviews and meta-analyses primarily focused on mindfulness- and/or acceptance-based interventions.^{11–18} Three reviews^{13,15,16} have reported a quantitative synthesis of pre-intervention to post-intervention change without comparing the effect against a comparator. Two of these three reviews reported a “small” pre-intervention to post-intervention change in weight¹⁵ or body mass index (BMI)¹³ while the other study¹⁶ reported a null effect on BMI. Critically, only one review¹⁴ reported a meta-analytic synthesis that compared the effectiveness of mindfulness- and acceptance-based interventions with those in other active interventions and control arms using appropriate statistical methods. A small but significant difference in weight or BMI was reported at post-intervention, favouring mindfulness and acceptance-based interventions over comparator arms. Subgroup analysis suggested that the effect may only hold when the comparator is waitlist control. In that review, there was no restriction on the minimum follow-up duration and outcomes were analysed at 1-month post-intervention (or the closest measurement to this). Thus, the pooled estimates reflected a mix of very short-term and longer term effects. Moreover, without a restriction on minimum BMI, these findings are less relevant from a policy perspective because behavioural weight management programmes are intended for adults with overweight/obesity.^{19,20} This concern is

compounded by the finding that a lower BMI was associated with a larger effect size.

To our knowledge, no head-to-head trial exists that has compared the effectiveness of different types of 3wCBT on weight management. In the absence of head-to-head trials, network meta-analysis can estimate the indirect evidence on the comparative effectiveness of different types of 3wCBT. The proposed mechanism for the superior effects of 3wCBT is through improvements in eating behaviour and psychological outcomes, so it is also important to synthesize evidence on the impact of 3wCBT on these outcomes. In addition, evidence synthesis of the effect of 3wCBT on eating behaviour and psychological outcomes has been limited to pre-intervention to post-intervention change^{13,15,16} and has not considered longer follow-up periods.

To address these knowledge gaps, we conducted the most comprehensive, inclusive, and relevant review and quantitative synthesis of available evidence to date. We included different types of 3wCBT beyond mindfulness and acceptance-based interventions. Our main objectives were (a) to evaluate the effectiveness of 3wCBTs on weight management by pooling the pre-intervention to post-intervention change effect estimates across all study types, (b) to compare the effectiveness of 3wCBTs on weight management against no/minimal interventions and standard behavioural treatment (SBT) separately using random-effects pairwise meta-analysis of randomized control trials (RCTs), (c) to estimate the comparative effectiveness of different types of 3wCBTs on weight management using random-effects network meta-analysis of RCTs, (d) to evaluate the impact of 3wCBT on eating behaviour and psychological and physical health outcomes, and (e) to provide a detailed description of intervention characteristics and to identify whether any of these are associated with better weight change outcomes by using meta-regression.

2 | METHODS

2.1 | Protocol and registration

The protocol was registered on Prospero (CRD42018088255) prior to article screening.²¹

2.2 | Eligibility criteria

Participants were community-dwelling adults (≥ 18 years) with overweight or obesity ($\text{BMI} \geq 25 \text{ kg/m}^2$) seeking assistance with weight management. Studies had to include a 3wCBT intervention for the purpose of weight management. Multi-component interventions

(eg, including diet and physical activity advice) were acceptable, with no restriction placed on the proportion of the intervention using 3wCBT. Interventions could be of any duration. Comparisons were (a) no/minimal intervention, (b) SBT, or (c) no comparator (single-arm pre-intervention to post-intervention studies). We defined SBT as structured programmes providing diet and/or physical activity advice and standard behaviour change techniques (eg, goal setting, self-monitoring, problem solving, social support). The primary outcome was body weight or BMI. Studies needed to measure this pre-intervention and at least 3-months post-baseline. Secondary outcomes were stress, anxiety, depression, meta-cognition, eating attitudes, eating behaviours, body satisfaction, quality of life, blood pressure, lipids, glycaemia, and adherence to treatment. All outcomes reported at 3-months from baseline and beyond were extracted. All settings apart from laboratories were eligible. We included research articles, theses, and dissertations reporting RCTs, non-RCTs, prospective cohort and case series studies.

2.3 | Information sources

Databases (MEDLINE, CINAHL, Embase, Cochrane database [CENTRAL], PsycINFO, AMED, ASSIA, and Web of Science) were searched by ERL from inception with no restrictions, using keywords and subject heading searches relating to the concepts: (a) third-wave CBTs and (b) overweight, obesity, or weight management (see Table S1). The initial search was conducted on 16 January 2018, and an updated search was conducted on 25 September 2019. Reference lists of eligible studies and relevant reviews were searched, and authors of relevant abstracts were contacted to identify whether findings had been accepted for publication.

2.4 | Study selection

Titles and abstracts, then full texts, were screened independently by two of three researchers, with a third reviewer adjudicating uncertainty or disagreement. Study authors were contacted to resolve any questions about eligibility. Non-English language texts were translated into English by colleagues who were fluent in that language.

2.5 | Data collection process

Data were extracted independently by two of four researchers using a form based on the Cochrane data extraction form,²² the Consolidated Standards of Reporting Trials (CONSORT) 2010 statement,²³ and the Template for Intervention Description and Replication (TIDieR) checklist²⁴ and cross-checked for consistency. Attempts were made to contact authors to retrieve missing data. If there was no response after two attempts, we used the data available in the published work.

2.6 | Risk of bias

Two researchers assessed studies independently using the Risk of Bias 2 tool (RoB 2)²⁵ or the Risk Of Bias In Non-randomized Studies of Interventions tool (ROBINS-I),²⁶ dependent upon study design. A third reviewer adjudicated inconsistency. The quality of evidence was assessed using the Grading of Recommendations Assessment, Development and Evaluation (GRADE) approach, which classifies studies as "high", "moderate", "low", or "very low" quality.²⁷

2.7 | Missing data

For the primary outcomes, where standard deviations (SDs) for mean change were missing and not provided following author correspondence, these were imputed using the following methods, in order of prioritization: (a) imputed from other time points within same study, (b) estimated from *t* statistics, Cohen's *d*, *p*-values or confidence intervals (CIs),²⁸⁻³² (c) estimated using a correlation coefficient of .97, based on empirical data from seven studies^{28,30,33-37} (17 estimates) that reported SDs for baseline, follow-up, and mean change.³⁸⁻⁴⁶ Insufficient data prevented this approach for secondary outcomes, so we used a correlation of .7 as in previous studies.^{47,48}

2.8 | Synthesis of results

Stata/SE v.14.2⁴⁹ was used for all statistical analyses. Following guidance,⁵⁰ we focused on 95% CIs, rather than statistical significance. For example, unlike conventional interpretations, we did not outright interpret an effect estimate "non-significant" if the lower or upper bound of the 95% CI was slightly above/below the null value; we interpreted them as "suggestive" of an effect.

2.8.1 | Pooled estimates of intervention-specific effects from all study types

Intervention-specific effects (post-intervention minus pre-intervention) were estimated by pooling effect estimates from all study designs. Due to heterogeneity in outcome measurement, effect estimates were reported as standardized mean change from the random-effects meta-analysis.⁵¹ Effect estimates were reported at the earliest measurement post-intervention (≥ 3 months from baseline) and at 3, 6, 9, 12, 18, 24, and 36 months from baseline. Outcomes falling between these time points were included with the closest time point.

2.8.2 | Intervention comparisons: Direct evidence from pairwise meta-analysis of RCTs

The direct effect comparing 3wCBT against (a) no/minimal intervention and (b) SBT was estimated using random-effects⁵¹ pairwise

meta-analysis of RCTs. The standardized mean difference (SMD) calculated using Hedges' method and 95% CI were reported.⁵²

2.8.3 | Intervention comparisons: Indirect and mixed evidence from network meta-analysis of RCTs

To compare types of 3wCBT, random-effects network meta-analysis of RCTs was conducted to estimate the indirect and mixed (direct plus indirect) evidence.⁵³ Basic assumptions were checked conceptually and statistically.⁵³ For example, to avoid violating the transitivity assumption, which requires that the comparator arm (eg, the waitlist control) is comparable across the trials, the comparators (SBT and no/minimal intervention) were not pooled/used together. Similarly, the intervention arms were dropped (namely resistance exercise³⁵ and food environment modification⁵⁴) if they were not comparable with other intervention arms. The consistency assumption was checked statistically to see if the direct and indirect effect estimates were comparable enough to pool them together into the mixed evidence.^{55,56} Effect estimates were reported as SMD and 95% CIs. The relative ranking probability of each intervention being the best treatment was estimated using rankograms.⁵⁷

2.8.4 | Sensitivity analysis

The influence of individual studies on weight change of 3wCBT compared with no/minimal intervention or SBT was examined using influence plots, where one study was removed at a time to see its effect on the overall estimate.⁵⁸

2.8.5 | Meta-regression on intervention and study characteristics

Where at least 10 studies provided relevant data,⁵⁹ meta-regression was used to identify potential sources of heterogeneity for pre-specified characteristics: number of sessions (continuous; <12 vs. ≥12 sessions), duration of intervention (<3 months vs. ≥3 months, <6 months vs. ≥6 months), method of delivery (face-to-face vs. remote; group vs. individual), and risk of bias (low, some concerns, high).

2.8.6 | Secondary outcomes

Only a small number of studies reported the secondary outcomes at each follow-up time point, so the first time point post-intervention was used and network meta-analysis was not conducted. Pooled and pairwise meta-analyses were conducted for secondary outcomes reported in two or more studies.

For consistency, we defined "change" as post-intervention minus pre-intervention values, so a negative change estimate indicates that the outcome decreased after the intervention.

3 | RESULTS

After duplicate removal, 8755 titles and abstracts were screened and 215 full text articles were assessed. Two additional studies and four articles related to already included studies were identified from reference lists⁶⁰⁻⁶⁴ and contacting an author.⁶⁵ Fifty articles reporting 37 studies met the inclusion criteria (Figure 1). Two studies were excluded from the meta-analyses due to cointerventions (pharmacotherapy⁶⁶ and bariatric surgery³⁹). Thirty-five studies were used in the pooled-effects meta-analysis of pre-intervention to post-intervention changes, and the 21 RCT design studies were used in the network meta-analysis that compared different interventions.

3.1 | Study characteristics

Seventeen studies^{30-33,36,41,45,46,66-74} used a two-group RCT, four^{35,40,44,54} used a three-group RCT, and one used a two-group cluster RCT design.⁴² Fourteen studies^{28,29,34,38,43,64,65,75-80} used a pre-intervention to post-intervention one-group design, and one study was a non-randomized three-group study.³⁹ The majority of studies were conducted in the United States ($n = 28$). The other studies were conducted in New Zealand,³⁴ Italy,³⁹ United Kingdom,^{41,42,77} the Netherlands,^{43,64} Finland,⁴⁴ and Portugal⁷⁰ (Table 1).

3.1.1 | Participant characteristics

Studies included 2726 participants and the sample size ranged from 10^{29,64} to 283.⁵⁴ Seventy-five percent of participants were female ($n = 2035/2726$), with 12 studies^{28,32-34,38,45,64,65,67,70,80} focusing exclusively on females. Mean age was 46 years (ranged from 21³³ to 58 years⁴⁵), and mean BMI was 35.6 kg/m² (Table 1 and Table S2a,b).

3.1.2 | Intervention characteristics

Twenty-two studies evaluated MBCT,^{28,29,31-33,35,36,38,40,42,45,46,64,65,67-70,74,77,78} eleven evaluated ACT-based interventions,^{30,34,44,54,71-73,75,76,79,80} three evaluated DBT^{39,43,66} (one⁶⁶ in combination with pharmacotherapy), and one evaluated CFT.⁴¹ Twenty-seven studies^{28-33,35,36,40,44-46,64,65,67-73,75,77,78,80,81} used primarily face-to-face, group-format delivery. One study⁴³ had an initial individual face-to-face session before delivery of group sessions. Five other studies^{38,39,41,42,66} used face-to-face group sessions along with another mode: emails,⁴² telephone calls,^{39,41} individual diet counselling³⁸ and a website for pharmacology support.⁶⁶ One study⁷⁶ used individual face-to-face lifestyle counselling and telephone delivery. Three delivered interventions on an individual, remote basis using email⁷⁴ and online website,^{34,79} two of these included telephone support.^{74,79} One study⁴⁴ had two intervention arms with the same content delivered face to face or through mobile telephone. Most interventions include home-based skills practice between sessions.




FIGURE 1 PRISMA flow diagram

Intervention duration varied, with two lasting less than a week,^{28,30} twelve studies between 1 and 3 months,^{29,32,33,36,42,44,65,66,77,79,80} and nine studies^{31,34,40,43,67,68,70,76,78} between 3½ and 5½ months in length. Thirteen studies lasted for 6 months or more, with five of these being 12 months in length.^{39,45,54,72,73} Hamel et al⁶⁴ did not report intervention length.

All studies were delivered on a weekly or alternating weekly basis, apart from two: a one-off 1-day workshop and 5-day residential retreat.^{28,30} Several interventions had an “active phase,” then an extended period with less regular sessions or telephone follow-up.^{38,40,45,54,68-73} Most interventions were implemented at a university,^{30,35,36,40,42,45,46,54,66,68} with other venues including primary care units and hospitals,^{41,70,76,77} yoga retreat centres,^{28,65} a community and oncology practice,³⁸ YMCA,²⁹ and participants' place of employment⁸⁰ (Table 1 and Table S3a,b).

3.2 | Risk of bias

Of the RCTs, the risk of bias was rated as 'high' in four,^{36,41,42,74} 'some concern' in eleven,^{30,31,33,35,40,45,46,66,69-71} and 'low' in seven studies^{32,44,54,67,68,72,73} (Table S4a). Of the 15 non-RCTs, the risk of bias was rated as 'serious' in nine,^{28,38,39,64,65,78-80} and 'moderate' in six^{29,34,43,75-77} studies (Table S4b). The quality of the evidence was

different for different comparisons, dependent on studies included. For the comparison between 3wCBT and no/minimal intervention at post-intervention, the quality of evidence was 'high' (three studies),^{40,44,67} for the comparison between 3wCBT and SBT, the quality of evidence was 'moderate' at post-intervention (nineteen studies)^{30-33,35,36,40-42,45,46,54,68-74} and 'high' at 12 months (five studies),^{45,54,68,72} 18 months (three studies),^{68,71,73} 24 months (two studies),^{72,73} and 36 months (one study,⁷² direct evidence only) from baseline. Details of study quality for all comparisons are reported Table S5a-c.

3.3 | Intervention effects on body weight or BMI

Twenty-five studies^{28-33,35,36,38,42,44-46,54,65,67-69,73-78} reported an absolute weight change (kg or lb), four studies^{71,72,79,80} reported percent change from baseline weight, and six studies^{34,40,41,43,64,70} reported BMI change.

3.3.1 | Pooled estimates of intervention-specific effects from all study types

Standardized mean change in weight or BMI for 3wCBT was -0.84 (95% CI: -1.06 , -0.62 ; $N = 35$; $I^2 = 93\%$) from baseline to post-

TABLE 1 Characteristics of included studies

First author, year	Population		Intervention			Comparison		Outcomes			
	N	Age, years ^a	BMI, kg/m ^{2a}	Female; N, %	Intervention (s)	Primary delivery mode	Group or individual	Length, months	Comparison	Measurements	Time points, months ^b
Randomized controlled trials											
Blevins, 2008 ³³	41	20.7 (1.4)	29.6 (1.9)	41 (100)	MBCT	Face to face	Group	2	SBT	Weight; anxiety; depression; binge eating; body dissatisfaction	2, 5
Carpenter, 2017 ⁷⁴	75	47.3 (10.0)	31.5 (2.3)	69 (92)	MBCT	Telephone and email	Individual	6	SBT	Weight; anxiety; depression; stress; psychological flexibility; binge eating; emotional eating; mindful eating	6
Daubenmier, 2011 ^{67,87}	47	MBCT: 40.4 (8.0); No/min: 41.4 (6.7)	31.2 (4.8)	47 (100)	MBCT	Face to face	Group	4	No/min	Weight; anxiety; stress; disinhibition; emotional eating; dietary restraint;	4
Daubenmier, 2016 ^{68,88}	194	MBCT: 47.2 (13.0); SBT: 47.8 (12.4)	MBCT: 35.4 (3.5); SBT: 35.6 (3.8)	MBCT: 79 (79); SBT: 81 (86)	MBCT	Face to face	Group	5½	SBT	Weight; DBP; SBP; fasting glucose; HbA1c; HDL; LDL; TG; TG HDL ratio; waist circumference	3, 6, 12, 18
Davis, 2008 ³⁵	71	45.1 (8.3)	32.9 (3.7)	63 (89)	MBCT	Face to face	Group	6	SBT, SBT + RE	Weight; mindfulness; disinhibition; emotional eating; dietary restraint; hunger; body dissatisfaction	3, 6
Goldbacher, 2016 ³¹	79	45.6 (10.5)	36.2 (4.1)	75 (95)	MBCT	Face to face	Group	5	SBT	Weight; emotional eating	5
Kristeller, 2014 ⁴⁰	150	46.6	40.3	132 (88)	MBCT	Face to face	Group	5¼	SBT, No/min	BMI; depression; binge eating; disinhibition; dietary restraint; hunger	6, 9
Lee, 2017 ³⁶	53	47.7 (11.3)	34.5 (4.8)	48 (91)	MBCT	Face to face	Group	3	SBT	Weight; stress; mindfulness; mindful eating; DBP; SBP; waist circumference	3, 6, 9

(Continues)

TABLE 1 (Continued)

First author, year	Population			Intervention			Comparison		Outcomes		
	N	Age, years ^a	BMI, kg/m ^{2a}	Female; N, %	Intervention (s)	Primary delivery mode	Group or individual	Length, months	Comparison	Measurements	Time points, months ^b
McKee, 2014 ⁴²	60	37.6 (13.5) ^c	32.6 (4.9) ^c	40 (72) ^c	MBCT	Face to face and email	Group and individual	2	SBT	Weight; mindfulness; waist circumference	2, 3
Miller, 2012 ^{69,89}	68	MBCT: 53.9 (8.2); SBT: 54.0 (7.0) ^c	MBCT: 36.2 (1.2); SBT: 36.1 (1.2) ^c	33 (63.5) ^c	MBCT	Face to face	Group	6	SBT	Weight; anxiety; depression; mindfulness; disinhibition; dietary restraint; hunger; fasting glucose; HbA1c; waist circumference	3, 4, 6
Palmeira, 2017 ^{37,70}	73	MBCT: 42.0 (8.8); SBT: 42.7 (8.4)	MBCT: 34.8 (5.26); SBT: 33.7 (4.8)	73 (100)	MBCT	Face to face	Group	3½	SBT	BMI; QoL; mindfulness; disinhibition; emotional eating; total cholesterol; waist circumference	3.5
Raja-Khan, 2017 ^{32,61}	86	44.5 (12.5)	38.9 (8.7)	86 (100)	MBCT	Face to face	Group	2	SBT	Weight; anxiety; depression; stress; mindfulness; DBP; SBP; fasting glucose; HbA1c; LDL	2, 4
Smith, 2018 ⁴⁵	40	MBCT: 58.6 (4.7); SBT: 58.6 (5.2) ^c	MBCT: 34.7 (4.3); SBT: 38.2 (7.1) ^c	36 (100) ^c	MBCT	Face to face	Group	12	SBT	Weight; binge eating	1½, 4, 9, 12
Spadaro, 2017 ^{46,90}	49	45.2 (8.2) ^c	32.5 (3.7) ^c	40 (87) ^c	MBCT	Face to face	Group	6	SBT	Weight; mindfulness; disinhibition; dietary restraint; hunger	3, 6
Butryn, 2017 ^{54,91}	283	ACT: 53.2 (9.4); SBT: 53.0 (9.3); BT + E: 53.4 (10.3)	ACT: 35.2 (4.6); SBT: 35.0 (5.2); BT + E: 35.4 (5.2)	ACT: 84 (82); SBT: 67 (76); BT + E: 72 (77)	ACT	Face to face	Group	12	SBT, BT + E	Weight	6, 12
Fletcher, 2011 ³⁰	72	ACT: 53.1 (11.1); SBT: 52.1 (12.6)	ACT: 36.2 (0.6); SBT: 34.7 (0.6)	60 (83)	ACT	Face to face	Group	1 day	SBT	Weight; anxiety; depression; stress; psychological flexibility; DBP; SBP	1 week (not weight/BMI), 3
(Continues)											

(Continues)

TABLE 1 (Continued)

First author, year	Population		Intervention		Comparison		Outcomes	
	N	Age, years ^a	BMI, kg/m ^{2a}	Female; N, %	Primary delivery mode	Group or individual	Length, months	Measurements
Forman, 2013 ⁷¹	128	45.7 (12.8)	34.1 (3.6)	NR	ACT	Group	10	Weight
Forman, 2016 ^{72,92}	190	51.6 (10.1)	36.9 (5.8)	156 (82)	ACT	Group	12	Weight
Lillis, 2016 ^{62,73,93}	162	50.2 (10.9)	37.6 (5.3)	138 (85)	ACT	Group	12	Weight; disinhibition
Sairanen, 2017 ^{44, 60,63,94}	219	49.5 (7.4)	31.3 (2.9)	185 (85)	ACT	Group Face to face Mobile	2	No/min Weight; mindfulness; psychological flexibility; disinhibition; emotional eating; dietary restraint; intuitive eating
Loader, 2013 ⁴¹	36	45.4 (9.5)	46.7 (0.3)	25 (69)	CFT	Group and individual	6	BMI; disinhibition; emotional eating; dietary restraint
Adler, 2008 ⁶⁶	17	49.4 (11.4)	37.7 (10.1)	15 (88)	DBT + O	Group and individual	3	SBT + O 3, 4.5
Preintervention to post-intervention studies								
Braun, 2012 ²⁸	37	Range: 32-65	NR	NR	MBCT	Group	5 days	N/A Weight; mindfulness; stress
Braun, 2016 ⁶⁵	S1: 22; S2: 21	S1: 48.2 (14.3); S2: 49.4 (10.7)	S1: 30.8 (4.2); S2: 35.5 (6.8)	S1: 22 (100); S2: 21 (100)	MBCT	Group	2½	N/A Weight; emotional eating; mindful eating
Chung, 2016 ³⁸	26	50.1 (9.0) ^c	35.1 (4.0) ^c	22 (100) ^c	MBCT	Group and individual	6	N/A Weight; mindful eating
Dalen, 2010 ²⁹	10	44 (8.7)	36.9 (6.2)	7 (70)	MBCT	Group	1½	N/A Weight; anxiety; binge eating; depression; dietary restraint; disinhibition; stress
Hamel, 2010 ⁶⁴	10	50.4 (13.2)	29.1 (3.2)	9 (90)	MBCT	Group	NR	N/A BMI; emotional eating; hunger; mindful eating; QoL

(Continues)

TABLE 1 (Continued)

First author, year	Population		Intervention		Comparison		Outcomes	
	N	Age, years ^a	BMI, kg/m ^{2a}	Female; N, %	Primary delivery mode	Group or individual	Length, months	Time points, months ^b
Hanson 2019 ⁷⁷	53	45.6 (11.3)	48.5 (9.2)	16 (30.2)	MBCT	Group	2	Weight; emotional eating ½, 1, 1½, 2, 8
Lundgren, 2003 ⁷⁸	33	44.8 (9.0) ^c	31.1 (3.6) ^c	16 (84) ^c	MBCT	Group (unclear)	5	Weight; QoL
Andalocio 2018 ⁷⁶	23	NR	39.9	21 (91.3)	ACT	Individual	4	Weight; waist circumference
Boucher, 2016 ³⁴	40	44.8 (3.1)	32.9 (6.0)	40 (100)	ACT	Individual	3½	BMI; intuitive eating
Bradley, 2017 ^{79,95}	20	54.3 (12.1)	NR	17 (85)	ACT	Individual	2½	Weight
Forman, 2009 ⁸⁰	29	43.7 (9.8)	35.8 (5.4)	29 (100)	ACT	Group	3	Weight; dietary restraint; disinhibition; emotional eating; mindfulness; QoL
Niemeier, 2012 ⁷⁵	21	52.2 (7.6)	32.8 (3.4)	19 (91)	ACT	Group	6	Weight; dietary restraint; disinhibition; hunger
Gallé, 2017 ³⁹	153	DBT:34 (3.8) IIT: 33 (4.2) SBT: 32 (5.1)	45.8 (6.4)	153 (100)	DBT	Group and individual	12	Weight
Roosen, 2012 ⁴³	35	39.2 (11.0)	35.4 (2.6)	30 (86)	DBT	Group and individual	5	BMI; depression; dietary restraint; disinhibition; emotional eating

Abbreviations: ACT, acceptance and commitment therapy; BMI, body mass index; BT + E, behaviour therapy with environmental change; CFT, compassion-focused therapy; DBT, dialectical behavioural therapy; DBT + O, DBT + Orlistat; IIT, interpersonal individual treatment; MBCT, mindfulness-based cognitive behavioural therapy; No/min, no/minimal intervention; NR, not reported; QoL, quality of life; RE, resistance exercise; S1, Study 1; S2, Study 2; SBT, standard behavioural treatment; SBT + O, SBT + Orlistat.

^aMean (SD) or range.

^bTime since randomization/baseline.

^cAmong completers/participants included in analysis.

^d6 and 9 months recorded but not reported in the article.

intervention (equating an absolute weight change of 5.5 kg). Weight change by types of 3wCBT at different time points is reported in Table S6. There was a pattern of weight loss (relative to baseline) for DBT up to 12 months, for MBCT up to 18 months, and for ACT up to 36 months. However, there was high heterogeneity and few studies at later time points. There was no evidence of weight loss following CFT, but this was based on one poor-quality study⁴¹ at a single 3 month follow-up from baseline.

3.3.2 | Intervention comparisons: Direct evidence

Figure 2 summarizes the results of the pairwise random-effects meta-analysis, and Table S5a presents the quality of evidence for all direct comparisons using GRADE.²⁷ Most individual studies were small and did not find evidence of a difference between interventions. However, when findings were meta-analysed, there was high-quality evidence to suggest greater weight loss for 3wCBT compared with no/minimal intervention at post-intervention and 9 month follow-up from baseline. There was moderate-quality evidence based on 19 studies,^{30-33,35,36,40-42,45,46,54,68-74} suggesting that 3wCBT had greater weight loss than SBT at post-intervention, and high-quality evidence suggesting that 3wCBT had greater weight loss than SBT at 12 months (five studies^{45,54,68,72,73}) and 24 months (two studies^{72,73}). Only ACT interventions provided data for the 24 month comparison. Estimates at 6 and 18 months also suggested greater weight loss for 3wCBT versus SBT, but there was no evidence of a difference between the two groups at 3 and 9 months.

3.3.3 | Sensitivity analysis

In the influence plot analysis, removal of one study at a time did not have any effect on the overall effects estimates from the pairwise meta-analysis of weight change.

3.3.4 | Intervention comparisons: Indirect and mixed evidence

Network meta-analysis was conducted up to 18 months post-baseline, as only a single pairwise comparison (ACT vs. SBT) was reported at 24 and 36 months. Intervention networks at each time point are summarized in Figure 3. Estimates from the network meta-analysis are summarized in Figure 4. ACT produced greater weight loss than no/minimal intervention at post-intervention and 9 months; comparisons at 3 and 6 months, however, did not provide evidence of superior effectiveness of ACT. Comparisons between ACT and SBT suggested greater weight loss for ACT post-intervention. There was no evidence of a difference at other time points. Compared with MBCT, ACT had greater weight loss at 9 months; comparisons at other time points did not show evidence of a difference.

Comparisons between no/minimal intervention and MBCT did not provide evidence of a difference at any time point, and there was no consistent pattern of effects. Comparisons between MBCT and SBT suggested that SBT was more effective at 9 months, but estimates at 12 and 18 months suggested that MBCT was favoured. When CFT was compared with the other interventions, CIs were wide with no comparisons favouring CFT.

When interventions were relatively ranked, ACT was the best intervention post-intervention and at 3, 6, and 9 months post-baseline. MBCT was the best ranking intervention at 12 and 18 months post-baseline; however, this was based on only five studies (two MBCT) and three studies (one MBCT), respectively (Figure S1).

In terms of absolute weight change, for example, the SMD in weight between 3wCBT and SBT equates to a difference of 0.6 kg post-intervention and 1.4 kg at 24-month follow-up from baseline.

3.4 | Interventions effects on secondary outcomes

Pooled arm-specific estimates (standardized mean change) of the effect of 3wCBT (combined) on secondary outcomes are presented in Figure S2. Pairwise estimates (SMD) from RCTs comparing 3wCBT and no/minimal intervention are presented in Figure S3; those comparing 3wCBT and SBT are presented in Figures 5A, 5B, and 5C.

3.4.1 | Psychological outcomes

Pooled arm-specific estimates showed a reduction in anxiety (N = 7),^{29,30,32,33,67,69,74} depression (N = 9),^{29,30,32,33,40,43,69,74,78} and stress (N = 8)^{28-30,32,36,67,74,78} following 3wCBT. When compared with no/minimal intervention, estimates suggested greater reductions in anxiety for 3wCBT (N = 1),⁶⁷ a greater reduction in depression for 3wCBT (N = 1),⁴⁰ but found no evidence of a difference in stress (N = 1).⁶⁷ Pairwise comparisons found no evidence of differences between 3wCBT and SBT for anxiety (N = 5),^{30,32,33,69,74} depression (N = 6),^{30,32,33,40,69,74} or stress (N = 4).^{30,32,36,74}

Pooled arm-specific estimates (N = 4)^{64,70,78,80} suggested an increase in quality of life following 3wCBT. One study⁷⁰ reported a greater increase in quality of life in 3wCBT versus SBT.

Pooled arm-specific estimates (N = 9)^{32,35,36,42,44,46,69,70,80} showed an increase in mindfulness with 3wCBT. The estimate from one study⁴⁴ suggested a greater increase in mindfulness for 3wCBTs versus no/minimal intervention. Pairwise estimates (N = 7)^{32,35,36,42,46,69,70} suggested a greater increase in mindfulness in 3wCBT versus SBT.

Pooled arm-specific estimates (N = 5)^{30,34,44,74,75} showed that psychological flexibility increased after 3wCBT. One study⁴⁴ showed a greater increase in psychological flexibility for 3wCBT versus




FIGURE 2 Weight change comparing third-wave CBT and no/minimal or standard behavioural treatment from random-effects pairwise meta-analysis. Time points are months since baseline unless otherwise specified. CBT, cognitive behaviour therapy; MBCT, mindfulness-based cognitive behaviour therapy; No/min, no/minimal intervention; SBT, standard behavioural treatment; SMD, standardized mean difference




FIGURE 3 Network of interventions at different follow-up from baseline time points. Nodes are weighted by the number of studies involved in each intervention while the edges are weighted by the number of studies involved in each comparison. Time points are months since baseline unless otherwise specified. ACT, acceptance and commitment therapy; CFT, compassion-focused therapy; MBCT, mindfulness-based cognitive behaviour therapy; No/min, no/minimal intervention; SBT, standard behavioural treatment

no/minimal intervention. Pairwise estimates ($N = 2$)^{30,74} found no evidence of a difference between 3wCBT and SBT.

3.4.2 | Eating behaviour

Pooled arm-specific estimates showed a decrease in binge eating ($N = 6$)^{29,33,40,45,74,78} disinhibition ($N = 14$)^{29,35,40,41,43,44,46,67,69,70,73,75,78,80} and hunger ($N = 7$)^{29,35,40,46,69,75,78} and suggested a decrease in emotional eating ($N = 13$)^{31,35,41,43,44,64,65,67,70,74,77,80} following 3wCBT. Compared with no/minimal intervention, three studies^{40,44,67} showed a greater decrease in disinhibition favouring 3wCBT, one study⁴⁰ reported a greater decrease in binge eating and hunger favouring 3wCBT, and two studies^{44,67} showed a greater decrease in emotional eating favouring 3wCBT. Pairwise estimates comparing 3wCBT and SBT found a greater decrease in disinhibition for 3wCBT ($N = 7$)^{35,40,41,46,69,70,73} but no evidence of a difference between the groups in binge eating ($N = 4$)^{33,40,45,74} hunger ($N = 4$)^{35,40,46,69} or emotional eating ($N = 5$)^{31,35,41,70,74}. Pooled arm-specific estimates showed an increase in dietary restraint ($N = 12$)^{29,35,40,41,43,44,46,67,69,75,78,80} intuitive eating ($N = 2$)^{34,44} and mindful eating ($N = 6$)^{36,38,64,65,74} following 3wCBT. Pairwise estimates comparing 3wCBT and no/minimal intervention showed a greater increase in dietary restraint ($N = 3$)^{40,44,67} and a greater increase in intuitive eating ($N = 1$)⁴⁴ in the 3wCBT group. Pairwise estimates comparing 3wCBT and SBT found a greater increase in

mindful eating for 3wCBT ($N = 2$)^{36,74} but no evidence of a difference in dietary restraint ($N = 5$)^{35,40,41,46,69}. No studies compared intuitive eating in 3wCBT versus SBT.

Pooled estimates ($N = 6$)^{33,35,43,65,78} showed no evidence of a change in body dissatisfaction following 3wCBT; pairwise comparisons showed no evidence of a difference between 3wCBT and SBT ($N = 2$)^{33,35}.

3.4.3 | Physical health outcomes

Pooled arm-specific estimates ($N = 4$)^{30,32,36,68} suggested a reduction in diastolic blood pressure and systolic blood pressure following 3wCBT, but pairwise estimates showed no evidence of differences between 3wCBT and SBT. Pooled arm-specific estimates ($N = 3$)^{32,68,69} suggested a reduction in fasting glucose and HbA_{1c} following 3wCBT, and pairwise estimates suggested greater reductions in fasting glucose and HbA_{1c} for 3wCBT versus SBT. There was no evidence of changes in high-density lipoprotein cholesterol ($N = 1$)⁶⁸ low-density lipoprotein cholesterol ($N = 2$)^{32,68} or total cholesterol ($N = 1$) following 3wCBT,⁷⁰ and no evidence of differences between 3wCBT and SBT in any of these outcomes. One study⁶⁸ reported a decrease in triglyceride and triglyceride-to-HDL ratio following 3wCBT and a greater decrease of triglyceride in 3wCBT compared with SBT with no evidence of a change for triglyceride-to-HDL ratio between 3wCBT and SBT. Pooled arm-specific estimates ($N = 6$)^{36,42,68-70,76} showed a decrease in waist




FIGURE 4 Summary of weight change from network meta-analysis at different follow-up from baseline time points. Time points are months since baseline unless otherwise specified. ACT, acceptance and commitment therapy; CFT, compassion-focused therapy; MBCT, mindfulness-based cognitive behaviour therapy; No/min, no/minimal intervention; SBT, standard behavioural treatment

circumference following 3wCBT, but pairwise comparisons found no evidence of a difference between 3wCBT and SBT.

3.5 | Meta-regression of intervention characteristics

A sufficient number of studies for meta-regression ($N \geq 10$) were only available at post-intervention and at 3- and 6- months since baseline for 3wCBTs versus SBT. Prespecified study and intervention characteristics were examined in the meta-regression at these time points including number of sessions, duration of intervention, method of delivery, and risk of bias (Table S7), and none were found to have any impact on the effect estimates on weight or BMI reported in the pairwise meta-analysis. There were too few studies in each stratum to analyse the potential effects of comorbidities (eg, diabetes). Due to the small number of studies, subgroup analysis was not conducted.

3.6 | Intervention adherence

There was substantial heterogeneity and poor reporting of attendance and adherence outcomes, limiting our ability to conduct any

meaningful quantitative analysis (Table S8). Only 22 studies reported any attendance information, but, for all these studies, attendance was at least 60% at group sessions overall, and eight^{31,35,42,46,54,69,71,72} out of 11 RCTs reporting attendance information for each group had a 3wCBT group with higher attendance than the control arm. Information provided on adherence included minutes of home meditation practice, number of mindful meals per week, food and exercise diaries, and completion of online modules. Generally, within each study, there seemed to be a spread of engagement in the home practice aspect of interventions. This also varied with interventions delivered via internet: one study⁴⁴ found a 91% median completion of all modules, and another³⁴ found a mean of 32%.

4 | DISCUSSION

This comprehensive systematic review and network meta-analysis found high-quality evidence suggesting that 3wCBT results in greater weight loss than no/minimal intervention. Importantly, it also found moderate-quality evidence that suggests that 3wCBT results in greater weight loss than SBT at post-intervention and high-quality evidence from a small number of studies indicating that 3wCBT

(A) Psychological outcomes




FIGURE 5 Changes in secondary outcomes comparing third-wave cognitive behaviour therapy and standard behavioural treatment at earliest time point post-intervention using random-effects pairwise meta-analysis. A, Psychological outcomes. B, Eating behaviours. C, Physical health outcomes. For variables with asterisks (*), estimates to the right of the dotted line indicate a desired change in favour of third-wave CBTs; for all other variables, it is to the left of the dotted line. ACT, acceptance and commitment therapy; CBT, cognitive behaviour therapy; MBCT, mindfulness-based cognitive behaviour therapy; SMD, standardized mean difference

results in greater weight loss than SBT at longer term follow-up. However, it is important to note that 3wCBTs did not consistently outperform other interventions across shorter follow-up times (eg, 3- and 9- months) and that differences in weight change between 3wCBT and SBT were small (approximately 0.6 kg difference post-intervention and 1.4 kg difference at 24 -months). Future research is

needed to establish the clinical significance of these small differences in weight change.

The finding that 3wCBT is potentially more effective than SBT contrasts with the report by Roche et al¹⁴ that acceptance- and mindfulness-based interventions were only more effective than waitlist control arms. Conversely, estimates for the difference in

(B) Eating behaviours




FIGURE 5 (Continued)

weight loss between CBT and no/minimal intervention in our analysis are slightly smaller. This may be because Roche et al¹⁴ merged short- and long-term follow-up data and included participants with a BMI less than 25kg/m², either of which may be associated with larger effects. Comparisons with other reviews that purport to have compared 3wCBT with other approaches are more challenging because of

less appropriate statistical approaches. Both Carrière et al¹⁵ and Rogers et al¹³ combined the estimates from both RCTs and single-arm pre-intervention to post-intervention studies. While the analytic methods used in Carrière et al¹⁵ is unclear, Rogers et al¹³ used only the post-intervention estimates for the RCTs (which ignores baseline differences between groups that may be influential in smaller

(C) Physical health outcomes




FIGURE 5 (Continued)

studies),⁸² and effect estimates were weighted by sample size, not SD. Consequently, the reported study-specific estimates are different

between Rogers et al¹³ and Roche et al¹⁴ even though both label the effect as Hedges' g.^{15,59}

Our statistical approach was more rigorous. In the pairwise meta-analysis, we only included RCTs, we applied a consistent definition of 'change' estimate as pre-intervention minus post-intervention estimates, we used appropriate methods for pairwise comparisons, and we further separated the effectiveness against no/minimal control or SBTs. We reported the effect estimates by follow-up times to have more insights of the short-term and longer term effects, and we restricted our analytic population to those with a BMI greater than or equal to 25 kg/m² as this is more relevant from a clinical and policy perspective.

To our knowledge, this is also the first review to report on the comparative effectiveness of individual 3wCBT types, which have never been tested in head-to-head trials. Network meta-analysis found that ACT-based interventions had the most consistent evidence of effectiveness indicating greater weight loss compared with SBT at post-intervention and 12- and 24-month follow-up from baseline; comparisons at other time points or with other 3wCBT types either appeared to favour ACT or did not show evidence of a difference. ACT was ranked as the best intervention up to 12 months and was the only 3wCBT to have outcomes at 24 and 36 months. Network estimates suggested that MBCT resulted in greater weight loss than SBT at 12 and 18 months, but favoured SBT at 9 months, and there was no evidence that MBCT was more effective than no/minimal intervention. This suggests that we should interpret the finding that MBCT was the highest ranking intervention at 12 and 18 months with some caution. Only four studies evaluated a 3wCBT approach other than acceptance or mindfulness and these were of low quality and short follow-up. Although we identified three studies using DBT, all used non-randomized pre-intervention to post-intervention design and one was combined with pharmacotherapy; therefore, they were not included in the pairwise or network meta-analysis, limiting conclusions on DBT effectiveness. CFT was found to have no evidence for weight loss; however, this finding is based upon one unpublished thesis,⁴¹ which was deemed to be of high risk of bias and of very low quality. To date, the evidence provides strongest support for the superiority of acceptance-based interventions. It is possible that the superiority of the acceptance-based programmes in this context is due to its focus on values and willingness to reduce experiential avoidance. However, more research is needed to confirm these differences and identify the mechanisms of action.

Changes in secondary outcomes were generally in the desired direction. Following 3wCBT, there was evidence suggestive of reductions in depression, anxiety, and stress and increases in quality of life, mindfulness, and psychological flexibility. There were similar reductions in binge eating, dietary disinhibition, hunger, and emotional eating and increases in dietary restraint, intuitive eating, and mindful eating following 3wCBT. This is consistent with previous reviews that have reported on these outcomes.^{13,15,16} Pairwise comparisons suggest that most improvements in these outcomes were greater than for no/minimal intervention. Focusing on comparisons of 3wCBT versus SBT, pairwise comparisons showed that reductions in disinhibition and increases in mindful eating were greater for 3wCBT but no differences in other psychological factors. These could represent important mechanisms by which 3wCBT has a greater effect on

weight control than SBT and warrant further investigation. Few studies reported changes in blood pressure, glycaemia, lipid profile, and waist circumference, and pairwise comparisons only suggested greater reductions in glycaemia.

Meta-regression did not identify any specific intervention characteristics (eg, duration, mode of delivery, number of sessions) that were more effective than others. This may have been due to the small number of studies. Similar to traditional behavioural weight management programmes, the majority of interventions were delivered in a group face-to-face format. Such delivery has often been found to be effective in weight loss,⁸³ with the group providing peer support and regular encouragement, particularly for those experiencing social isolation. However, closed-group programmes led by clinical psychologists can be logistically difficult and costly to deliver, particularly in the context of national health services providing free or discounted health care. Increasingly, standard behavioural weight management interventions are moving to more scalable methods of delivery to increase reach and reduce cost. Only four interventions^{34,44,74,79} in our review used remote delivery through internet or mobile phone. Similarly, a review of online mindful eating interventions¹⁸ found only two studies focusing on weight loss, and none were delivered through mobile telephone, highlighting a dearth of investigation into this research area. Greater consideration may need to be given to the use of digital technology to facilitate intervention delivery, with its 24-hour accessibility, improved scalability, and increased reach.⁸⁴

5 | STRENGTHS AND LIMITATIONS

The study of 3wCBT for weight management is in its relative infancy, with the earliest included RCTs from 2008³³ (MBCT) and 2011³⁰ (ACT). While a number of RCTs have been conducted, most have had small samples and short follow-up and few provide high-quality evidence. Only three RCTs reported outcomes beyond 12 months despite 3wCBT being hypothesized to have particular benefit for long-term outcomes. However, these studies with longer term follow-up had low risk of bias and provided high-quality evidence. The small number of studies limited our network meta-analysis to up to 18 months post-baseline and meant that there were insufficient studies to conduct meaningful meta-regression on the most effective intervention components and characteristics beyond 6 months. Many studies also had very small sample sizes. The small number of studies and small sample sizes meant that many of our estimates had wide CIs, thereby limiting the power to detect a difference. Many of the studies included in this review had high or serious risk of bias. However, it should be noted that we used a stringent assessment tool, and heterogeneity for many outcomes was low. For some studies, this may also reflect the slowness of the obesity field to adopt standards for trial reporting (eg, CONSORT),²³ rather than the quality of the research itself. It is also important to note that the studies with longer term follow-up (ie, 12 and 24 months from baseline) were of high quality, based upon the GRADE assessment tool,²⁷ so we can have greater confidence in the findings at these time points.

There was heterogeneity in the content of 3wCBT programmes, with a combination of standardized, modified, and novel programmes that varied in length and practice time. Some studies evaluated interventions that used combinations of different third-wave therapies, which may obscure potential differences between types of therapy. However, this is a reflection of how these interventions are used. In attempting to collate adherence and attendance data, we found a low number of studies reporting intervention fidelity information and substantial variability in reporting. 3wCBTs seem to have comparable attendance and attrition rates to standard behavioural programmes,⁸⁵ suggesting that they are an acceptable and feasible option. Lack of information stopped us from conducting a meta-regression to try to identify sources of heterogeneity in attendance or adherence.

Studies in our review had a high proportion of female participants; this is typical in weight loss programmes and mindfulness interventions.⁸⁶ This limits generalizability of findings to males⁸³ and warrants purposive recruitment of males to studies and weight loss programmes per se. Furthermore, several studies lacked detail of participant demographics such as ethnicity and socio-economic status; more complete reporting would enable us to understand the extent of the generalizability of results.

There are several strengths to this review. RCTs and pre-intervention to post-intervention studies were included in our pooled estimates, but only RCTs were included in pairwise and network meta-analysis to provide the estimates of the comparative effectiveness against a comparator. By conducting a network analysis, we could estimate comparisons between different types of 3wCBT that have not been directly compared, incorporating direct, indirect, and mixed evidence in our evaluations of the evidence. To maximize on relevant research, we included unpublished theses and contacted authors regarding abstracts in conference proceedings. Unlike previous reviews,¹⁴ we restricted our analytic population to those with a BMI ≥ 25 kg/m² to make our results more relevant to health care policies that recommend weight management interventions for people with overweight/obesity.

6 | CONCLUSIONS

This systematic review and network meta-analysis found moderate-quality evidence suggesting that 3wCBT results in a small increase in weight loss compared with SBT at post-intervention. It found high-quality evidence from a small number of studies suggesting that 3wCBT results in greater weight loss than SBT at 12 and 24 month follow-up from baseline. Evidence specifically appears to support the use of acceptance-based programmes. Larger, high-quality trials are needed in this area to better understand who these interventions work for and how they work, so that we can target these interventions appropriately and identify the most crucial components and "active ingredients." Future research should also consider how we deliver these interventions in a cost-effective way that maximizes scalability while maintaining effectiveness.

ACKNOWLEDGEMENTS

This study is funded by the National Institute for Health Research (NIHR) Programme Grants for Applied Research RP-PG-0216-20010. The views expressed are those of the authors and not necessarily those of the NIHR or the Department of Health and Social Care. ALA and SJG are supported by the Medical Research Council (MC_UU_12015/4). SJG is an NIHR senior investigator. The University of Cambridge has received salary support in respect of SJG from the National Health Service in the East of England through the Clinical Academic Reserve. We want to thank the patient user group panel for assisting with the refinement of the research question and interpretation of results. We would like to thank individuals within the unit that assisted with article language translation, Eleanor Barker for help in developing the search strategy and Rebecca Jones for assisting with database searches. We would also like to thank all corresponding authors of articles that provided additional information or clarity on their studies.

CONFLICTS OF INTEREST

AA is the chief investigator on two publically funded (MRC, NIHR) trials where the intervention is provided by WW (formerly Weight Watchers) at no cost outside the submitted work. AJH reports receiving personal fees from Slimming World, outside the submitted work. CAH reports education work and consultancy Oviva, Orexigen Therapeutics, Kasteck, Ethicon, Mundipharma, Consilient Health, Nestle, and Novo Nordisk, outside the submitted work. ERL, NI, SB, and SJG have no conflicts of interest.

ORCID

Emma R. Lawlor  <https://orcid.org/0000-0002-0742-0476>

Nazrul Islam  <https://orcid.org/0000-0003-3982-4325>

Sarah Bates  <https://orcid.org/0000-0002-6358-0060>

Amy L. Ahern  <https://orcid.org/0000-0001-5069-4758>

REFERENCES

- Curioni CC, Lourenc PM. Long-term weight loss after diet and exercise: a systematic review. *Int J Obes*. 2005;29(10):1168-1174. <https://doi.org/10.1038/sj.ijo.0803015>
- Forman EM, Butryn ML, Manasse M, Bradley LE. Acceptance-based behavioral treatment for weight control: a review and future directions. *Curr Opin Psychol*. 2015;2:87-90. <https://doi.org/10.1016/j.copsyc.2014.12.020>
- Hartmann-Boyce J, Johns DJ, Jebb SA, Summerbell C, Aveyard P. Behavioural weight management programmes for adults assessed by trials conducted in everyday contexts: systematic review and meta-analysis. *Obes Rev*. 2014;15(11):920-932. <https://doi.org/10.1111/obr.12220>
- Barte JCM, Ter Bogt NCW, Bogers RP, et al. Maintenance of weight loss after lifestyle interventions for overweight and obesity, a systematic review. *Obes Rev*. 2010;11(12):899-906. <https://doi.org/10.1111/j.1467-789X.2010.00740.x>
- Dombrowski S, Knittle K, Avenell A, Araujo-Soares V, Snihotta F. Long term maintenance of weight loss with non-surgical interventions in obese adults: systematic review and meta-analyses of randomised controlled trials. *BMJ*. 2014;348(6):g2646. <https://doi.org/10.1136/bmj.g2646>

6. Dietz WH, Baur LA, Hall K, et al. Management of obesity: improvement of health-care training and systems for prevention and care. *Lancet*. 2015;385(9986):2521-2533. [https://doi.org/10.1016/S0140-6736\(14\)61748-7](https://doi.org/10.1016/S0140-6736(14)61748-7)
7. NICE guideline. Obesity: identification, assessment and management. Clinical guideline [CG189]. 2014.
8. Hayes SC, Villatte M, Levin M, Hildebrandt M. Open, aware, and active: contextual approaches as an emerging trend in the behavioral and cognitive therapies. *Annu Rev Clin Psychol*. 2011;7:141-168. <https://doi.org/10.1146/annurev-clinpsy-032210-104449>
9. Hayes SC. Acceptance and commitment therapy, relational frame theory, and the third wave of behavioral and cognitive therapies. *Behav Ther*. 2004;35(4):639-665. [https://doi.org/10.1016/S0005-7894\(04\)80013-3](https://doi.org/10.1016/S0005-7894(04)80013-3)
10. Linardon J, Fairburn CG, Fitzsimmons-Craft EE, Wilfley DE, Brennan L. The empirical status of the third-wave behaviour therapies for the treatment of eating disorders: a systematic review. *Clin Psychol Rev*. 2017;58:125-140. <https://doi.org/10.1016/j.cpr.2017.10.005>
11. Olson KL, Emery CF. Mindfulness and weight loss: a systematic review. *Psychosom Med*. 2015;77(1):59-67. <https://doi.org/10.1097/PSY.0000000000000127>
12. O'Reilly GA, Cook L, Spruijt-Metz D, Black DS. Mindfulness-based interventions for obesity-related eating behaviours: a literature review. *Obes Rev*. 2014;15(6):453-461. <https://doi.org/10.1111/obr.12156>
13. Rogers JM, Ferrari M, Mosely K, Lang CP, Brennan L. Mindfulness-based interventions for adults who are overweight or obese: a meta-analysis of physical and psychological health outcomes. *Obes Rev*. 2017;18(1):51-67. <https://doi.org/10.1111/obr.12461>
14. Roche AI, Kroska EB, Denburg NL. Acceptance- and mindfulness-based interventions for health behavior change: Systematic review and meta-analyses. *J Contextual Behav Sci*. 2019;13:74-93. <https://doi.org/10.1016/j.jcbs.2019.06.002>
15. Carrière K, Khoury B, Günak MM, Knäuper B. Mindfulness-based interventions for weight loss: a systematic review and meta-analysis. *Obes Rev*. 2018;19(2):164-177. <https://doi.org/10.1111/obr.12623>
16. Ruffault A, Czernichow S, Hagger MS, et al. The effects of mindfulness training on weight-loss and health-related behaviours in adults with overweight and obesity: a systematic review and meta-analysis. *Obes Res Clin Pract*. 2017;11(5 Suppl 1):90-111. <https://doi.org/10.1016/j.orcp.2016.09.002>
17. Katterman SN, Kleinman BM, Hood MM, Nackers LM, Corsica JA. Mindfulness meditation as an intervention for binge eating, emotional eating, and weight loss: a systematic review. *Eat Behav*. 2014;15(2):197-204. <https://doi.org/10.1016/j.eatbeh.2014.01.005>
18. Lyzwinski LN, Caffery L, Bambling M, Edirippulige S. A systematic review of electronic mindfulness-based therapeutic interventions for weight, weight-related behaviors, and psychological stress. *Telemed e-Health*. 2018;24(3):173-184. <https://doi.org/10.1089/tmj.2017.0117>
19. Public Health England. National Mapping of Weight Management Services; Provision of Tier 2 and Tier 3 Services in England. 2015.
20. Curry SJ, Krist AH, Owens DK, et al. Behavioral weight loss interventions to prevent obesity-related morbidity and mortality in adults US preventive services task force recommendation statement. *JAMA*. 2018;320(11):1163-1171. <https://doi.org/10.1001/jama.2018.13022>
21. Lawlor ER, Islam N, Griffin SJ, Hill AJ, Hughes CA, Ahern AL. Third-wave cognitive behaviour therapies for weight management: systematic review and network meta-analysis protocol. *BMJ Open*. 2018;8(7):e023425. <https://doi.org/10.1136/bmjopen-2018-023425>
22. The Cochrane Public Health Group. Cochrane Public Health Group Data Extraction and Assessment Template. 2011:1-22.
23. Schulz KF, Altman DG, Moher D, CONSORT Group. CONSORT 2010 Statement: updated guidelines for reporting parallel group randomised trials. *BMC Med*. 2010;8:18. <https://doi.org/10.1186/1741-7015-8-18>
24. Hoffmann TC, Glasziou PP, Boutron I, et al. Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide. *BMJ*. 2014;348:g1687. <https://doi.org/10.1136/bmj.g1687>
25. Higgins JPT, Sterne JAC, Savović J, et al. A revised tool for assessing risk of bias in randomized trials. In: Chandler J, McKenzie J, Boutron I, Welch V, eds. *Cochrane Database of Systematic Reviews*. Cochrane Methods; 2016 Issue 10 (Suppl 1). <https://doi.org/10.1002/14651858.CD201601>
26. Sterne JA, Hernán MA, Reeves BC, et al. ROBINS-I: A tool for assessing risk of bias in non-randomised studies of interventions. *BMJ*. 2016;355:i4919. <https://doi.org/10.1136/bmj.i4919>
27. Guyatt GH, Oxman A, Vist G, et al. GRADE an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ*. 2008;336(7650):924-926. <https://doi.org/10.1136/bmj.39489.470347.AD>
28. Braun TD, Park CL, Conboy LA. Psychological well-being, health behaviors, and weight loss among participants in a residential Kripalu yoga-based weight loss program. *Int J Yoga Therap*. 2012;22:9-22.
29. Dalen J, Smith BW, Shelley BM, Sloan AL, Leahigh L, Begay D. Pilot study: Mindful Eating and Living (MEAL): Weight, eating behavior, and psychological outcomes associated with a mindfulness-based intervention for people with obesity. *Complement Ther Med*. 2010;18(6):260-264. <https://doi.org/10.1016/j.ctim.2010.09.008>
30. Fletcher LB. A Mindfulness and Acceptance-based Intervention for Increasing Physical Activity and Reducing Obesity. *Dissertation*. 2011.
31. Goldbacher E, La Grotte C, Komaroff E, Vander Veur S, Foster GD. An initial evaluation of a weight loss intervention for individuals who engage in emotional eating. *J Behav Med*. 2016;39(1):139-150. <https://doi.org/10.1007/s10865-015-9678-6>
32. Raja-Khan N, Agito K, Shah J, et al. Mindfulness-based stress reduction in women with overweight or obesity: a randomized clinical trial. *Obesity*. 2017;25(8):1349-1359. <https://doi.org/10.1002/oby.21910>
33. Blevins NC. Mindfulness Meditation as an Intervention for Body Image and Weight Management in College Women: A Pilot Study. *Dissertation*. 2008.
34. Boucher S, Gray A, Lillis J, Tyka TL, Horwath CC. Teaching intuitive eating and acceptance and commitment therapy skills via a web-based intervention: a pilot single-arm intervention study. *JMIR Res Protoc*. 2016;5(4):e180. <https://doi.org/10.2196/resprot.5861>
35. Davis KK. Effect of Mindfulness Meditation and Home-Based Resistance Exercise on Weight Loss, Weight Loss Behaviors, and Psychosocial Correlates in Overweight Adults. *Dissertation*. 2008.
36. Lee TM. Comparing Mindfulness-Enriched Weight Management to Current Standard Practices. *Dissertation*. 2017. <https://doi.org/10.13023/ETD.2017.054>
37. Palmeira L, Cunha M, Pinto-Gouveia J. Processes of change in quality of life, weight self-stigma, body mass index and emotional eating after an acceptance-, mindfulness- and compassion-based group intervention (Kg-Free) for women with overweight and obesity. *J Health Psychol*. 2017;24(8):1-14. <https://doi.org/10.1177/1359105316686668>
38. Chung S, Zhu S, Friedmann E, et al. Weight loss with mindful eating in African American women following treatment for breast cancer: a longitudinal study. *Support Care Cancer*. 2016;24(4):1875-1881. <https://doi.org/10.1007/s00520-015-2984-2>
39. Gallé F, Cirella A, Salzano AM, Di Onofrio V, Belfiore P, Liguori G. Analyzing the effects of psychotherapy on weight loss after laparoscopic gastric bypass or laparoscopic adjustable gastric banding in patients with borderline personality disorder: a prospective study. *Scand J Surg*. 2017;106(4):299-304. <https://doi.org/10.1177/1457496917701670>

40. Kristeller J, Wolever RQ, Sheets V. Mindfulness-based eating awareness training (MB-EAT) for binge eating: a randomized clinical trial. *Mind*. 2014;5(3):282-297. <https://doi.org/10.1007/s12671-012-0179-1>
41. Loader KA. A compassionate mind approach to self-help for treatment seeking obese adults: a randomised controlled trial. *Dissertation*. 2013.
42. McKee HC, Ntoumanis N. Developing self-regulation for dietary temptations: intervention effects on physical, self-regulatory and psychological outcomes. *J Behav Med*. 2014;37(6):1075-1081. <https://doi.org/10.1007/s10865-014-9557-6>
43. Roosen MA, Safer D, Adler S, Cebolla A, van Strien T. Group dialectical behavior therapy adapted for obese emotional eaters: a pilot study. *Nutr Hosp*. 2012;27(4):1141-1147. <https://doi.org/10.3305/nh.2012.27.4.5843>
44. Sairanen E, Tolvanen A, Karhunen L, et al. Psychological flexibility mediates change in intuitive eating regulation in acceptance and commitment therapy interventions. *Public Health Nutr*. 2017;20(9):1681-1691. <https://doi.org/10.1017/s1368980017000441>
45. Smith BW, Shelley BM, Sloan AL, Colleran K, Erickson K. A preliminary randomized controlled trial of a mindful eating intervention for post-menopausal obese women. *Mind*. 2018;9(3):836-849. <https://doi.org/10.1007/s12671-017-0824-9>
46. Spadaro KC, Davis KK, Sereika SM, Gibbs BB, Jakicic JM, Cohen SM. Effect of mindfulness meditation on short-term weight loss and eating behaviors in overweight and obese adults: a randomized controlled trial. *J Complement Integr Med*. 2017;15(2). <https://doi.org/10.1515/jcim-2016-0048>
47. Godfrey KM, Gallo LC, Afari N. Mindfulness-based interventions for binge eating: a systematic review and meta-analysis. *J Behav Med*. 2015;38(2):348-362. <https://doi.org/10.1007/s10865-014-9610-5>
48. Vocks S, Tuschen-Caffier B, Pietrowsky R, Rustenbach SJ, Kersting A, Herpertz S. Meta-analysis of the effectiveness of psychological and pharmacological treatments for binge eating disorder. *Int J Eat Disord*. 2010;43(3):205-217. <https://doi.org/10.1002/eat.20696>
49. StataCorp. *Stata Statistical Software*. 2015.
50. Wasserstein RL, Lazar NA. The ASA's statement on *p*-values: context, process, and purpose. *Am Stat*. 2016;70(2):129-133. <https://doi.org/10.1080/00031305.2016.1154108>
51. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials*. 1986;7(3):177-188. [https://doi.org/10.1016/0197-2456\(86\)90046-2](https://doi.org/10.1016/0197-2456(86)90046-2)
52. Hedges LV, Olkin I. *Statistical Methods for Meta-Analysis*. Academic Press; 2014.
53. Salanti G, Del Giovane C, Chaimani A, Caldwell DM, Higgins JPT. Evaluating the quality of evidence from a network meta-analysis. *PLoS ONE*. 2014;9(7):e99682. <https://doi.org/10.1371/journal.pone.0099682>
54. Butryn ML, Forman EM, Lowe MR, Gorin AA, Zhang F, Schaumberg K. Efficacy of environmental and acceptance-based enhancements to behavioral weight loss treatment: the ENACT trial. *Obesity*. 2017;25(5):866-872. <https://doi.org/10.1002/oby.21813>
55. Higgins JPT, Jackson D, Barrett JK, Lu G, Ades AE, White IR. Consistency and inconsistency in network meta-analysis: concepts and models for multi-arm studies. *Res Synth Methods*. 2012;3(2):98-110. <https://doi.org/10.1002/jrsm.1044>
56. Dias S, Welton NJ, Caldwell DM, Ades AE. Checking consistency in mixed treatment comparison meta-analysis. *Stat Med*. 2010;29(7-8):932-944. <https://doi.org/10.1002/sim.3767>
57. Salanti G, Ades AE, Ioannidis JPA. Graphical methods and numerical summaries for presenting results from multiple-treatment meta-analysis: an overview and tutorial. *J Clin Epidemiol*. 2011;64(2):163-171. <https://doi.org/10.1016/j.jclinepi.2010.03.016>
58. Tobias A. Assessing the influence of a single study in the meta-analysis estimate. *Stata Tech Bull*. 1999;47:15-17. <https://www.stata-press.com/journals/stbcontents/stb47.pdf>
59. Higgins JPT, Green S (Eds). *Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 (Updated March 2011)*. The Cochrane Collaboration; 2011 www.handbook.cochrane.org.
60. Lappalainen R, Sairanen E, Järvelä E, et al. The effectiveness and applicability of different lifestyle interventions for enhancing wellbeing: the study design for a randomized controlled trial for persons with metabolic syndrome risk factors and psychological distress. *BMC Public Health*. 2014;14:310. <https://doi.org/10.1186/1471-2458-14-310>
61. Raja-Khan N, Agito K, Shah J, et al. Mindfulness-based stress reduction for overweight/obese women with and without polycystic ovary syndrome: design and methods of a pilot randomized controlled trial. *Contemp Clin Trials*. 2015;41:287-297. <https://doi.org/10.1016/j.cct.2015.01.021>
62. Lillis J, Niemeier HM, Ross KM, et al. Weight loss intervention for individuals with high internal disinhibition: design of the Acceptance Based Behavioral Intervention (ABBI) randomized controlled trial. *BMC Psychol*. 2015;3(1):17. <https://doi.org/10.1186/s40359-015-0075-2>
63. Mattila E, Lappalainen R, Välikynen P, Sairanen E, Lappalainen P. Usage and dose response of a mobile acceptance and commitment therapy app: secondary analysis of the intervention arm of a randomized controlled trial. *JMIR Mhealth Uhealth*. 2016;4(3):e90. <https://doi.org/10.2196/mhealth.5241>
64. Hamel REG. Weight loss and psychosocial effects of a mindfulness-based intervention in overweight adults: a pilot study. *Dissertation*. 2010.
65. Braun TD, Noggle JJ, Gorin AA, Garivaltis H, Conboy LA, Park CL. Group-based yogic weight loss with Ayurveda-inspired components: a pilot investigation of female yoga practitioners and novices. *Int J Yoga Therap*. 2016;26:55-72. <https://doi.org/10.17761/1531-2054-26.1.55>
66. Adler S. Orlistat/Alli compared to Orlistat/Alli plus dialectical behavior therapy in overweight binge eaters: a randomized control trial. *Dissertation*. 2008.
67. Daubenmier J, Kristeller J, Hecht FM, et al. Mindfulness intervention for stress eating to reduce cortisol and abdominal fat among overweight and obese women: an exploratory randomized controlled study. *J Obes*. 2011;2011:1-13. <https://doi.org/10.1155/2011/651936>
68. Daubenmier J, Moran PJ, Kristeller J, et al. Effects of a mindfulness-based weight loss intervention in adults with obesity: a randomized clinical trial. *Obesity*. 2016;24(4):794-804. <https://doi.org/10.1002/oby.21396>
69. Miller CK, Kristeller JL, Headings A, Nagaraja H, Miser WF. Comparative effectiveness of a mindful eating intervention to a diabetes self-management intervention among adults with type 2 diabetes: a pilot study. *J Acad Nutr Diet*. 2012;112(11):1835-1842. <https://doi.org/10.1016/j.jand.2012.07.036>
70. Palmeira L, Pinto-Gouveia J, Cunha M. Exploring the efficacy of an acceptance, mindfulness & compassionate-based group intervention for women struggling with their weight (Kg-Free): a randomized controlled trial. *Appetite*. 2017;112:107-116. <https://doi.org/10.1016/j.appet.2017.01.027>
71. Forman EM, Butryn ML, Juarascio AS, et al. The Mind Your Health project: a randomized controlled trial of an innovative behavioral treatment for obesity. *Obesity*. 2013;21(6):1119-1126. <https://doi.org/10.1002/oby.20169>
72. Forman EM, Butryn ML, Manasse SM, et al. Acceptance-based versus standard behavioral treatment for obesity: results from the Mind Your Health randomized controlled trial. *Obesity*. 2016;24(10):2050-2056. <https://doi.org/10.1002/oby.21601>
73. Lillis J, Niemeier HM, Thomas JG, et al. A randomized trial of an acceptance-based behavioral intervention for weight loss in people with high internal disinhibition. *Obesity*. 2016;24(12):2509-2514. <https://doi.org/10.1002/oby.21680>

74. Carpenter KM, Vickerman KA, Salmon EE, Javitz HS, Epel ES, Lovejoy JC. A randomized pilot study of a phone-based mindfulness and weight loss program. *Behav Med*. 2017;6(4):1-11. <https://doi.org/10.1080/08964289.2017.1384359>
75. Niemeier HM, Leahey T, Palm Reed K, Brown RA, Wing RR. An acceptance-based behavioral intervention for weight loss: a pilot study. *Behav Ther*. 2012;43(2):427-435. <https://doi.org/10.1016/j.beth.2011.10.005>
76. Andalcio AA. Implementation of a Lifestyle Intervention Program in the Primary Care Setting to Decrease Body Mass Indices in Obese Adults. *Dissertation*. 2018.
77. Hanson P, Shuttlewood E, Halder L, et al. Application of mindfulness in a tier 3 obesity service improves eating behavior and facilitates successful weight loss. *J Clin Endocrinol Metab*. 2019;104(3):793-800. <https://doi.org/10.1210/je.2018-00578>
78. Lundgren JD. A Mindfulness-Based Behavioral Treatment for Weight Loss. *Dissertation*. 2003.
79. Bradley LE, Forman EM, Kerrigan SG, et al. Project HELP: a remotely delivered behavioral intervention for weight regain after bariatric surgery. *Obes Surg*. 2017;27(3):586-598. <https://doi.org/10.1007/s11695-016-2337-3>
80. Forman EM, Butryn ML, Hoffman KL, Herbert JD. An open trial of an acceptance-based behavioral intervention for weight loss. *Cogn Behav Pract*. 2009;16(2):223-235. <https://doi.org/10.1016/j.cbpra.2008.09.005>
81. Butryn ML, Phelan S, Hill JO, Wing RR. Consistent self-monitoring of weight: a key component of successful weight loss maintenance. *Obesity*. 2007;15(12):3091-3096. <https://doi.org/10.1038/oby.2007.368>
82. Johnsen TJ, Friborg O. The effects of cognitive behavioral therapy as an anti-depressant treatment is falling: a meta-analysis. *Psychol Bull*. 2015;141(4):747-768. <https://doi.org/10.1037/bul0000015>
83. Borek AJ, Abraham C, Greaves CJ, Tarrant M. Group-based diet and physical activity weight-loss interventions: a systematic review and meta-analysis of randomised controlled trials. *Appl Psychol Health Well-Being*. 2018;10(1):62-86. <https://doi.org/10.1111/aphw.12121>
84. Beleigoli AM, Andrade AQ, Cançado AG, Paulo MN, Maria De Fátima HD, Ribeiro AL. Web-based digital health interventions for weight loss and lifestyle habit changes in overweight and obese adults: systematic review and meta-analysis. *J Med Internet Res*. 2019;8(21):e298. <https://doi.org/10.2196/jmir.9609>
85. Caplan W, Bowman JD, Pronk NP. Weight-loss outcomes: a systematic review and meta-analysis of weight-loss clinical trials with a minimum 1-year follow-up. *J Am Diet Assoc*. 2007;107(10):1755-1767. <https://doi.org/10.1016/j.jada.2007.07.017>
86. Pagoto SL, Schneider KL, Oleski JL, Luciani JM, Bodenlos JS, Whited MC. Male inclusion in randomized controlled trials of lifestyle weight loss interventions. *Obesity*. 2012;20(6):1234-1239. <https://doi.org/10.1038/oby.2011.140>
87. Daubenmier J, Lin J, Blackburn E, et al. Changes in stress, eating, and metabolic factors are related to changes in telomerase activity in a randomized mindfulness intervention pilot study. *Psychoneuroendocrinology*. 2012;37(7):917-928. <https://doi.org/10.1016/j.psyneuen.2011.10.008>
88. Mason AE, Epel ES, Aschbacher K, et al. Reduced reward-driven eating accounts for the impact of a mindfulness-based diet and exercise intervention on weight loss: data from the SHINE randomized controlled trial. *Appetite*. 2016;100:86-93. <https://doi.org/10.1016/j.appet.2016.02.009>
89. Miller CK, Kristeller JL, Headings A, Nagaraja H. Comparison of a mindful eating intervention to a diabetes self-management intervention among adults with type 2 diabetes: a randomized controlled trial. *Health Educ Behav*. 2014;41(2):145-154. <https://doi.org/10.1177/1090198113493092>
90. Spadaro KC. Weight loss: Exploring self-regulation through mindfulness meditation. *Dissertation*. 2008.
91. Schumacher LM, Godfrey KM, Forman EM, Butryn ML. Change in domain-specific but not general psychological flexibility relates to greater weight loss in acceptance-based behavioral treatment for obesity. *J Contextual Behav Sci*. 2019;12:59-65. <https://doi.org/10.1016/j.jcbs.2019.01.008>
92. Forman EM, Manasse SM, Butryn ML, Crosby RD, Dallal DH, Crochiere RJ. Long-term follow-up of the Mind Your Health project: acceptance-based versus standard behavioral treatment for obesity. *Obesity*. 2019;27(4):565-571. <https://doi.org/10.1002/oby.22412>
93. Lillis J, Thomas JG, Niemeier HM, Wing RR. Exploring process variables through which acceptance-based behavioral interventions may improve weight loss maintenance. *J Contextual Behav Sci*. 2017;6(4):398-403. <https://doi.org/10.1016/j.jcbs.2017.07.005>
94. Järvelä-Reijonen E, Karhunen L, Sairanen E, et al. The effects of acceptance and commitment therapy on eating behaviour and diet delivered through face-to-face contact and a mobile app: a randomized controlled trial. *Int J Behav Nutr Phys Act*. 2018;15(1):22. <https://doi.org/10.1186/s12966-018-0654-8>
95. Bradley LE. A remote, acceptance-based intervention for weight regain after bariatric surgery. *Dissertation*. 2015.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of this article.

How to cite this article: Lawlor ER, Islam N, Bates S, et al. Third-wave cognitive behaviour therapies for weight management: A systematic review and network meta-analysis. *Obesity Reviews*. 2020;21:e13013. <https://doi.org/10.1111/obr.13013>