Vertically-aligned Silicon Carbide Nanowires as Metal-free Photocatalyst under Visible Light

Jindui Hong,^{*a} Seyyed Shayan Meysami,^a Vitaliy Babenko,^{a,b} Santamon Luanwuthi,^a Jesus Acapulco,^a Philip Holdway,^a Nicole Grobert^{*a}

^a*Department of Materials, University of Oxford, Parks Road, Oxford OX1 3PH, United Kingdom*

^b*Present address: Centre of Advanced Photonics and Electronics, University of Cambridge, 9 JJ Thomson Ave, Cambridge CB3 0FA, United Kingdom*

^{*}*E-mail: jindui.hong@materials.ox.ac.uk; nicole.grobert@materials.ox.ac.uk*

Figure S1. SEM images of (a, b) VACNTs, (c, d) SiO₂-VACNTs, (e, f) VASiC/CNTs and (g, h) VASiCs showing the infiltration of SiO₂ gel in VACNTs and the presence of aligned nanowires in VASiC/CNTs and VASiCs.

Figure S2. TEM images of (a-c) VACNTs, (d-f) SiO₂-VACNTs, (g-i) VASiC/CNTs and (j-m) VASiCs showing the non-uniform and thicker diameter of VASiCs than VACNTs.

Figure S3. STEM image and EDS mapping for VACNTs, SiO₂-VACNTs and VASiC/CNTs showing the presence of SiO₂ in SiO₂-VACNTs and the presence of CNTs in VASiC/CNTs. C (orange), Fe (red), Si (green) and O (blue).

Figure S4. (a) Mass spectra of the gas residuals from the synthesis of VASiCs showing the main signal of Ar; (b) zoom-in view of the mass spectra showing signals of Ar and CO. m/z 20: $^{40}\text{Ar}^{2+}$; m/z 28: $[\text{CO}]^+$; m/z 36: $^{36}\text{Ar}^+$; m/z 40: $^{40}\text{Ar}^+$.