



Direct Imaging Discovery of Giant Exoplanet β Pictoris d: A Decade-long Game of Hide-and-seek

Ben J. Sutcliffe^{1,2,44} , Markus J. Bonse^{3,4,44} , Valentin Christiaens^{5,6,7} , Clémence Fontanive^{1,2} , Elisabeth C. Matthews^{8,9} , Luke T. Parker¹⁰ , Tim D. Pearce¹¹ , Jayne L. Birkby¹⁰ , Beth A. Biller^{1,2} , Trent J. Dupuy^{1,2} , Emily O. Garvin¹² , Leyla Iskandarli¹³ , Jens Kammerer³ , Yifan Zhou¹⁴ , Robert J. De Rosa¹⁵ , Aarynn L. Carter¹⁶ , Sasha Hinkley¹⁷ , Matthew A. Kenworthy¹⁸ , William O. Balmer¹⁹ , Iain Hammond⁸ , James Mang^{20,45} , Caroline V. Morley²⁰ , Mark J. Neeser³ , Olivier Absil^{7,46} , Anthony Boccaletti²¹ , Mariangela Bonavita¹ , Brendan P. Bowler²² , Xueqing Chen^{1,2} , Felix A. Dannert¹² , Julien H. Girard¹⁶ , Markus Kasper³ , Anne-Marie Lagrange^{21,23} , Pengyu Liu^{1,2} , Gilles Orban de Xivry⁷ , Michael Poon²⁴ , Sascha P. Quanz^{12,25} , Benoît Serra³ , Johanna M. Vos²⁶ , Kevin Wagner²⁷ , Jason Wang^{28,29} , Bernhard Schölkopf^{4,30} , Guido Agapito³¹ , Alex Agudo Berbel³² , Dániel Apai^{27,33,34} , Andrea Baruffolo³⁵ , Martin Black³⁶ , Marco Bonaglia³¹ , Runa Briguglio³¹ , Yixian Cao³² , Luca Carbonaro³¹ , Lee Chapman³⁶ , Giovanni Cresci³¹ , Yigit Dallilar³⁷ , Richard Davies³² , Matthias Deysenroth³² , Ivan Di Antonio³⁸ , Amico Di Cianno³⁸ , Gianluca Di Rico³⁸ , David Doelman¹⁸ , Mauro Dolci³⁸ , Frank Eisenhauer³² , Simone Esposito³¹ , Debora Ferruzzi³¹ , Helmut Feuchtgruber³² , Natascha Förster-Schreiber³² , Kyle Franson^{39,47} , Reinhard Genzel³² , Stefan Gillessen³² , Eileen C. Gonzales⁴⁰ , Michael Hartl³² , Jean Hayoz¹² , Heinrich Huber³² , Christoph Keller⁴¹ , Kateryna Kravchenko³² , Jarron Leisenring²⁷ , John Lightfoot³⁶ , David Lunney³⁶ , Dieter Lutz³² , Mike Macintosh³⁶ , Filippo Mannucci³¹ , Stanimir Metchev⁴² , Thomas Ott³² , David Pearson³⁶ , Alfio Puglisi³¹ , Sebastian Rabien³² , Christian Rau³² , Armando Riccardi³¹ , Bernardo Salasnich³⁵ , Taro Shimizu³² , Frans Snik¹⁸ , Eckhard Sturm³² , Genaro Suárez⁹ , Linda Tacconi³² , Xianyu Tan⁴³ , William Taylor³⁶ , Christopher Waring³⁶ , and Marco Xompero³¹

¹ Institute for Astronomy, University of Edinburgh, Royal Observatory, Blackford Hill, Edinburgh, EH9 3HJ, UK; ben.sutcliffe@roe.ac.uk

² Centre for Exoplanet Science, University of Edinburgh, Edinburgh, EH9 3HJ, UK

³ European Southern Observatory, Karl-Schwarzschild-Straße 2, 85748, Garching bei München, Germany; markus.bonse@eso.org

⁴ Max Planck Institute for Intelligent Systems, Max-Planck-Ring 4, 72076, Tübingen, Germany

⁵ Université Paris-Saclay, Université Paris Cité, CEA, CNRS, AIM, F-91191 Gif-sur-Yvette, France

⁶ Université Paris-Saclay, CNRS, Institut d'Astrophysique Spatiale, 91405 Orsay, France

⁷ STAR Institute, Université de Liège, Allée du Six Août 19c, 4000 Liège, Belgium

⁸ Max-Planck-Institut für Astronomie, Königstuhl 17, D-69117 Heidelberg, Germany

⁹ Department of Astrophysics, American Museum of Natural History, Central Park West at 79th Street, New York, NY 10024, USA

¹⁰ Astrophysics, Department of Physics, University of Oxford, Denys Wilkinson Building, Keble Road, Oxford, OX1 3RH, UK

¹¹ Department of Physics, University of Warwick, Gibbet Hill Road, Coventry CV4 7AL, UK

¹² ETH Zürich, Institute for Particle Physics and Astrophysics, Wolfgang-Pauli-Strasse 27, 8093 Zürich, Switzerland

¹³ Breakthrough Listen, Department of Physics, University of Oxford, Oxford OX1 3RH, UK

¹⁴ Department of Astronomy, University of Virginia, 530 McCormick Road, Charlottesville, VA 22904, USA

¹⁵ European Southern Observatory, Alonso de Córdova 3107, Vitacura, Casilla 19001, Santiago, Chile

¹⁶ Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

¹⁷ University of Exeter, Physics Building, Stocker Road, Exeter EX4 4QL, UK

¹⁸ Leiden Observatory, Leiden University, Postbus 9513, 2300 RA Leiden, The Netherlands

¹⁹ Department of Physics & Astronomy, Johns Hopkins University, 3400 N. Charles Street, Baltimore, MD 21218, USA

²⁰ Department of Astronomy, University of Texas at Austin, 2515 Speedway, Austin, TX 78712, USA

²¹ LIRA, Observatoire de Paris, Université PSL, CNRS, Sorbonne Université, Univ. Paris Diderot, Sorbonne Paris Cité, CY Cergy Paris Université, 5 place Jules Janssen, 92195 Meudon, France

²² Department of Physics, University of California, Santa Barbara, Santa Barbara, CA 93106, USA

²³ Université Grenoble Alpes, CNRS, IPAG, F-38000 Grenoble, France

²⁴ Department of Astronomy and Astrophysics, University of Toronto, 50 St. George Street, Toronto, ON M5S 3H4, Canada

²⁵ ETH Zürich, Department of Earth and Planetary Sciences, Sonneggstrasse 5, 8092 Zürich, Switzerland

²⁶ School of Physics, Trinity College Dublin, Dublin 2, Ireland

²⁷ Department of Astronomy and Steward Observatory, University of Arizona, 933 N. Cherry Avenue, Tucson, AZ 85721, USA

²⁸ Department of Physics and Astronomy, Northwestern University, 2145 Sheridan Road, Evanston, IL 60208-3112, USA

²⁹ Center for Interdisciplinary Exploration and Research in Astrophysics, 1800 Sherman Avenue, Northwestern University, Evanston, IL 60201, USA

³⁰ ELLIS Institute, Maria-von-Linden-Str. 2, 72076, Tübingen, Germany

³¹ INAF—Osservatorio Astrofisico di Arcetri, Largo E. Fermi 5., 50125, Firenze, Italy

³² Max Planck Institute for Extraterrestrial Physics, Giessenbachstraße 1, 85748 Garching, Germany

³³ Lunar and Planetary Laboratory, The University of Arizona, Tucson, AZ 85721, USA

³⁴ Department of Earth, Atmospheric and Planetary Sciences, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

³⁵ INAF—Osservatorio Astronomico di Padova, Vicolo dell'Osservatorio 5, 35122, Padova, Italy

³⁶ STFC UK Astronomy Technology Centre, Royal Observatory, Blackford Hill, Edinburgh, EH9 3HJ, UK

³⁷ I. Physikalisches Institut, Universität zu Köln, Zùlpicher Straße 77, 50937, Köln, Germany

³⁸ INAF—Osservatorio Astronomico d'Abruzzo, Via Mentore Maggini, 64100, Teramo, Italy

³⁹ Department of Astronomy & Astrophysics, University of California, Santa Cruz, 1156 High Street, Santa Cruz, CA 95064, USA

⁴⁰ Department of Physics and Astronomy, San Francisco State University, 1600 Holloway Avenue, San Francisco, CA 94132, USA

⁴¹ National Solar Observatory, 3665 Discovery Drive, Boulder, CO 80303, USA

⁴² Department of Physics & Astronomy, Western University, 1151 Richmond Street, London, ON N6A 3K7, Canada⁴³ Tsung-Dao Lee Institute and School of Physics and Astronomy, Shanghai Jiao Tong University, 1 Lisuo Road, Shanghai 200127, People's Republic of China
Received 2026 May 26; revised 2026 June 16; accepted 2026 June 17; published 2026 July 15

Abstract

We report the direct imaging discovery of a third exoplanet in the β Pictoris (β Pic) system. We detect β Pictoris d (β Pic d) in noncoronagraphic observations obtained with the Very Large Telescope (VLT) Enhanced Resolution Imager and Spectrograph (ERIS), as well as multi-epoch archival datasets from the JWST Near Infrared Camera (NIRCam) and VLT/SPHERE. Astrometric measurements over an 11 yr baseline demonstrate that it is consistent with a gravitationally bound source with orbital motion. Joint multi-planet orbit fits of all three planets in the system yield a semimajor axis of $26.0^{+2.2}_{-6.1}$ au and inclination $89.0^{+0.7}_{-0.6}$ deg for planet d. β Pic d has a larger orbital semimajor axis than the other known planets in the system, but is coplanar with the inner two planets, and its orbit is consistent with sculpting the inner edge of the debris disk. β Pic d has a contrast of $\Delta L' = 12.11 \pm 0.15$ mag, with colors and luminosity that closely match those of 51 Eri b, another exoplanet in the β Pic moving group. Its VLT/ERIS and JWST/NIRCam colors are distinct from those of free-floating planetary-mass objects of a similar age and temperature. Its red F410M – F444W color indicates strong CO₂ absorption in its atmosphere and suggests significant enhancement in metals compared to free-floating objects. From the ATMO hot-start evolutionary models, we estimate an effective temperature of 600^{+45}_{-60} K and mass of $2.4 \pm 0.6 M_{\text{Jup}}$, which also closely matches similar estimates for 51 Eri b. β Pic d is among the lowest-mass exoplanets imaged from the ground. This discovery highlights the deep sensitivity achievable with ground-based imaging in the mid-infrared and the discovery potential of future high-contrast observations with the Extremely Large Telescope.

Unified Astronomy Thesaurus concepts: Exoplanets (498); Exoplanet astronomy (486); Exoplanet systems (484); Direct imaging (387); High contrast techniques (2369); Orbital motion (1179); Debris disks (363); Very Large Telescope (1767); James Webb Space Telescope (2291)

1. Introduction

Since the detection of the first directly imaged exoplanets such as β Pictoris b (β Pic b; A.-M. Lagrange et al. 2010) and HR 8799 bcd (C. Marois et al. 2008), near-infrared (near-IR) direct imaging has revealed a population of dozens of young $>3 M_{\text{Jup}}$ planets with effective temperatures of 800–2000 K (see T. Currie et al. 2023a, for a recent review). Many of these planets reside in multi-planet systems (e.g., J. Milli et al. 2017; M. Keppler et al. 2018; S. Y. Haffert et al. 2019; A. J. Bohn et al. 2020; S. Hinkley et al. 2023; L. M. Close et al. 2025; R. F. van Capelleve et al. 2025a, 2025b; C. Lawlor et al. 2026), including both of the two emblematic first directly imaged exoplanetary systems; HR 8799 e (C. Marois et al. 2010) was announced soon after the other three HR 8799 planets, and β Pictoris c (β Pic c) was detected indirectly in 2019 (A.-M. Lagrange et al. 2019a). Following this trend, we report the discovery of a third giant exoplanet in the β Pic system. It seems likely that other directly imaged exoplanetary systems harbor additional lower-mass planets that are not detectable with current instrumentation.

Recent improvements in sensitivity in mid-infrared (mid-IR) instrumentation have enabled high-contrast imaging observations of a growing number of cold ($\lesssim 700$ K) giant exoplanets at 3–5 μm wavelengths close to their peak blackbody emission, including the detections of ϵ Ind A b (E. C. Matthews et al.

2024), TWA 7 b (K. A. Crotts et al. 2025; A.-M. Lagrange et al. 2025), and 14 Her c (D. C. Bardalez Gagliuffi et al. 2025) with JWST. At young ages (<100 Myr), this translates into the possibility to detect $<3 M_{\text{Jup}}$ exoplanets, which are more challenging to detect in the near-IR. While space-based facilities offer inherent stability and access to longer wavelengths, the largest ground-based telescopes remain highly complementary for mid-IR detections of such planets. This is in part because of the greater light-collecting power and angular resolution provided by their mirror sizes, which exceed that of JWST, e.g., the Very Large Telescope (VLT) Enhanced Resolution Imager and Spectrograph (ERIS; R. Davies et al. 2023).

With an age of 23 ± 8 Myr (R. A. Lee et al. 2024) and distance of 19.63 ± 0.06 pc (Gaia Collaboration et al. 2016, 2023), the A6V star β Pictoris (β Pic) hosts two known exoplanetary companions, β Pic b (A.-M. Lagrange et al. 2010) and β Pic c (A.-M. Lagrange et al. 2019a; M. Nowak et al. 2020). The outer planet, β Pic b, has a dynamical mass of $9.3^{+2.6}_{-2.5} M_{\text{Jup}}$ and a semimajor axis of 10.26 ± 0.10 au (A. M. Lagrange et al. 2020; G. M. Brandt et al. 2021), while β Pic c has a comparable dynamical mass of $8.89 \pm 0.75 M_{\text{Jup}}$ (J. Kammerer et al. 2025) but a much closer-in orbital semimajor axis of 2.68 ± 0.02 au (S. Lacour et al. 2021). Photometric and multiresolution spectroscopic observations of β Pic b find a partially cloudy atmosphere with water and carbon monoxide, $T_{\text{eff}} \sim 1700$ K, $\log(g) \sim 4.2$ dex, and rotational velocity $v \sin(i) \sim 20 \text{ km s}^{-1}$ (e.g., M. Bonnefoy et al. 2013; T. Currie et al. 2013; K. M. Morzinski et al. 2015; J. Chilcote et al. 2017; GRAVITY Collaboration et al. 2020; T. Stolker et al. 2020; J. Kammerer et al. 2024; L. T. Parker et al. 2024; R. Landman et al. 2024; M. Houll   et al. 2025; M. Ravet et al. 2025). Spectroscopy in the K band for β Pic c obtained with VLTI/GRAVITY (J. Kammerer et al. 2025) suggests a similar effective temperature and early L spectral type as β Pic b.

⁴⁴ These two authors contributed equally to this work and should be considered joint first authors.

⁴⁵ NSF Graduate Research Fellow.

⁴⁶ O.A. is a Research Director of the Fonds de la Recherche Scientifique—FNRS.

⁴⁷ NASA Sagan Fellow.



Original content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](https://creativecommons.org/licenses/by/4.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Table 1
List of Observations of β Pic Used in this Work, as well as the Astrometry and Photometry of β Pic d Detections

ID	Instrument	Obs. Date (yy:mm: dd UT)	Filter	Exp. (s)	Seeing (arcsec)	Field Rot. (deg)	S/N	Separation (mas)	PA (deg)	Δ Mag. (mag)	App. Mag. (mag)
1	VLT/ERIS	2025-12-03	L'	7395	0.71	63.9	14.9	1126.9 ± 3.9	210.35 ± 0.22	12.11 ± 0.15	15.57 ± 0.16
2	JWST/ NIRCam	2025-03-22	F410M	47040	...	9.9	5.1	1104 ± 20	211.1 ± 1.0	11.6 ± 0.2	15.2 ± 0.2
3	JWST/ NIRCam	2023-03-18	F444W	3667	...	10.2	6.6	1012 ± 20	210.4 ± 1.0	11.6 ± 0.2	14.7 ± 0.2
4	VLT/SPHERE	2020-02-08	$H3$	2024	0.60	30.7	5.9	784 ± 15	209.4 ± 1.0	<15.7	<19.2
5	VLT/SPHERE	2019-03-11	$K2$	2808	0.72	32.1	6.2	719 ± 12	209.6 ± 1.0	<14.9	<18.4
6	VLT/SPHERE	2014-12-08	$K1$	1980	0.77	26.1	7.9	375 ± 20	208.0 ± 2.0	14.1 ± 0.7	17.6 ± 0.7
		...	$K2$	...	...		7.2	...	...	13.6 ± 0.6	17.1 ± 0.6
A	VLT/NaCo	2011-12-11	L'	4800	0.57	54.1	...	...	...	...	...

Note. Epochs with a detection are assigned an ID number for convenient reference; the VLT/NaCo nondetection is indicated with “A”. For the JWST/NIRCam datasets, field rotation refers to the angular difference in roll positions. The S/N values for the ground-based observations are corrected for small number statistics (D. Mawet et al. 2014; M. J. Bonse et al. 2023), while we report *spaceKLIP* values for JWST/NIRCam (J. H. Girard et al. 2022; J. Kammerer et al. 2022; A. L. Carter et al. 2023). Due to SDI self-subtraction, limits are reported for datasets 4 and 5. Additional details are provided in the main text.

β Pic also possesses a large edge-on circumstellar debris disk (spanning between ~ 30 and 200 au from the star; B. A. Smith & R. J. Terrile 1984; I. Rebollido et al. 2024; Y. Han et al. 2026) that has long been connected to potential exoplanet formation in this system (e.g., H. Scholl et al. 1993; D. Lazzaro et al. 1994; C. J. Burrows et al. 1995; D. Mouillet et al. 1997; S. R. Heap et al. 2000; Z. Wahhaj et al. 2003; W. R. F. Dent et al. 2014; Y. Han et al. 2023; N. Skaf et al. 2023). The known planets are believed to have played a significant role in sculpting disk structures, in particular, the significant warp observed in the β Pic disk, with the inner ~ 80 au tilted by $\sim 5^\circ$ relative to the outer disk (C. J. Burrows et al. 1995; D. A. Golimowski et al. 2006). Both planets reside in near edge-on orbits coplanar with each other, although neither transit (L. van Sluijs et al. 2019; M. A. Kenworthy et al. 2021). The orbits of these planets are thought to be inclined relative to the outer disk, and hence drive the warp (D. Mouillet et al. 1997; J. L. Smallwood 2023). Recently, A. Lacquement et al. (2025) proposed the existence of an additional planet in the β Pic system to explain the observed inner debris disk edge at 30–50 au (Y. Han et al. 2026). They found that the two known planets are unable to sculpt the disk out to this radius, and that a planet capable of doing so would necessarily reside beyond the orbit of β Pic b.

In this Letter, we present the discovery of a third, wider-separation planet in the β Pic system, β Pic d, initially identified in noncoronagraphic direct imaging observations with VLT/ERIS. We confirm the planet through additional detections in earlier epochs of archival datasets from other instruments. We describe the corresponding data reduction methodology in Section 2. In Section 3, we use astrometry to confirm that the observed source is gravitationally bound to its host star and fit its orbital motion. We also use photometric measurements to begin characterizing the atmosphere of the planet and estimate its mass. We then discuss the implications of the results for the β Pic system in Section 4. Lastly, we summarize this work in Section 5. An independent and contemporaneous discovery of β Pic d using JWST/NIRSpec+MIRI observations is separately reported by A. Gibbs et al. 2026.

2. Observations and Data Reduction

We present results derived from observations obtained with four different instruments over 11 yr. A summary of these observations can be found in Table 1.

2.1. VLT/ERIS Observations

We observed the β Pic system on 2025 December 3 (02:15:33–05:18:07 UT) with the ERIS Near Infrared Camera System (NIX; D. Pearson et al. 2016) on the VLT as part of a UK Guaranteed Time Observations program with ESO program No. 116.28WQ.001 (PIs: Birkby and Biller). We used the 13 mas pixel $^{-1}$ camera and the L' -band filter, which has a central wavelength of 3.79 μ m, a bandwidth of 0.60 μ m, and zero-point flux density of 239.24 Jy in the Vega system.⁴⁸ To maximize sensitivity, we used the LM-pupil without a coronagraph. Compared to the Lyot stop of the vortex coronagraph (G. Orban de Xivry et al. 2024), this provides better core throughput, enabling deeper background limits. The observations were obtained in pupil-stabilized mode, which permits the use of angular differential imaging (ADI; C. Marois et al. 2006) for point-spread function (PSF) subtraction. We acquired 212 science frames on-target, each consisting of 4000 coadded frames with an integration time of 8.7214 ms per frame, for a total of 7395 s (~ 2 hr) after removing four frames with poor adaptive optics (AO) correction. We used nodding to facilitate background subtraction while avoiding regions of clustered bad pixels. These short individual exposure times ensured that the images remained unsaturated, which is essential to minimize artifacts caused by detector crosstalk and potential detector persistence.

We reduced the data using *eristools*, a Python package for processing VLT/ERIS high-contrast imaging data based on *PynPoint* (T. Stolker et al. 2019). The pipeline will be described in detail in a forthcoming publication (M. J. Bonse et al. 2026, in preparation). Using *eristools*, we perform the following steps: First, we

⁴⁸ Full filter information at: <https://svo2.cab.inta-csic.es/theory/fps/index.php?id=Paranal/ERIS.Lp>.




Figure 1. Gallery of images for the epochs at which β Pic d is detected (source indicated with a white arrow). The JWST/NIRCam images have been processed with a high-pass filter to reduce disk flux. The known planet β Pic b is the bright source in the northeast of the image in epochs 1–3. In epochs 4 and 5, β Pic b is within the inner mask used for data reduction. We note that for the 2014 December 8 epoch (epoch 6), β Pic b and β Pic d are nearly coincident; planet β Pic b (open circle) has been subtracted here to allow the much fainter β Pic d (arrow) to be seen. All images use the same normalized color scale and field of view, with ticks spaced at $1''$ intervals.

subtract dark frames and apply a flat-field correction using sky flats obtained at similar airmasses to those of the science frames. VLT/ERIS data also require correction for detector nonlinearity. We use *EsoRex* (ESO CPL Development Team 2015) to fit a second-order polynomial to each individual pixel. The resulting nonlinearity coefficients are imported into *eristools* to correct for the detector nonlinearity. Second, we subtract the background from each science frame using the frames at the opposite nod position. *eristools* identifies bad pixels in two steps. First, we generate a bad pixel map from the detector nonlinearity coefficients. We then identify any remaining outliers through a 4σ clipping within a local region of size 5×5 pixels. In the final step, we aligned all frames relative to each other using subpixel cross correlations, then centered them by fitting the unsaturated PSF with a Gaussian.

For the postprocessing, we applied a full-frame principal component analysis (PCA; A. Amara & S. P. Quanz 2012; R. Soummer et al. 2012) using *PynPoint* and 4S (M. J. Bonse et al. 2025), see T. D. Gebhard et al. (2022). We explored different sets of processing parameters. For the PCA, we calculated residuals with 1–50 principal components. For the 4S reductions, the images were spatially downsampled by a factor of 1.5 to keep the computational requirements manageable, and we varied the regularization parameter λ_{reg} between 10^5 and 5×10^6 . The final residual image shown in Figure 1 shows the 4S reduction with $\lambda_{\text{reg}} = 10^5$.

2.2. JWST/NIRCam Observations

We use two archival epochs of coronagraphic imaging observations targeting β Pic with the JWST (J. P. Gardner et al. 2006, 2023) Near Infrared Camera (NIRCam; M. J. Rieke et al. 2003, 2023).

The first of these was obtained on 2023 March 18 UT as part of GTO 1411 and was previously described by J. Kammerer et al. (2024). This dataset consists of observations in six filters, two in NIRCam’s short-wavelength channel (F182M, F210M) using the MASK210R coronagraphic occulting mask, and four in its long-wavelength channel (F250M, F300M, F335M, F444W) using the MASK335R mask. The short and long-wavelength observations have pixel scales of 31 and 63 mas pixel $^{-1}$, respectively. The total integration in each filter is given in Table 1 and in Appendix A. Each observation was obtained at two roll positions, separated by 10.2° , to enable PSF subtraction through ADI, and observations of a guide star, α Pictoris, were also acquired to allow reference-star differential imaging (RDI; e.g., J. E. Krist et al. 1998). The data reduction process for this dataset was carried out using the *pyKLIP* (J. J. Wang et al. 2015) and *spaceKLIP* (J. H. Girard et al. 2022; J. Kammerer et al. 2022; A. L. Carter et al. 2023) pipelines and is described in detail in J. Kammerer et al. (2024).

The second JWST/NIRCam epoch was obtained on 2025 March 21–22 (11:23:20–03:34:07 UT) as part of GO 4758 (Y. Zhou et al. 2026). These observations were acquired using

the F210M and F410M filters in dual-band coronagraphic imaging mode and using the MASK335R occulting mask. Each exposure covered 2352 s of integration time. This was repeated 10 times for each of two rolls, separated by 9.9, giving 23,520 s per roll and 47,040 s (~ 13.07 hr) in total. As with the previous epoch, PSF subtraction was performed using combined ADI and RDI using reference-star observations of α Pictoris. The data reduction process for this dataset is described in detail in Y. Zhou et al. (2026); however, this process followed the same steps as for the previous epoch (J. Kammerer et al. 2024). In brief, `spaceKLIP` was used to apply the stage 1 and 2 JWST pipelines⁴⁹ to produce flux-calibrated images, correct bad pixels, and recenter the images. For both epochs, a 5 pixel-wide high-pass filter was applied to the images prior to PSF subtraction with Karhunen–Loève Image Projection (KLIP; A. Amara & S. P. Quanz 2012; R. Soummer et al. 2012; L. Pueyo 2016) to remove smooth, spatially extended light from the debris disk. We performed the PSF subtraction for both epochs by using the `pyklip-pipeline.run_obs` function within `spaceKLIP` to apply KLIP with ADI+RDI, with three annuli, one subsection, and 50 Karhunen–Loève modes.

2.3. VLT/SPHERE Observations

We use three archival epochs obtained with the Spectro-Polarimetric High-contrast imager for Exoplanet REsearch (SPHERE; J.-L. Beuzit et al. 2019) on the VLT between 2014 December 8 UT and 2020 February 8 UT as part of ESO programs 60.A-9255(A), 60.A-9382(A), and 0104.C-0418(F), published in A.-M. Lagrange et al. (2019b) and A. M. Lagrange et al. (2020). These data were all obtained with SPHERE’s Infrared Dual-band Imager and Spectrograph (IRDIS; K. Dohlen et al. 2008) in either the K band using the $K12$ filter pair, or the H band using the $H23$ filter pair (see Table 1). The imaging was obtained in IRDIFS mode, meaning that simultaneous SPHERE Integral Field Spectrograph (IFS) observations were also acquired (R. U. Claudi et al. 2008). Out of 18 SPHERE observations of the system between 2014 and 2020, we focused on the three IRDIS sequences that were the most favorable for the detection of the β Pic d candidate based on observing conditions, total integration time, and field rotation. We do not detect the candidate in initial reductions of the accompanying IFS data and in additional IRDIS datasets (see the details in Appendix A). Therefore, we leave a deeper investigation of these data for a subsequent paper. For the three imaging epochs, we consider the nominal pixel scale and true north values of $12.255 \text{ mas pixel}^{-1}$ and -1.75° deg, respectively, given their stability over time (A.-L. Maire et al. 2016, 2021) and their consistency with the values inferred with dedicated astrometric calibration observations presented in A.-M. Lagrange et al. (2019b), A. M. Lagrange et al. (2020). The satellite spots were present in the entirety of all considered observations, enabling $\sim 1.5 \text{ mas}$ centering precision for the star when processing the image sequence (A.-L. Maire et al. 2021).

We reduced the SPHERE data using the `vcal-sphere` pipeline (V. Christiaens et al. 2023b), which leverages both the ESO recipe execution tool (v3.13.10) for basic calibration (dark subtraction, flat-fielding, linearity correction, bad pixel

identification, cube building) and routines from the Vortex Image Processing (VIP) for pre- and postprocessing (C. A. Gomez Gonzalez et al. 2017; V. Christiaens et al. 2023a). The VIP routines involved in the pipeline carry out a PCA-based sky subtraction, bad pixel clump identification and correction, coronagraphic image realignment based on the satellite spots, bad frame trimming, distortion correction, and modeling and subtraction of the stellar signals using PCA (A. Amara & S. P. Quanz 2012; R. Soummer et al. 2012).

For postprocessing, we considered two strategies to disentangle planet and speckle signals: ADI alone, and spectral differential imaging (SDI; W. Sparks & H. Ford 2002) followed by ADI through a two-step approach akin to the SADI method presented in V. Christiaens et al. (2019). In both cases, we tested PCA models using between 1 and 100 principal components. For the final images shown in Figure 1, we median-combined the reductions obtained with selected ranges of principal components: 5–40 for the 2014 ADI-only epoch, 10–40 for the 2019 SDI+ADI epoch, and 20–60 for the 2020 SDI+ADI epoch. Taking this additional median combination helps to further remove residual speckle noise while preserving the signal that is robust with varying numbers of principal components. We considered a single annular region for the 2019 and 2020 epochs, where the inner radius was set to $\sim 450 \text{ mas}$ in order to mask planet b, and the outer radius was limited to $1''$ so as not to significantly extend calculation time. For the 2019 and 2020 epochs, we used a threshold in rotation corresponding to a minimum motion of 1 FWHM in the middle of the annulus, while a smaller annulus (4 FWHM wide around the position of planet b) with no rotation threshold was considered for the 2014 epoch. For the latter, we obtained the final image after subtracting planet b from the data cube and carrying out a number of tests to assess the impact of this subtraction on the parameters estimated for planet d (see more details in Section 3.1 and Appendix B).

2.4. VLT/NaCo Observations

We reprocessed 10 archival L' -band datasets obtained with the NaCo (R. Lenzen et al. 2003; G. Rousset et al. 2003; J. H. V. Girard et al. 2012) instrument in the years between 2009 and 2015, as well as data from 2003 that led to the discovery of β Pic b (A.-M. Lagrange et al. 2009). All datasets, except the early data from 2003, were taken with pupil tracking and allowed for ADI. No coronagraph was used for any observation. For preprocessing, we use an automated pipeline based on `PynPoint` (T. Stolker et al. 2019) to perform the following steps: First, standard calibrations (dark subtraction and flat-field correction) are applied, followed by the removal of the thermal background using averaged sky frames. We correct bad pixels using a 4σ -clipping algorithm within a 5×5 local region. Next, we align the frames via subpixel cross correlation. Temporal binning is not performed prior to PSF subtraction in order to preserve temporal resolution and improve the alignment of frames. Finally, we use a PCA-based criterion to reject low-quality frames, including those affected by an open AO loop. As with the ERIS data, we use PCA (A. Amara & S. P. Quanz 2012; R. Soummer et al. 2012) and 4S (M. J. Bonse et al. 2025) for PSF subtraction. We varied the number of principal components between 1 and 100, and tested regularization of 4S between 10^4 and 10^6 . For PSF subtraction of the 2003 data, we first subtract the reference star HR 2435, which was used in

⁴⁹ https://spaceklip.readthedocs.io/en/latest/tutorials/tutorial_NIRCam_reductions.html

A.-M. Lagrange et al. (2009). We then create a pseudo-pupil-tracking dataset by rotating the frames based on their parallactic angle to align the spider pattern. Finally, we process the data in a similar manner to the newer NACO datasets using PCA.

3. Results

3.1. Detections of β Pic d

We present the final images from the epochs at which we detect β Pic d in Figure 1. We detect a clear point source, the putative β Pic d, southwest of the star in the VLT/ERIS data with a signal-to-noise ratio (S/N) of 14.9. We detect a compatible point source in the longest wavelength filters for each of the two JWST/NIRCam epochs, and for the three VLT/SPHERE epochs, including detections in both the $K12$ and $H23$ bands. We do not detect the source in the VLT/NaCo epochs. We summarize all of the detections in Table 1, and nondetections in Appendix A.

As we progressively recovered the β Pic d candidate in more datasets, we used `Octofitter` (W. Thompson et al. 2023) and `orbitize!` (S. Blunt et al. 2020, 2024) to fit preliminary single-planet orbits to the later epochs and predict the expected position of the candidate in earlier datasets. For the earliest SPHERE epoch, from 2014, these indicated that the β Pic d candidate would likely be very close to β Pic b. We therefore first subtracted the much brighter planet b from the 2014 datacube using the position and flux found by the negative fake companion (NegFC) method (A.-M. Lagrange et al. 2010; O. Wertz et al. 2017; T. Stolker et al. 2019). After removing β Pic b, we processed this cube with PCA-ADI and identified a point source located in the wings of the PSF of planet b (~ 45 mas apart). This detection at $S/N \sim 7.5$ is significant (D. Mawet et al. 2014; M. J. Bonse et al. 2023). It is also robust with respect to the uncertainties on the parameters of planet b used for the subtraction, although the latter dominate the uncertainty budget for the parameters estimated for the planet d candidate. We provide additional details in Appendix B.

For the 2019 and 2020 VLT/SPHERE epochs, we combined SDI and ADI to achieve a deeper contrast. This was necessary as the β Pic d candidate falls close to the extreme-AO corrected radius of VLT/SPHERE. At these separations, fast varying speckles affect the contrast achieved with ADI only. We only detect the candidate when combining SDI and ADI to remove these speckles.

At the epochs of available VLT/NaCo observations (between 2003 and 2015), our preliminary orbit fits predicted the candidate to be $< 0.5''$ from the star. We therefore did not detect the candidate at these separations in any of the VLT/NaCo observations because it was too close to the star or the much brighter planet b. However, as the sensitivity of these images at wider angular separation was sufficient to recover a source with the L' -band flux of the β Pic d candidate, these observations remained highly valuable for ruling out the possibility that the point source was a background source (see Section 3.3).

3.2. Astrometric and Photometric Measurements

We measured the astrometry and photometry of the source in the different datasets using NegFC as described in the

following subsections and report these values in Table 1 for each detection.

3.2.1. VLT/ERIS

Due to the high computing requirements of 4S, we run the NegFC injection for ERIS based on PCA. A high-pass filter similar to that of the JWST reduction was applied to prevent contamination due to the disk signal. The fitting error is very stable for 5–15 PCA components, with uncertainties of 0.6 mas in separation, $0.03''$ in PA, and 0.02 mag in photometry. In addition to the fitting errors from the NegFC injection, we consider the following sources of error for the astrometry and photometry: (1) a precision of ~ 0.25 pixels on centering of the star; (2) an additional uncertainty in the PA of 0.023° on the true north calibration (see the ERIS User Manual);⁵⁰ (3) the variability in the unsaturated reference PSF, which contributes an uncertainty of 0.125 mag.

3.2.2. JWST/NIRCam

For the JWST/NIRCam measurements, we used the routines in `spaceKLIP`, following the method applied by A. L. Carter et al. (2023) and J. Kammerer et al. (2024). We used these routines to model the KLIP-processed PSF of the candidate, accounting for companion self-subtraction, and fit these to the images using Markov Chain Monte Carlo (MCMC) sampling as integrated in `spaceKLIP` with `emcee` (D. Foreman-Mackey et al. 2013). We adopt conservative errors on the astrometry to account for uncertainties in the position of the host star behind the coronagraphic mask and the distortion correction in `spaceKLIP`, as noted by J. Kammerer et al. (2024), which combined introduce systematic uncertainties of ~ 10 mas. We combine this in quadrature with the statistical uncertainties from the MCMC fit and round up to the nearest 10 mas in $\Delta R.A.$ and $\Delta Dec.$ Although the disk flux was well removed by the high-pass filter during data reduction, we also adopt conservative errors on the photometry to account for the impact of any residual disk flux. We do this by measuring the approximate disk flux at the location of the β Pic d candidate and combining this as an error in quadrature with the uncertainties from the MCMC fit.

3.2.3. VLT/SPHERE

We detail our method for measuring the astrometry and photometry of the β Pic d candidate in the 2014 VLT/SPHERE data, for which β Pic b was first removed, in Appendix B.2. We estimate contrasts of $\Delta K1 = 14.1 \pm 0.7$ mag and $\Delta K2 = 13.6 \pm 0.6$ mag at this epoch. The considered errors on both the astrometry and photometry are conservative due to the uncertainty caused by the proximity to β Pic b, which is ~ 100 times brighter than β Pic d in $K1$ and $K2$.

For the 2019 and 2020 VLT/SPHERE epochs obtained by combining SDI and ADI, the uncertainty associated with the PA is dominated by residual speckle noise. We estimate this from the injection and recovery of 180 fake companions at the same flux and radial separation, but different azimuths, than that estimated for the planet d candidate. In comparison, the uncertainty associated with the true north is negligible

⁵⁰ https://www.eso.org/sci/facilities/paranal/instruments/eris/doc/ERIS_User_Manual_v117.1.pdf




Figure 2. The astrometry of β Pic d relative to its host star at each epoch (circles) compared to its expected position if the candidate were a stationary background source (corresponding crosses). The spiral line indicates the expected motion of a stationary background source, as calculated using the Gaia DR3 proper motion and parallax of β Pic and projected backward in time from the recent VLT/ERIS epoch (2025 December 3). The thick gray line represents the plane of the β Pic debris disk (PA $\sim 30^\circ$, I. Rebolledo et al. 2024).

($0.04''$; A.-L. Maire et al. 2021). The uncertainty associated with the radial separation for these SDI+ADI detections is dominated by the poorly quantified amount of SDI self-subtraction. During SDI, we scale the shortward band image (both in physical size and in flux) to match the speckle pattern of the longward band. This involves a flux scaling and radial motion outward of the companion by ~ 0.6 – 0.8 FWHM (at the 2019 and 2020 epochs), before subtraction. We adopted a conservative radial separation uncertainty that scales inversely with the radial motion of the planet between the two channels. This term reflects the maximum centroid displacement that could be introduced by SDI self-subtraction, which we consider negligible once the radial motion is larger than 1 FWHM. A dedicated two-channel forward-modeling approach is needed to accurately estimate the planet’s astrometry and its photometry in both channels, which is beyond the scope of this Letter and will be presented in a subsequent paper. Here, we report a lower limit on the flux in the longest wavelength band, either $K2$ or $H3$. This limit corresponds to the case where planet self-subtraction is negligible when the shorter-wavelength image is rescaled and subtracted from the longer-wavelength image before ADI. Using NegFC provides a flux estimate that is corrected for ADI self-subtraction, hence leading to an overall lower (upper) limit in flux (magnitude) contrast in $K2$ and $H3$. For $K2$, this limit is consistent with the estimate from 2014 obtained with ADI alone.

3.3. Common Proper Motion Analysis

We plot the astrometry of the β Pic d candidate relative to its host star at each epoch in Figure 2, alongside the corresponding positions that would be expected for a stationary background source at the same epochs, based on

the parallax (50.93 ± 0.15 mas) and proper motion ($\mu_\alpha = 5.16 \pm 0.20$ mas yr $^{-1}$, $\mu_\delta = 84.04 \pm 0.19$ mas yr $^{-1}$) of β Pic (Gaia DR3; Gaia Collaboration et al. 2016, 2023). At all epochs, the measured position of the candidate differs significantly from that expected of a background object, with the overall motion across the ~ 11 yr baseline deviating from the background track at the $>24\sigma$ level. Furthermore, the motion of the candidate appears to be coplanar with the edge-on β Pic debris disk (indicated with a thick gray line), as for the two known planets in the system.

As an additional check, we verified that no source was visible at the expected background location in the VLT/NaCo epochs taken in the L' filter. Although we did not detect the β Pic d candidate in these datasets, the sensitivity of the dataset (ID: A, Table 1) is sufficient to detect a background source if one were present. We inject an artificial source with $L' = 12.11$ mag at the background position on 2011 December 11 and recover the signal at an S/N of 6.3. Therefore, we conclude that the candidate is not a background source and that the observed motion is that of a gravitationally bound companion moving outward in its orbit.

3.4. Orbit Fitting

With an 11 yr baseline and significant orbital motion observed across available epochs, we were able to perform an analysis of β Pic d’s orbit and place initial constraints on its configuration. The detailed description of our orbit fitting procedure is presented in Appendix C, and we briefly summarize the key points here. We performed a joint multi-planet analysis of β Pic b, c, and d using *orvara* v1.1.2 (T. D. Brandt et al. 2021), complementing our new relative astrometry for planet d with literature measurements of the inner planets, along with radial velocity (RV) data and absolute astrometry of the star⁵¹. Since planet d is expected to have negligible influence on the star’s acceleration, we conservatively chose to neglect the mass of β Pic d in the fits, thereby assuming that the RVs and absolute astrometry are not sensitive to planet d (within measurement uncertainties), while still accounting for the motion of the star under the influence of the inner two planets when fitting the relative astrometry of planet d. This was enforced by setting a Gaussian prior of $10^{-6} \pm 10^{-9} M_\odot$ on planet d’s mass.

Results for the semimajor axis (a), eccentricity (e), and inclination (i) of planet d are shown in Figure 3, while the remaining orbital parameters are presented in C.2, along with results for planets b and c for which almost all orbital parameters are well constrained. The best constrained orbital elements for planet d are the inclination and longitude of the ascending node, characterizing the orientation of its orbit ($i_d = 89.0^{+0.7}_{-0.6}$ deg, $\Omega_d = 210.8^{+0.6}_{-0.4}$ deg), which indicate an orbit closely aligned with those of planets b ($i_b = 88.993^{+0.010}_{-0.011}$ deg, $\Omega_b = 211.787^{+0.010}_{-0.008}$ deg) and c ($i_c = 88.93^{+0.11}_{-0.09}$ deg, $\Omega_c = 211.08^{+0.05}_{-0.04}$ deg), despite adopting uninformed priors and without enforcing coplanarity with the known planetary system architecture. Since planet d is constrained only by relative astrometry, our inferred value of Ω_d is subject to the usual 180° degeneracy of relative astrometric orbits, as the observations alone cannot distinguish whether the orbital motion is directed toward or away from the observer. We therefore restricted the fit

⁵¹ We note that *orvara* reports the orbital parameters of the planets, not the star.




Figure 3. Marginalized posterior distributions for the semimajor axis, eccentricity, and inclination of planet d from our orbital fit, along with their joint posteriors. Lighter and darker shades show the complete posterior samples and the subset with orbit-crossing solutions removed, respectively. Contours over the joint posteriors indicate the 1σ and 2σ credible regions.

to values of Ω_d corresponding to the same direction of orbital motion as the other planets in the system. The alternative, degenerate solution corresponds to a coplanar but retrograde orbit and is obtained through the transformation $\lambda_d \rightarrow \lambda_d + 180^\circ$ and $\omega_d \rightarrow \omega_d + 180^\circ$, with the corresponding shift $\Omega_d \rightarrow \Omega_d + 180^\circ$.

On the other hand, a strong covariance is seen between e_d and a_d , with multiple families of solutions consistent with observations: low- e orbits at moderate a ; a branch of higher- e orbits extending to lower a and corresponding to our observation epochs being near apoastron; and a branch of higher- e orbits at larger a corresponding to our observation epochs being near periastron. As a result, although there is a weak preference for lower eccentricities, we can only place an upper limit on $e_d < 0.90$ at 2σ . Likewise, the semimajor axis posterior is bimodal, with peaks for the two sets of solutions at about 18 and 26 au.

In Figure 4, however, we compare our orbit fit to the relative astrometry measurements for planet d, as well as a deprojected view of the β Pic system. There is a subset of high- e , low- a orbital solutions for planet d that appear to cross the orbit of planet b (and in some cases, even planet c) and would thus likely be unstable. While the lack of mass measurement for planet d prevents any advanced dynamical stability analysis at this stage, we used the analytical criterion of D. P. Whitmire et al. (1998, Equation (6)), which provides a geometric test for whether two coplanar orbits intersect for any eccentricities and orientations. From this, we determined that 28% of the posterior solutions are in configurations where the orbits of planets b and d would intersect (light-blue orbits in Figure 4), making them likely unphysical. We thus discarded these solutions from our posterior samples.

The final posteriors after removal of orbit-crossing solutions are shown in the darker contours in Figure 3. Considering only the noncrossing orbits eliminates the highest- e configurations

of planet d, resulting in $e_d < 0.44$ at 2σ . Without the highly eccentric orbits, the semimajor axis distribution becomes unimodal, at $26.0^{+2.2}_{-6.1}$ au. Table 2 reports results for planet d's main orbital elements derived from the final posterior distribution restricted to noncrossing solutions (see Appendix C.2 for the remaining orbital parameters). We note that i_d and Ω_d are largely unaffected by the removal of orbit-crossing posterior solutions, as expected for truly independently constrained parameters. However, other orbital parameter posteriors are influenced by our Bayesian priors, particularly the uniform e_d prior, and therefore should be treated with caution.

3.5. Photometric Analysis and Mass Estimate

In this Section we use the color and luminosity of β Pic d to infer an effective temperature and mass, and study the planet's atmosphere. In Figure 5, we present color–magnitude diagrams (CMDs) comparing the ERIS and JWST photometry of β Pic d against other directly imaged planets, field brown dwarfs, and evolutionary model tracks, with young objects highlighted (see Appendix D for the details). There are five known planets and several free-floating brown dwarfs in the β Pic moving group (BPMG; B. Zuckerman et al. 2001); all are coeval and formed in a similar environment, providing a pristine comparative sample.

While the $L' - F444W$ color of β Pic d matches those of brown dwarfs, its $L' - F410M$ and $F410M - F444W$ colors diverge significantly from those of the brown dwarfs. Recent observations of warmer planetary companions have similarly shown very faint F410M magnitudes for their luminosity (W. O. Balmer et al. 2025b, 2026). The F444W filter has a pivot wavelength of $4.402 \mu\text{m}$ and $\Delta\lambda = 1.024 \mu\text{m}$, while the F410M filter covers a subset of the same wavelengths, with a pivot wavelength of $4.083 \mu\text{m}$ and $\Delta\lambda = 0.436 \mu\text{m}$. F444W encompasses both the $4.3 \mu\text{m}$ CO₂ feature and the $4.6 \mu\text{m}$ CO fundamental bandhead. Thus, β Pic d's red F410M – F444W color in particular suggests that the majority of light in the F444W bandpass is emitted on the red side of the bandpass, i.e., strong CO₂ absorption is present on the blue side.

This strong CO₂ absorption is in turn suggestive of an elevated atmospheric metallicity; we find that Sonora Flame Skimmer models (J. Mang et al. 2026, in preparation) with a vertical mixing parameter of $K_{zz} = 4$ are able to match the $L' - F410M$ colors of both the brown dwarf and planet populations, assuming that planets have higher metallicities, and the chosen K_{zz} value here is consistent with that observed in previous studies of 51 Eri b (A. Madurowicz et al. 2025) and field brown dwarfs of similar temperature (e.g., S. Mukherjee et al. 2024). However, these models are also unable to explain the F444W magnitude of β Pic d and other planets in the sample. Following W. O. Balmer et al. (2025b), who applied ExoRem models (J.-L. Baudino et al. 2015; B. Charnay et al. 2018) to the spectrum of 51 Eri b, we show the best-fitting model photometry using the same grid in Figure 5. This model provides a reasonable fit to all β Pic d photometry, explaining both the blue $L' - F410M$ and red F410M – F444W colors.

β Pic d has $L' - F410M$ colors as well as VLT/SPHERE K_1 -band and ERIS L' -band absolute magnitudes that closely match (within 1σ) 51 Eri b (B. Macintosh et al. 2015; A. Rajan et al. 2017; M. Samland et al. 2017; T. J. Dupuy et al. 2022; S. B. Brown-Sevilla et al. 2023; N. Whiteford et al. 2023; W. O. Balmer et al. 2025b; A. Denis et al. 2026). Out of the current cohort of directly imaged exoplanets, 51 Eri b is the




Figure 4. Left: astrometry of β Pic d (white circles) relative to the host star (white star). We show 150 randomly drawn samples from our posteriors, color-coded by eccentricity from low to high (purple to orange) unless the sample crosses the orbit of β Pic b (light blue). The solid black line indicates the lowest- χ^2 orbit. Middle: the same information deprojected by the corresponding inclination for each orbit. The lowest- χ^2 orbits for planets b (dark gray) and c (light gray) are plotted for comparison. Right: projected separation from the host star as a function of time for all three planets. The 2014 VLT/SPHERE point illustrates the near-coincident projected positions of planets b and d as seen from Earth.

Table 2
Properties of β Pic d Based on Astrometric and Photometric Analysis

Parameter	Value	Note
a_d (au)	$26.0^{+2.2}_{-6.1}$	noncrossing
P_d (yr)	91^{+18}_{-27}	noncrossing
e_d	<0.44	noncrossing, 2σ c.i.
i_d (deg)	$89.0^{+0.7}_{-0.6}$	prior-independent
Ω_d (deg)	$210.8^{+0.6}_{-0.4}$	prior-independent
m_d (M_{Jup})	2.4 ± 0.6	estimated
$T_{\text{eff},d}$ (K)	600^{+45}_{-60}	estimated
R_d (R_{Jup})	1.26 ± 0.03	estimated

Note. Orbital parameters are derived either from the posteriors with planet b orbit-crossing solutions removed or the normal analysis if the measurement is independent of priors. Mass, T_{eff} , and radius estimates are from a combination of ATMO 2020 chemical equilibrium and strong nonequilibrium evolutionary models, assuming an age of 23 ± 8 Myr.

only exoplanet with a mid-T spectral type discovered to date (T6.5; A. Rajan et al. 2017). Like β Pic d, it is a BPMG member: the two planets are likely coeval and formed from similar material. W. O. Balmer et al. (2025b) recently derived $T_{\text{eff}} = 632 \pm 13$ K and $R = 1.30 \pm 0.03 R_{\text{Jup}}$ for 51 Eri b, based on forward modeling of photometry and spectroscopy from Gemini/GPI, VLT/SPHERE, and JWST/NIRCam with the ExoRem model grid, confirming earlier temperature estimates from A. Rajan et al. (2017) and M. Samland et al. (2017). T. J. Dupuy et al. (2022) derived an upper limit to the mass of 51 Eri b of $<9 M_{\text{Jup}}$ based on astrometric constraints, and S. B. Brown-Sevilla et al. (2023) estimated a mass range of $2\text{--}4 M_{\text{Jup}}$ from forward modeling of VLT/SPHERE spectroscopy.

Since 51 Eri b is such a close analog to β Pic d, we estimated a bolometric correction for β Pic d based on the colors of 51 Eri b following the method of Section 4.2 of K. Franson et al. (2023). This allows us to extrapolate a bolometric luminosity from the available photometry with a model-independent method. We adopt a bolometric correction of $\text{BC}(L') = 0.06 \pm 0.27$ mag from $\log(L_{\text{bol}}/L_{\odot}) = -5.55 \pm 0.10$ dex (W. O. Balmer et al. 2025b) and $L' = 16.20 \pm 0.11$ mag (A. Rajan et al. 2017) at a distance of 29.9 pc. Thus, for β Pic d, with $M_{L'} = 14.11 \pm 0.16$ mag, our BC implies $\log(L_{\text{bol}}/L_{\odot}) = -5.67 \pm 0.13$ dex. This

luminosity is consistent with a range of initial entropies, including some low enough to correspond to “cold-start” formation scenarios (e.g., M. S. Marley et al. 2007; G.-D. Marleau & A. Cumming 2014). If the cold-start scenario is indeed favored for β Pic d, β Pic would become the first system known to host planets formed through both hot- and cold-start pathways, with β Pic b and c representing the hot-start regime. However, to obtain an estimate of its mass from models, we adopt the conservative limiting case of “hot-start” initial entropy.

Using this bolometric luminosity estimate for β Pic d with an age range of 23 ± 8 Myr, we applied rejection sampling following T. J. Dupuy et al. (2023), B. E. Miles et al. (2023), and A. L. Carter et al. (2023) to estimate the physical properties of β Pic d. Adopting the ATMO 2020 evolutionary models (M. W. Phillips et al. 2020, both equilibrium and strong nonequilibrium grids), we estimate a mass of $2.4 \pm 0.6 M_{\text{Jup}}$, radius of $1.26 \pm 0.03 R_{\text{Jup}}$, and effective temperature of 600^{+45}_{-60} K (Table 2). Considering only the ERIS L' -band photometry derived in this Letter and interpolating directly from the same models, we find masses of $2\text{--}3.5 M_{\text{Jup}}$, consistent with the values derived through our luminosity approach, further validating our method. These values are very close matches to literature values for 51 Eri b. Thus, β Pic d is the second mid-T type exoplanet discovered and is a very close analog to 51 Eri b.

4. Discussion

4.1. Implications for the β Pic System

β Pic is an extremely well-studied and dynamically active system, comprising multiple giant planets, a warped debris disk, signs of recent giant collisions, and intense exocometary activity near the star. Our discovery of an additional, massive planet in the system naturally raises several dynamical questions.

First, is β Pic d consistent with the disk warp? The main disk spans roughly 30–200 au (Y. Han et al. 2026), with the inner ~ 80 au tilted by $\sim 5^\circ$ relative to the outer disk (C. J. Burrows et al. 1995; D. A. Golimowski et al. 2006). This warp was ascribed to planets b and c, whose shared orbital plane is tilted relative to the outer disk (D. Mouillet et al. 1997; S. Lacour et al. 2021; J. L. Smallwood 2023; I. Macias et al. 2026). Our discovery of β Pic d is consistent with this picture; the new planet shares the orbital plane of b and c, as required by




Figure 5. CMDs of β Pic d, shown alongside photometry for directly imaged planets in BPMG (blue stars) and around other stars (brown stars), as well as free-floating brown dwarfs in BPMG (blue circles) and the field (pink circles). While young (BPMG) and field-age brown dwarfs have similar colors as a function of magnitude, the planets are markedly fainter at F410M than brown dwarfs with comparable magnitude. We show evolutionary models from ATMO (M. W. Phillips et al. 2020) and atmosphere models from Sonora Flame Skimmer (J. Mang et al. 2026 in preparation) and ExoRem (J.-L. Baudino et al. 2015; B. Charnay et al. 2018). For the atmosphere models, we assume $\log g = 4.0$ dex and $R = 1.3 R_{\text{Jup}}$. Sonora Flame Skimmer models adopt solar C/O, $K_{\text{zz}} = 4$, and two metallicities ($[\text{Fe}/\text{H}] = 0.0$ and 1.0), while the ExoRem model has $[\text{Fe}/\text{H}] = 0.5$, $\text{C}/\text{O} = 0.65$, and is chosen based on the best-fit model of 51 Eri b in W. O. Balmer et al. (2025b); this model also provides the closest match to β Pic d of all models shown.

the warp model. This is further evidence that the inner and outer regions of β Pic are generally misaligned, perhaps hinting that some past dynamical event, such as a flyby, tilted the outer system.

Second, is β Pic d sculpting the disk's inner edge? The other planets are too far inward, so it has long been inferred that another planet lies beyond b and sets the disk edge through planetesimal scattering (e.g., A. Lacquemet et al. 2025). Our discovery of β Pic d supports this picture. The exact location of the disk edge is difficult to constrain, because the disk is viewed edge-on to Earth; a nonparametric fit to Atacama Large Millimeter/submillimeter Array (ALMA) disk data yields a shallow, uncertain radial profile, whose inner edge could be anywhere from 30 to 50 au (Y. Han et al. 2026). Assuming an inner edge of 40 ± 5 au, we apply the *SculptingPlanet* code⁵² (T. D. Pearce et al. 2022) to dynamically predict the properties of a planet sculpting the disk edge, assuming a star mass of $1.83 \pm 0.03 M_{\odot}$ and age of 23 ± 8 Myr. The result is shown in Figure 6. If the disk edge is sculpted by a single planet, then that planet would lie in the white region of the figure, with a minimum mass of $1.3 \pm 0.1 M_{\text{Jup}}$ and a maximum semimajor axis of 31 ± 4 au (higher masses and smaller semimajor axes are also compatible). This compares very favorably with our adopted parameters for β Pic d, with an estimated mass of $2.4 \pm 0.6 M_{\text{Jup}}$ and semimajor axis of $26.0^{+2.2}_{-6.1}$ au. We therefore confirm that β Pic d is consistent with a planet sculpting the disk's inner edge. Given the uncertainty on the edge location, we cannot dynamically rule out additional planets beyond d. Furthermore, the lack of significant eccentricity in the ALMA disk (J. B. Lovell et al. 2026) suggests a low eccentricity for β Pic d. If the planet were significantly eccentric and sufficiently massive, then it would impose an eccentricity on the disk (V. Faramaz et al. 2014; T. D. Pearce & M. C. Wyatt 2014); the lack of such disk




Figure 6. Dynamical prediction for a planet sculpting the disk inner edge (T. D. Pearce et al. 2022). The predicted planet should lie in the white region to 1σ , in agreement with our adopted parameters for β Pic d. Line 1 shows orbits within 5 Hill radii of the disk, along which the planet would lie, and line 2 is the minimum planet mass to sculpt the disk within the star's age. Shading around lines denotes 1σ uncertainties.

eccentricity would support an eccentricity on the lower end of our prior-dependent posterior for e_d (see Section 3.4).

Third, is β Pic d related to the exocomets? β Pic displays significant exocomet activity, with star-grazing comets transiting every few days (R. Ferlet et al. 1987; A. Lecavelier des Etangs et al. 2022). This likely requires planets to drive comets from a debris belt inward toward the star (e.g., A. J. Mustill et al. 2026). One possibility is that planets b and c excite bodies from a close-in asteroid belt at ~ 1 au (H. Beust et al. 2024), while an alternative is that planets scatter bodies inward from the outer belt at ~ 40 au (e.g. L. Rodet and D. Lai 2024, K. P. Jaworska and H. J. Hoeijmakers 2026). β Pic d could do the latter; it appears close enough to the outer belt to scatter debris, some of which could be passed inward via planets b and c toward the star. If β Pic d is scattering comets inward, then the back reaction of comets on the planet would drive

⁵² <https://github.com/TimDPearce/SculptingPlanet>

β Pic d outward into the belt, exposing it to fresh material and keeping the interaction going (e.g., A. Bonsor et al. 2014).

4.2. The BPMG as a Laboratory for Exoplanet Formation

The BPMG contains five imaged exoplanets (β Pic bcd, 51 Eri b, and AF Lep b), as well as several free-floating planetary mass objects with similar properties to the imaged exoplanets (e.g., M. C. Liu et al. 2013; Z. Zhang et al. 2021). Together, these objects form a cohort of objects with a range of temperatures and masses but similar ages, both in disks around stars of varying masses and in the form of isolated planetary mass objects. This makes the BPMG a valuable laboratory for studying different modes of exoplanet formation, as well as the atmospheres and evolution of these objects (e.g., K. I. Öberg et al. 2011; P. Mollière et al. 2022; E. Nasedkin et al. 2024; J. Wang 2025; J.-B. Ruffio et al. 2026; J. W. Xuan et al. 2026). However, we note here that as planet formation may proceed at different rates in different systems, these planets likely do not share the exact same age (e.g., J. B. Pollack et al. 1996; R. H. Durisen et al. 2007; T. Currie et al. 2009, 2013).

The colors of β Pic d and 51 Eri b are notably similar. Comparative spectroscopic studies of these planets and other BPMG objects will reveal how the composition and chemistry of their atmospheres differ, and ideally also the imprint of the formation environment on these atmospheres. For the exoplanet companions, eventual dynamical mass measurements for the full sample will then allow these properties to be traced specifically as a function of mass, composition, age, and formation environment.

4.3. A Game of Hide-and-seek: The Necessity of Long Time Baselines for Exoplanet Confirmation

One of the main lessons of this study is that patience, as well as repeated observations, are necessary to search for and robustly confirm common proper motion for directly imaged exoplanet candidates. Due to the edge-on viewing angle of the β Pic system, all three of the exoplanets spend significant amounts of time in undetectable positions behind the star or within the radii of coronagraphic masks. Earlier studies of the β Pic system missed β Pic d largely because this planet was either at an undetectable separation from the star or very close to β Pic b in a given epoch of observation. This was also the case for AF Lep b; although this system was targeted by several imaging surveys (e.g., J. M. Stone et al. 2018; E. L. Nielsen et al. 2019; R. Launhardt et al. 2020), the planet’s orbital motion and high orbital inclination meant that it was not discovered until much later (R. J. De Rosa et al. 2023; K. Franson et al. 2023; D. Mesa et al. 2023). Planets with orbital semimajor axes ≤ 30 au, which display significant orbital motion like β Pic d, will routinely require multiple epochs to confirm. In the era of extremely large telescopes (ELTs), when direct imaging begins to observe e.g., known RV exoplanets with orbital semimajor axes < 10 au, this problem will persist, as these planets may lie behind the coronagraph mask for large portions of their orbits.

Orbital motion is not the only concern here, as stars with significant proper motions of their own can, in some cases, mimic a common proper motion exoplanet companion, at least for a few epochs. This has led to the retraction of multiple claims of directly imaged exoplanets, when, after a sufficiently large time baseline, it has become clear that the exoplanet

candidate was in fact just a background star with significantly different proper motion than the rest of the background field (E. L. Nielsen et al. 2017; M. Kenworthy et al. 2025). However, supplemental information can also provide compelling evidence that a candidate companion is a bona fide exoplanet before common proper motion confirmation; in the case of β Pic d, its red colors (compared to the blue colors expected for background stars and galaxies), orbital motion along the plane of the known circumstellar disk, and consistency with the expected dynamical picture of the disk strongly supported that it was indeed an exoplanet.

4.4. The Potential of ELT/METIS, as Revealed by ERIS-NIX

Like JWST, ERIS-NIX opens up a new discovery space for young directly imaged exoplanets by achieving deep sensitivity at wider separations. This gain in sensitivity results from the reduced number of warm optical surfaces, made possible by mounting the instrument at the Cassegrain focus behind the deformable secondary mirror (R. Davies et al. 2023). ERIS-NIX is equipped with two coronagraphs designed for high-contrast imaging: a vector vortex coronagraph (G. Orban de Xivry et al. 2024) and a vector Apodizing Phase Plate coronagraph (vAPP; D. S. Doelman et al. 2021; M. A. Kenworthy et al. 2026); however, the loss of throughput from coronagraphic optics limits sensitivity outside the contrast-limited regime. The ERIS-NIX noncoronagraphic imaging reported here achieved deep sensitivity, enabling a high-S/N detection of β Pic d in the L' band. This high-S/N L' detection then facilitated our lower S/N detections at shorter wavelengths and tighter separations. These results highlight the discovery potential of the upcoming Mid-infrared ELT Imager and Spectrograph (METIS; B. Brandl et al. 2021) instrument for the Extremely Large Telescope (ELT), which will have both coronagraphic optics and the possibility to obtain ultra-deep, noncoronagraphic imaging with a 39 m aperture.

5. Conclusions

We present the discovery of β Pic d, a third, widely separated giant planet in the β Pic system, with detections in six epochs of observations covering a total baseline of 11 yr. Our findings are summarized below:

1. We detect β Pic d at a contrast of $\Delta/L = 12.11 \pm 0.15$ mag and separation of 1126.9 ± 3.9 mas in our VLT/ERIS observations from 2025 December 3 and at closer separations in earlier JWST/NIRCam and VLT/SPHERE observations in several filters. Our astrometry shows that its motion is consistent with a gravitationally bound source with orbital motion approximately in the plane of the β Pic disk, and rules out the possibility that it is a background star or galaxy. β Pic d is currently moving outward in its orbit and was likely obscured by its host star and β Pic b in archival VLT/NaCo observations from approximately 2009 to 2011.
2. We estimate a mass of $2.4 \pm 0.6 M_{\text{Jup}}$, radius of $1.26 \pm 0.03 R_{\text{Jup}}$, and T_{eff} of 600^{+45}_{-60} K for β Pic d, interpolating from the ATMO 2020 evolutionary models. Its colors differ from brown dwarfs at similar absolute magnitudes, and its red F410M – F444W color suggests the presence of CO₂ in its atmosphere.

3. We use astrometry, along with literature measurements of the inner planets and host star, to fit the orbit and measure an inclination and longitude of the ascending node that are consistent with being coplanar with the inner planets. After excluding orbits that cross β Pic b's orbit, we find a semimajor axis of $26.0^{+2.2}_{-6.1}$ au, which corresponds to a period of 91^{+18}_{-27} yr, and eccentricity <0.44 (2σ). Such orbits are consistent with theoretical predictions for a planet sculpting the debris-disk inner edge.
4. β Pic d is a close analog of the BPMG exoplanet 51 Eri b and is only the second exoplanet discovered with a likely mid-T spectral type. This makes the β Pic planetary system the first known to contain both definitively “hot-start” planets (b and c) and a potentially “cold-start” planet. β Pic d is the least massive exoplanet detected in the β Pic system to date, and is among the lowest-mass exoplanets imaged from the ground.

The discovery of β Pic d using a combination of ground and space-based observations demonstrates the complementarity of these facilities and the value of archival datasets for investigating and confirming newly identified companions, even in well-known systems. β Pic d was missed in earlier studies of the system, in part because the planet was too close to its host star or the far brighter β Pic b for many epochs of observation. This illustrates the importance of revisiting compelling targets when conducting planet searches. Furthermore, the strong L' -band detection of this planet with VLT/ERIS emphasizes the deep contrast sensitivity that large ground-based telescopes still offer in the era of JWST, and hints at the potential of ELT/METIS for detecting even fainter, lower-mass exoplanets at wavelengths closer to the peak of their emission.

Future observations of β Pic d will characterize its atmosphere in depth, further constrain its orbit, and yield deeper insights into the architecture and formation of this young planetary system. This discovery also highlights the value of the BPMG, which now contains five imaged exoplanets with a range of temperatures and masses but similar ages, as a laboratory for understanding planetary atmospheres and evolution. Spectroscopic studies and dynamical mass measurements of this sample will probe how the properties of these planets differ, and thus reveal the imprint of their formation environment on their atmospheres.

Acknowledgments

The authors would like to thank the referee of this Letter, Thayne Currie, whose thoughtful and constructive comments helped us to improve this work.

The authors thank the JWST GTO 1411 (PI: Stark) and GO 4758 (PI: Zhou) teams for providing reduced data products and valuable advice. BJS and BAB acknowledge funding by the UK Science and Technology Facilities Council (STFC) grant nos. ST/V000594/1 and UKRI1196. L.T.P. and J.L.B. acknowledge funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program under grant agreement No 805445. L.T.P. further acknowledges funding support from Brasenose College. J.L.B. further acknowledges the support of the Leverhulme Trust via the Philip Leverhulme Physics Prize. T.D.P. is supported by a UKRI Stephen Hawking Fellowship and a

Warwick Prize Fellowship, the latter made possible by a generous philanthropic donation. E.O.G. gratefully acknowledges the financial support from the Swiss National Science Foundation (SNSF) under project grant number 200020_200399. L.I. is supported by funding from Breakthrough Listen. The Breakthrough Prize Foundation funds the Breakthrough Initiatives, which manage Breakthrough Listen. S.H. acknowledges funding by the UK Science and Technology Facilities Council (STFC) grant No. ST/Y002792/1. This work has been carried out within the framework of the NCCR PlanetS supported by the Swiss National Science Foundation under grants 51NF40_182901 and 51NF40_205606. This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (COBREX; grant agreement #885593). J.M.V. acknowledges support from a Royal Society Research Ireland University Research Fellowship (URF/1/221932, RF/ERE/221108) and the European Union through the Exo-PEA ERC project (grant number 101164652). Views and opinions expressed are however those of the author (s) only and do not necessarily reflect those of the European Union or the European Research Council Executive Agency. Neither the European Union nor the granting authority can be held responsible for them. This material is based upon work supported by the National Aeronautics and Space Administration under agreement No. 80NSSC21K0593 for the program “Alien Earths.” E.C.G. acknowledges support from the Heising-Simons Foundation.

This work is based on observations collected at the European Organization for Astronomical Research in the Southern Hemisphere under ESO programs 116.28WQ.001, 60.A-9255(A), 60.A-9382(A), and 0104.C-0418(F). We would like to thank the support staff at Paranal Observatory for their assistance in obtaining these observations. This work is based in part on observations made with the NASA/ESA/CSA JWST. The data were obtained from the Mikulski Archive for Space Telescopes at the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-03127 for JWST. These observations are associated with JWST programs GTO 1411 (PI: Stark) and GO 4758 (PI: Zhou).

This research has made use of the NASA Exoplanet Archive, which is operated by the California Institute of Technology, under contract with the National Aeronautics and Space Administration under the Exoplanet Exploration Program (J. L. Christiansen et al. 2025). This research has made use of the Astrophysics Data System, funded by NASA under Cooperative Agreement 80NSSC21M00561. This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France (M. Wenger et al. 2000). This research made use of SAOImageDS9, a tool for data visualization supported by the Chandra X-ray Science Center (CXC) and the High Energy Astrophysics Science Archive Center (HEASARC) with support from the JWST Mission office at the Space Telescope Science Institute for 3D visualization (W. A. Joye & E. Mandel 2003). This work has made use of data from the European Space Agency (ESA) mission Gaia (<https://www.cosmos.esa.int/gaia>), processed by the Gaia Data Processing and Analysis Consortium (DPAC, <https://www.cosmos.esa.int/web/gaia/dpac/consortium>) (Gaia Collaboration et al. 2016, 2023). Funding for the DPAC

has been provided by national institutions, in particular the institutions participating in the Gaia Multilateral Agreement. This research has made use of the Spanish Virtual Observatory (SVO) Filter Profile Service “Carlos Rodrigo,” funded by MCIN/AEI/10.13039/501100011033/ through grant PID2023-146210NB-I00 (C. Rodrigo & E. Solano 2020). This work made use of the *whereistheplanet*⁵³ prediction tool (J. J. Wang et al. 2021) to initialize the removal of β Pic b for our NegFC analysis. This work has made use of the *species* package for synthetic photometry and mass estimates using model isochrones (T. Stolker et al. 2020).

Data Availability

A data package for this Letter will be made available through a Zenodo repository shortly after publication, at doi:[10.5281/zenodo.20246422](https://doi.org/10.5281/zenodo.20246422). This will include either the datasets themselves or links to the archives from which they can be downloaded, as well as the config files for our orbit fitting procedure. The JWST/NIRCam data presented in this Letter were obtained from the Mikulski Archive for Space Telescopes (MAST) at the Space Telescope Science Institute. The specific observations analyzed can be accessed via DOI: [10.17909/wk9r-7797](https://doi.org/10.17909/wk9r-7797).

Facilities: VLT:Yepun (ERIS-NIX, NaCo), VLT:Melipal (SPHERE), VLT:Antu (NaCo), JWST.

Software: This work makes use of the Python programming language,⁵⁴ in particular packages, including NumPy (C. R. Harris et al. 2020), backtracks (W. O. Balmer et al. 2025a), Octofitter (W. Thompson et al. 2023), orbitize (S. Blunt et al. 2020, 2024), orvara (T. D. Brandt et al. 2021), astropy (Astropy Collaboration et al. 2013, 2018; Astropy Collaboration et al. 2022), astroquery (A. Ginsburg et al. 2019), EsoRex (ESO CPL Development Team 2015), pyKLIP (J. J. Wang et al. 2015), spaceKLIP (J. H. Girard et al. 2022; J. Kammerer et al. 2022; A. L. Carter et al. 2023), STPSF (M. D. Perrin et al. 2012; M. D. Perrin et al. 2014), fourS (M. J. Bonse et al. 2025), eristools (M. J. Bonse et al. 2026, in preparation), PynPoint (A. Amara & S. P. Quanz 2012; T. Stolker et al. 2019), vcal-sphere (V. Christiaens et al. 2023b), vip_hci (C. A. Gomez Gonzalez et al. 2017; V. Christiaens et al. 2023a), SculptingPlanet (T. D. Pearce et al. 2022), species (T. Stolker et al. 2020), Matplotlib (J. D. Hunter 2007), SciPy (P. Virtanen et al. 2020), emcee (D. Foreman-Mackey et al. 2013), and uncertainties (E. O. Lebigot et al. 2025).

Author Contributions

B.J.S. and M.J.B. were equally responsible for leading this project as well as the preparation and presentation of this manuscript. B.J.S. led the coordination of the team. B.J.S. also led the analysis of the JWST/NIRCam data, and M.J.B. led the analysis of the VLT/ERIS and VLT/NaCo data. V.C. led the analysis of the VLT/SPHERE data with the help of M.J.B. for the preliminary orbit fits. C.F. and T.D. led the orbit fitting analysis. E.C.M. and B.A.B. led the interpretation of the photometry of β Pic d. L.T.P. and B.J.S. acquired the VLT/ERIS observations. T.D.P. led the analysis of the implications of this discovery for the β Pic disk and system as a whole. J.L.B. and B.A.B. led the UK ERIS GTO team and were the PIs of the VLT/ERIS dataset that led to the discovery of β Pic d. Authors E.O.G. through B.S. provided relevant scientific expertise and contributed to the preparation of the manuscript. All other authors (G.A. through M.X.) are listed alphabetically and made crucial contributions to acquiring the observations described in this Letter or designing and developing the VLT/ERIS instrument central to this work.

Appendix A Nondetections

In addition to the detection in the archival data, we processed several datasets that did not result in the detection of β Pic d. These nondetections are attributable to the planet being too close to the star or to poor observing conditions. Despite all SPHERE data being acquired in dual IRDIS+IFS modes, we only reduced the 2019 and 2020 IFS datasets corresponding to the IRDIS data, in which we did detect β Pic d. This is for two reasons. First, 16 out of 18 datasets were taken in IFS (*YJ*) + IRDIS (*H23*) mode, where the IFS data span a short wavelength coverage, which limits the efficiency of SDI, where planet d is expected to be extremely faint. This includes the 2020 IFS data. Second, among the two remaining SPHERE datasets, acquired in IFS (*YJH*) + IRDIS (*K12*) mode (i.e., the 2014 and 2019 epochs), planet d was found at a much shorter separation and in the wings of planet b in the 2014 IRDIS data. This leaves the 2019 *YJH* IFS dataset as the most amenable for a redetection of planet d at shorter wavelengths. All the investigated datasets are summarized in Table 3.

⁵³ <http://whereistheplanet.com/>

⁵⁴ Python Software Foundation; <https://www.python.org/>

Table 3
List of β Pic Observations Analyzed Here That Did Not Yield a Detection

Instrument	Program	Obs. Date (yy:mm:dd UT)	Filter	Exp. (s)	Seeing (arcsec)	Field Rot. (deg)	Comment
JWST/NIRCam	GO 4758	2025-03-22	F210M	47,040	...	9.9	
JWST/NIRCam	GO 4758	2023-03-18	F182M	3014	...	10.2	
JWST/NIRCam	GTO 1411	2023-03-18	F210M	3014	...	10.2	
JWST/NIRCam	GTO 1411	2023-03-18	F250M	3421	...	10.2	
JWST/NIRCam	GTO 1411	2023-03-18	F300M	3421	...	10.2	
JWST/NIRCam	GTO 1411	2023-03-18	F335M	3667	...	10.2	
VLT/SPHERE	0104.C-0418(F)	2020-02-08	<i>YJ</i>	2024	0.60	30.7	IFS dataset.
VLT/SPHERE	0104.C-0418(D)	2019-12-22	<i>H23</i>	1260	0.59	30.7	Short total integration.
VLT/SPHERE	60.A-9382(A)	2019-03-11	<i>YJH</i>	2808	0.72	32.1	IFS dataset.
VLT/SPHERE	1100.C-0481(V)	2018-09-12	<i>H23</i>	1144	0.57	26.1	Short total integration and high airmass (~ 1.5).
VLT/SPHERE	0104.C-0418(F)	2016-10-14	<i>H23</i>	3872	0.72	53.5	Wind-driven halo. Tentative detection of d.
VLT/SPHERE	096.C-0241(E)	2016-03-26	<i>H23</i>	1664	1.65	19.9	
VLT/SPHERE	096.C-0241(G)	2016-01-20	<i>H23</i>	2328	1.15	29.8	
VLT/SPHERE	096.C-0241(B)	2015-11-28	<i>H23</i>	2104	1.13	40.0	Wind-driven halo.
VLT/SPHERE	095.C-0298(H)	2015-02-05	<i>H23</i>	3384	0.75	43.3	Very close proximity to b. Tentative detection of d.
VLT/NaCo	094.C-0149(D)	2015-01-23	<i>L'</i>	4699	0.99	30.6	Poor seeing and little field rotation.
VLT/NaCo	090.C-0396(D)	2012-12-16	<i>L'</i>	3360	0.72	67.6	β Pic d is too close to the star (< 300 mas).
VLT/NaCo	088.C-0358(C)	2011-12-11	<i>L'</i>	4800	0.57	54.1	β Pic d is too close to the star (< 300 mas).
VLT/NaCo	086.C-0341(B)	2010-09-28	<i>L'</i>	6000	0.80	51.7	β Pic d is too close to the star (< 300 mas).
VLT/NaCo	284.C-5057(A)	2010-04-07	<i>L'</i>	3360	1.23	18.3	β Pic d is too close to the star (< 300 mas).
VLT/NaCo	284.C-5057(A)	2010-03-21	<i>L'</i>	3360	1.36	19.9	β Pic d is too close to the star (< 300 mas).
VLT/NaCo	084.C-0739(A)	2009-12-29	<i>L'</i>	6060	0.83	61.0	β Pic d is too close to the star (< 300 mas).
VLT/NaCo	084.C-0739(A)	2009-12-26	<i>L'</i>	4800	0.86	44.0	β Pic d is too close to the star (< 300 mas).
VLT/NaCo	084.C-0739(A)	2009-12-25	<i>L'</i>	4800	1.18	50.8	β Pic d is too close to the star (< 300 mas).
VLT/NaCo	072.C-0624(B)	2003-11	<i>L'</i>	4113	0.69	...	No pupil tracking.

Note. The VLT/NaCo data taken in 2003-11 were collected in field-tracking between 2003-11-10 and 2003-11-17. Additional frames of the reference star HR 2435 were used for RDI.

Appendix B

Subtraction of β Pic b in the 2014 VLT/SPHERE Data

In 2014, β Pic d was very close to β Pic b, which outshines the faint companion by several orders of magnitude. To detect and characterize β Pic d, we used the NegFC technique implemented in VIP (C. A. Gomez Gonzalez et al. 2017; V. Christiaens et al. 2023a). We used NegFC twice: first, to model and subtract β Pic b to enable the detection of β Pic d, and second, to fit the astrometry and photometry of β Pic d. The analysis was carried out separately for the *K1* and *K2* bands.

B.1. Detection of β Pic d in 2014

Unlike explicit forward models (e.g., L. Pueyo 2016), which estimate the signature of a companion in the PCA residuals, the NegFC technique inserts a negative fake companion into the raw frames before PCA. This approach achieves a better accuracy than forward models to recover the astrometry and photometry of bright companions (e.g., F. Cantalloube et al. 2024). We refer to the data after the fake companion is subtracted as the *empty cube*. The empty cube is treated as a new dataset; that is, a new PCA basis is calculated to obtain the residual image. Using Nelder–Mead minimization, we optimized the position and flux of the inserted negative companion by minimizing an objective function defined as the standard deviation of the residual pixel intensities within a 1 FWHM aperture centered on the first-guess location of β Pic b. This process revealed β Pic d next to the original position of

β Pic b (see Figure 7, for residuals in the *K1* filter). We assessed the robustness of this detection by applying annular PCA with 1–100 components on the optimal empty cube, i.e., the cube that minimizes the NegFC objective function. The dataset offers a total field rotation of 26.1° , resulting in rotational motion of 171 mas. To preserve as many frames as possible for accurate subtraction of the speckle noise with PCA, we did not apply a rotation threshold. While the speckle pattern changes as a function of the principal components, the detection of β Pic d is robust for a wide range of components (5–40). The left column of Figure 7 shows residual images calculated on the original ADI cubes, including β Pic b. As the number of principal components increases, the signal of β Pic b is altered, resulting in a stripe pattern. This effect is caused by the brightness of the companion, which directly influences the PCA basis. The robust recovery of β Pic d for a wide range of principal components confirms that it is not a residual from a poor subtraction of β Pic b.

The residual image shown in Figure 1 was obtained by median combining all residual images between five and 40 components, and then averaging the result over the *K1* and *K2* filters.

B.2. Astrometry and Photometry of β Pic d

As for the other epochs, we estimated the astrometry and photometry of β Pic d using NegFC, but this time on the empty cube after subtracting β Pic b. To propagate the uncertainties associated with the subtraction of β Pic b into the retrieved




Figure 7. PSF-subtracted images obtained with annular PCA on the original 2014 VLT/SPHERE datacube (left column) and the cube resulting from the subtraction of our best estimate of β Pic b (right column), shown for 10, 20, and 40 principal components. The residuals shown are the results for the K1 filter. The ticks are spaced by 500 mas intervals.

parameters of β Pic d, we used the MCMC implementation of NegFC in VIP to draw samples from the posterior distribution of the position and flux of β Pic b. For each posterior sample of the β Pic b parameters, we generated a new empty cube and repeated the NegFC inference for β Pic d. We performed this procedure for 30 randomly selected posterior samples. For each realization, we considered two aperture radii for the objective function, 1 FWHM and 2 FWHM, and six 20-component PCA ranges between 10–30 and 60–80 components. Within each range, the corresponding PCA residuals were median-combined before evaluating the NegFC objective function. These choices reflect the facts that the speckle noise and residuals from β Pic b vary depending on the number of principal components, and that either choice of aperture size yielded plausible residual-minimization solutions. Through this analysis, we found that the uncertainty associated with the subtraction of β Pic b dominates the error budget of the inferred β Pic d parameters. By contrast, injection-and-recovery tests of individual fake planets with the inferred β Pic d parameters, but placed at different azimuths, yielded negligible additional uncertainties.

The final adopted uncertainties reflect the spread in the retrieved β Pic d parameters induced by varying the PCA range, the aperture radius used for the NegFC objective function, and the posterior sample of the β Pic b parameters used to construct the empty cube. We estimated contrast uncertainties for β Pic d of ~ 0.7 and ~ 0.6 mag in the K1 and K2 bands, respectively. Our final astrometric uncertainties

reflect the spread of radial separation and PA values inferred for both K1 and K2 (Table 1).

Appendix C Orbit Analysis

We performed an analysis of β Pic d’s orbit using our new relative astrometry as well as data from the literature. We used `orvara` v1.1.2 (T. D. Brandt et al. 2021), which can fit relative astrometry for multiple planets, host-star RVs, and absolute astrometry from Hipparcos and Gaia DR3. It accounts for the astrometric offsets of any planets interior to a given planet and neglects the influence of outer planets, which has been shown to be valid for the orbits of β Pic c and β Pic b to $\sim 20 \mu\text{as}$ over 20 yr (G. M. Brandt et al. 2021). The same work also showed that the impact of ignoring β Pic c when fitting β Pic b astrometry led to systematics of ~ 10 mas. Given that β Pic b’s orbit is about $4\times$ larger than β Pic c, its influence on β Pic d’s astrometry could be even larger, although the timescale of orbit variations is correspondingly longer, which may reduce its influence.

In the following, we adopted priors that are uniform in $\log M_*$ and $\log \sigma_{\text{jit}}$ for the host star and $\log a$, e , Ω , ω , λ , and $\sin i$, for all planets. To improve convergence, we used a start file to initialize our chains at values near the highest-likelihood parameters found in preliminary exploratory chains, and we imposed loose Gaussian priors on the masses of planets b and c of 0.009 ± 0.018 and $0.008 \pm 0.016 M_\odot$, respectively. Mainly due to the uncertain nature of the mass of β Pic d, we chose to neglect its mass in our orbit fits. By construction, it would not influence `orvara`’s fit of the inner two planets at all, and it is expected to be much smaller in mass than the inner planets, so it should induce a comparatively negligible acceleration on the host star ($\propto m_d/r_d^2$). We used 100 walkers, five parallel-tempering temperatures, 10^6 steps, thinned every 50th step, and we discarded the first 50% of steps as burn-in. The configuration files used for our orbit fit are provided in the Zenodo data package connected to this manuscript.

C.1. Literature Data

To perform our multi-planet orbit analysis, we included literature measurements of relative astrometry and RVs, as well as the most recent Hipparcos-Gaia Catalog of Accelerations (HGCA; T. D. Brandt 2021). We used the host-star RVs from A.-M. Lagrange et al. (2019a) cleaned of stellar activity as presented by T. Vandal et al. (2020). For the innermost planet β Pic c, only VLT/IRISA has resolved it, and we used all four reprocessed measurements from S. Lacour et al. (2021). For β Pic b, we use the following: all nine VLT/NACO measurements from 2003 to 2011 processed by G. Chauvin et al. (2012); the 2008 M' -band detection from T. Currie et al. (2011); the 2012 epoch from M. Bonnefoy et al. (2013); eight Gemini/NICI and Magellan/MagAO measurements over 2009–2012 from E. L. Nielsen et al. (2014, which excludes one discrepant CH_4S measurement from 2012); 14 Gemini/GPI measurements spanning 2013–2018 from J. J. Wang et al. (2016) as reported in E. L. Nielsen et al. (2020, which excludes one 2013 measurement with an anomalously high PA); all 34 VLT/SPHERE measurements spanning 2014–2020 from A.-M. Lagrange et al. (2019b) and A. M. Lagrange et al. (2020); and all seven VLT/IRISA measurements from 2018 to 2021 as reprocessed by S. Lacour

Table 4
Posterior from the Orvara Analysis

Parameter		Mode $\pm 1\sigma$	95.4% c.i.
M_* ($M_\odot$)	*	$1.789^{+0.024}_{-0.027}$	1.737, 1.838
σ_{jit} (m s $^{-1}$)	*	48^{+10}_{-8}	33, 70
m_b (M_{Jup})	*	$8.7^{+2.6}_{-2.5}$	3.5, 13.7
a_b (au)	*	10.07 ± 0.03	10.00, 10.13
$e \sin \omega_b$	*	$0.131^{+0.017}_{-0.016}$	0.097, 0.164
$e \cos \omega_b$	*	$0.2966^{+0.0023}_{-0.0022}$	0.2919, 0.3010
e_b	...	$0.105^{+0.004}_{-0.003}$	0.098, 0.113
ω_b (deg)	...	$23.7^{+3.1}_{-2.6}$	17.9, 29.3
i_b (deg)	*	$88.993^{+0.010}_{-0.011}$	88.972, 89.013
Ω_b (deg)	*	$211.787^{+0.010}_{-0.008}$	211.771, 211.806
λ_b (deg)	*	$322.6^{+0.8}_{-0.6}$	321.2, 324.1
T_b^{peri} (yr)	...	1990.27 ± 0.04	1990.19, 1990.35
P_b (yr)	...	$23.77^{+0.15}_{-0.16}$	23.45, 24.08
m_c (M_{Jup})	*	$8.7^{+0.8}_{-0.7}$	7.2, 10.4
a_c (au)	*	2.70 ± 0.03	2.65, 2.76
$e \sin \omega_c$	*	$-0.504^{+0.026}_{-0.022}$	-0.545, -0.444
$e \cos \omega_c$	*	$-0.237^{+0.024}_{-0.016}$	-0.268, -0.191
e_c	...	$0.307^{+0.019}_{-0.025}$	0.254, 0.346
ω_c (deg)	...	$-114.9^{+2.6}_{-2.4}$	-120.3, -109.9
i_c (deg)	*	$88.93^{+0.11}_{-0.09}$	88.74, 89.14
Ω_c (deg)	*	$211.08^{+0.05}_{-0.04}$	211.00, 211.18
λ_c (deg)	*	288^{+23}_{-13}	266, 329
T_c^{peri} (yr)	...	$2019.490^{+0.018}_{-0.027}$	2019.429, 2019.525
P_c (yr)	...	$3.29^{+0.07}_{-0.03}$	3.23, 3.42
a_d (au)	*	...	14.9, 36.7
$e \sin \omega_d$	*	...	-0.44, 0.52
$e \cos \omega_d$	*	$-0.80^{+0.69}_{-0.04}$	-0.85, 0.48
e_d	...	$0.10^{+0.31}_{-0.10}$	0.00, 0.90
ω_d (deg)	...	...	-170, 180
i_d (deg)	*	$88.9^{+0.9}_{-0.7}$	86.5, 90.7
Ω_d (deg)	*	$210.9^{+0.7}_{-0.6}$	209.6, 213.5
λ_d (deg)	*	270^{+17}_{-64}	135, 299
T_d^{peri} (yr)	...	2003 ± 10	1985, 2030
P_d (yr)	...	...	42, 165

Note. An “*” next to a parameter indicates that it was directly varied in the MCMC; any other parameters were computed from combinations of these. For parameters lacking their mode $\pm 1\sigma$ values, the posterior was strongly multimodal. The reference epoch is 2010.0.

et al. (2021). The HGCA does not contain a significant detection of astrometric acceleration ($\chi^2 = 0.935$), but it can still be useful in constraining the planet masses, as the Hipparcos and Gaia proper motion anomaly nondetections have uncertainties of about 0.4 and 0.3 mas yr $^{-2}$, respectively, so we include it.

C.2. Results

The best-fit orbit, i.e., the parameters that minimize χ^2 , resulted in $\chi^2 = 161$ for the relative astrometry alone, which is reasonable given that we fit 84 measurements with ≈ 21 free parameters (approximate because some are also constrained by RVs) and thereby 147 degrees of freedom. The subset of our new relative astrometry for planet d independently has $\chi^2 = 3.2$ for 6 degrees of freedom, implying that the astrometric errors from our analysis of VLT/ERIS, VLT/SPHERE, and JWST/NIRCam data are reasonable.

We report the resulting orbital parameters of all three planets in Table 4. The marginalized posteriors (Figure 8) for many of the orbital parameters are quite narrow and thus not influenced by our priors. These include all of the parameters for planet b and all but the mean longitude at the reference epoch for planet c ($\lambda_c = 288^{+23}_{-13}$ deg). While uniform priors on angles like λ are generally well justified, the same is not true for astrophysical parameters like eccentricity and semimajor axis. The parameters that we determine to be prior-independent for β Pic d are the inclination i_d and longitude of ascending node Ω_d . In Table 5, we report results for planet d from the subset of orbital solutions that do not cross the orbit of planet b.

In addition to our baseline prior, we also experimented with allowing planet d to have some mass in our orbit fits. However, this seemed to result in attempts to “overfit” the residuals of the astrometric and radial accelerations (after removing b and c) by exploring unrealistically massive and eccentric orbits for planet d. Given that we expect planet d to have a negligible impact on the host-star acceleration, these experiments justify our choice of a massless prior on planet d (implemented as $10^{-6} \pm 10^{-9} M_\odot$ in orvara).




Figure 8. Marginalized posterior distributions for our orbital parameters of planets d (gray), c (red), and b (yellow) in the β Pic system. Contour lines and shaded regions in the joint posterior indicate the 1σ and 2σ credible regions. The posteriors for planet d include the full posterior distributions from the fits (light gray) along with the posteriors after removal of solutions crossing the orbits of the inner planets (dark gray). As planet d was set to be massless, it is not shown in the first column. The orbital parameters shown are the semimajor axes, eccentricities, inclinations, arguments of periaapsis ω , longitudes of ascending node Ω , and mean longitudes λ at reference epoch $T_{\text{ref}} = 2010.0$.

Table 5Posteriors for β Pic d Excluding Orbits That Cross β Pic b's Orbit

Parameter		Mode $\pm 1\sigma$	95.4% c.i.
a_d (au)	*	$26.0^{+2.2}_{-6.1}$	17.9, 39.7
$e \sin \omega_d$	*	$-0.26^{+0.30}_{-0.09}$	-0.42, 0.21
$e \cos \omega_d$	*	$-0.57^{+0.62}_{-0.05}$	-0.64, 0.54
e_d	...	$0.10^{+0.12}_{-0.10}$	0.00, 0.44
ω_d (deg)	...	...	-170, 180
i_d (deg)	*	$89.0^{+0.7}_{-0.6}$	87.7, 90.3
Ω_d (deg)	*	$210.8^{+0.6}_{-0.4}$	210.0, 212.0
λ_d (deg)	*	270^{+10}_{-33}	214, 302
T_d^{peri} (yr)	...	2000^{+21}_{-7}	1982, 2032
P_d (yr)	...	91^{+18}_{-27}	56, 186

Note. See Table 4 for details; those posteriors were adopted as input for this subsample. An “*” next to a parameter indicates that it was directly varied in the MCMC; any other parameters were computed from combinations of these.

Appendix D CMD Data

To construct the CMD shown in Figure 5, we use literature photometry and spectroscopy for a variety of sources. Planet observations for HR 8799 bcde are from W. O. Balmer et al. (2025b), G. Bryden et al. (2026), C. Marois et al. (2008, 2010); for HIP 65426 b they are from A. L. Carter et al. (2023), A. C. Cheetham et al. (2019); for HIP 99770 b from W. O. Balmer et al. (2026), T. Currie et al. (2023b); for 51 Eri b from W. O. Balmer et al. (2025b), A. Rajan et al. (2017); for AF Lep b from M. J. Bonse et al. (2025), K. Franson et al. (2024); for β Pic b from J. Kammerer et al. (2024), T. Currie et al. (2013), Y. Zhou et al. (2026), and for β Pic d from this work.

For brown dwarfs, we plot synthetic photometry derived from observed spectra. The field-age object photometry is derived from JWST spectra published in S. A. Beiler et al. (2024). To assemble known BPMG members, we used the UltracoolSheet (W. M. J. Best et al. 2025), which includes a synthesis of membership information derived from Banyan Σ (J. Gagné et al. 2018) and the literature, based on the approach described in A. Sanghi et al. 2023. This yielded a list of 18 field ultracool dwarf systems with membership determinations from the UltracoolSheet T. Dupuy et al. (2018), J. D. Kirkpatrick et al. (2021), and Z. Zhang et al. (2021). The CMD also shows PSO J318.5338–22.8603, which was observed with JWST (P. Mollière et al. 2025), as well as all BPMG members that have SPHEREx (J. J. Bock et al. 2026) spectra in the library presented by J. Gagné et al. (2026).

ORCID iDs

Ben J. Sutcliffe <https://orcid.org/0000-0002-9962-132X>
 Markus J. Bonse <https://orcid.org/0000-0003-2202-1745>
 Valentin Christiaens <https://orcid.org/0000-0002-0101-8814>
 Clémence Fontanive <https://orcid.org/0000-0002-2428-9932>
 Elisabeth C. Matthews <https://orcid.org/0000-0003-0593-1560>
 Luke T. Parker <https://orcid.org/0009-0009-8137-9991>
 Tim D. Pearce <https://orcid.org/0000-0001-5653-5635>
 Jayne L. Birkby <https://orcid.org/0000-0002-4125-0140>

Beth A. Biller <https://orcid.org/0000-0003-4614-7035>
 Trent J. Dupuy <https://orcid.org/0000-0001-9823-1445>
 Emily O. Garvin <https://orcid.org/0000-0003-2530-9330>
 Leyla Iskandarli <https://orcid.org/0009-0000-8422-2222>
 Jens Kammerer <https://orcid.org/0000-0003-2769-0438>
 Yifan Zhou <https://orcid.org/0000-0003-2969-6040>
 Robert J. De Rosa <https://orcid.org/0000-0002-4918-0247>
 Aarynn L. Carter <https://orcid.org/0000-0001-5365-4815>
 Sasha Hinkley <https://orcid.org/0000-0001-8074-2562>
 Matthew A. Kenworthy <https://orcid.org/0000-0002-7064-8270>
 William O. Balmer <https://orcid.org/0000-0001-6396-8439>
 Iain Hammond <https://orcid.org/0000-0003-1502-4315>
 James Mang <https://orcid.org/0000-0001-5864-9599>
 Caroline V. Morley <https://orcid.org/0000-0002-4404-0456>
 Mark J. Neeser <https://orcid.org/0009-0002-2455-7500>
 Olivier Absil <https://orcid.org/0000-0002-4006-6237>
 Anthony Boccaletti <https://orcid.org/0000-0001-9353-2724>
 Mariangela Bonavita <https://orcid.org/0000-0002-7520-8389>
 Brendan P. Bowler <https://orcid.org/0000-0003-2649-2288>
 Xueqing Chen <https://orcid.org/0009-0005-9339-2369>
 Felix A. Dannert <https://orcid.org/0000-0002-5476-2663>
 Julien H. Girard <https://orcid.org/0000-0001-8627-0404>
 Markus Kasper <https://orcid.org/0000-0002-8425-6606>
 Anne-Marie Lagrange <https://orcid.org/0000-0002-2189-2365>
 Pengyu Liu <https://orcid.org/0000-0001-7047-0874>
 Gilles Orban de Xivry <https://orcid.org/0000-0002-4790-415X>
 Michael Poon <https://orcid.org/0000-0001-7739-9767>
 Sascha P. Quanz <https://orcid.org/0000-0003-3829-7412>
 Benoît Serra <https://orcid.org/0009-0005-5104-5772>
 Johanna M. Vos <https://orcid.org/0000-0003-0489-1528>
 Kevin Wagner <https://orcid.org/0000-0002-4309-6343>
 Jason Wang <https://orcid.org/0000-0003-0774-6502>
 Bernhard Schölkopf <https://orcid.org/0000-0002-8177-0925>
 Guido Agapito <https://orcid.org/0000-0002-6382-2613>
 Dániel Apai <https://orcid.org/0000-0003-3714-5855>
 Andrea Baruffolo <https://orcid.org/0000-0002-1114-4355>
 Marco Bonaglia <https://orcid.org/0000-0002-4236-2339>
 Runa Briguglio <https://orcid.org/0000-0002-0495-0543>
 Yixian Cao <https://orcid.org/0000-0001-5301-1326>
 Luca Carbonaro <https://orcid.org/0000-0003-1492-1591>
 Giovanni Cresci <https://orcid.org/0000-0002-5281-1417>
 Richard Davies <https://orcid.org/0000-0003-4949-7217>
 Ivan Di Antonio <https://orcid.org/0000-0003-3260-4389>
 Amico Di Cianno <https://orcid.org/0000-0003-4508-5645>
 Gianluca Di Rico <https://orcid.org/0000-0002-5213-8269>
 David Doelman <https://orcid.org/0000-0003-0695-0480>
 Mauro Dolci <https://orcid.org/0000-0001-8000-5642>
 Simone Esposito <https://orcid.org/0000-0002-3114-677X>
 Debora Ferruzzi <https://orcid.org/0000-0003-1475-5976>
 Kyle Franson <https://orcid.org/0000-0003-4557-414X>
 Reinhard Genzel <https://orcid.org/0000-0002-2767-9653>
 Stefan Gillessen <https://orcid.org/0000-0002-5708-0481>
 Eileen C. Gonzales <https://orcid.org/0000-0003-4636-6676>
 Jean Hayoz <https://orcid.org/0000-0003-3768-5712>
 Christoph Keller <https://orcid.org/0000-0002-1368-841X>
 Jarron Leisenring <https://orcid.org/0000-0002-0834-6140>
 Dieter Lutz <https://orcid.org/0000-0003-0291-9582>

Mike Macintosh  <https://orcid.org/0009-0000-5761-3332>
 Filippo Mannucci  <https://orcid.org/0000-0002-4803-2381>
 Stanimir Metchev  <https://orcid.org/0000-0003-3050-8203>
 Thomas Ott  <https://orcid.org/0000-0003-1572-0396>
 Alfio Puglisi  <https://orcid.org/0000-0001-9553-8804>
 Armando Riccardi  <https://orcid.org/0000-0001-5460-2929>
 Bernardo Salasnich  <https://orcid.org/0000-0002-7502-2701>
 Taro Shimizu  <https://orcid.org/0000-0002-2125-4670>
 Frans Snik  <https://orcid.org/0000-0003-1946-7009>
 Eckhard Sturm  <https://orcid.org/0000-0002-0018-3666>
 Genaro Suárez  <https://orcid.org/0000-0002-2011-4924>
 Linda Tacconi  <https://orcid.org/0000-0002-1485-9401>
 Xianyu Tan  <https://orcid.org/0000-0003-2278-6932>
 Christopher Waring  <https://orcid.org/0009-0009-3690-0821>
 Marco Xompero  <https://orcid.org/0000-0002-5565-084X>

References

- Amara, A., & Quanz, S. P. 2012, *MNRAS*, **427**, 948
 Astropy Collaboration, Price-Whelan, A. M., Lim, P. L., et al. 2022, *ApJ*, **935**, 167
 Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, *AJ*, **156**, 123
 Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, *A&A*, **558**, A33
 Balmer, W. O., Kammerer, J., Pueyo, L., et al. 2025b, *AJ*, **169**, 209
 Balmer, W. O., Otten, G. P. P. L., & Stolker, T. 2025a, backtracks: a python package to compare relative astrometry with background helical motion, v0.6.0, Zenodo, doi:10.5281/zenodo.14838369
 Balmer, W. O., Pueyo, L., Messier, A., et al. 2026, *ApJL*, **1001**, L26
 Bardalez Gagliuffi, D. C., Balmer, W. O., Pueyo, L., et al. 2025, *ApJL*, **988**, L18
 Baudino, J.-L., Bézard, B., Boccaletti, A., et al. 2015, *A&A*, **582**, A83
 Beiler, S. A., Cushing, M. C., Kirkpatrick, J. D., et al. 2024, *ApJ*, **973**, 107
 Best, W. M. J., Dupuy, T. J., Liu, M. C., et al. 2025, The UltracoolSheet: Photometry, Astrometry, Spectroscopy, and Multiplicity for 4000+ Ultracool Dwarfs and Imaged Exoplanets, v2.1.0, Zenodo, doi:10.5281/zenodo.15802304
 Beust, H., Milli, J., Morbidelli, A., et al. 2024, *A&A*, **683**, A89
 Beuzit, J.-L., Vigan, A., Mouillet, D., et al. 2019, *A&A*, **631**, A155
 Blunt, S., Wang, J., Hirsch, L., et al. 2024, *JOSS*, **9**, 6756
 Blunt, S., Wang, J. J., Angelo, I., et al. 2020, *AJ*, **159**, 89
 Bock, J. J., Aboobaker, A. M., Adamo, J., et al. 2026, *ApJ*, **999**, 139
 Bohn, A. J., Kenworthy, M. A., Ginski, C., et al. 2020, *ApJL*, **898**, L16
 Bonnefoy, M., Boccaletti, A., Lagrange, A.-M., et al. 2013, *A&A*, **555**, A107
 Bonse, M. J., Garvin, E. O., Gebhard, T. D., et al. 2023, *AJ*, **166**, 71
 Bonse, M. J., Gebhard, T. D., Dannert, F. A., et al. 2025, *AJ*, **169**, 194
 Bonsor, A., Raymond, S. N., Augereau, J.-C., & Ormel, C. W. 2014, *MNRAS*, **441**, 2380
 Brandl, B., Bettonvil, F., van Boekel, R., et al. 2021, *Msngr*, **182**, 22
 Brandt, G. M., Brandt, T. D., Dupuy, T. J., Li, Y., & Michalik, D. 2021, *AJ*, **161**, 179
 Brandt, T. D. 2021, *ApJS*, **254**, 42
 Brandt, T. D., Dupuy, T. J., Li, Y., et al. 2021, *AJ*, **162**, 186
 Brown-Sevilla, S. B., Maire, A.-L., Mollière, P., et al. 2023, *A&A*, **673**, A98
 Bryden, G., Llop-Sayson, J., Beichman, C. A., et al. 2026, *AJ*, **171**, 301
 Burrows, C. J., Krist, J. E., Stapelfeldt, K. R., & WFPC2 Investigation Definition Team 1995, AAS Meeting, **187**, 32.05
 Cantalloube, F., Christiaens, V., Cantero, C., et al. 2024, *SPIE*, **13097**, 1309713
 Carter, A. L., Hinkley, S., Kammerer, J., et al. 2023, *ApJL*, **951**, L20
 Charnay, B., Bézard, B., Baudino, J.-L., et al. 2018, *ApJ*, **854**, 172
 Chauvin, G., Lagrange, A.-M., Beust, H., et al. 2012, *A&A*, **542**, A41
 Cheetham, A. C., Samland, M., Brems, S. S., et al. 2019, *A&A*, **622**, A80
 Chilcote, J., Pueyo, L., De Rosa, R. J., et al. 2017, *AJ*, **153**, 182
 Christiaens, V., Casassus, S., Absil, O., et al. 2019, *MNRAS*, **486**, 5819
 Christiaens, V., Gonzalez, C., Farkas, R., et al. 2023a, *JOSS*, **8**, 4774
 Christiaens, V., Hammond, I., Juillard, S., Kokoulina, E., & Balsalobre-Ruza, O. 2023b, VCAL-SPHERE: Hybrid pipeline for reduction of VLT/SPHERE data, Astrophysics Source Code Library, ascl:2311.002
 Christiansen, J. L., McElroy, D. L., Harbut, M., et al. 2025, *PSJ*, **6**, 186
 Claudi, R. U., Turatto, M., Gratton, R. G., et al. 2008, *SPIE*, **7014**, 70143E
 Close, L. M., van Capelleveen, R. F., Weible, G., et al. 2025, *ApJL*, **990**, L9
 Crotts, K. A., Carter, A. L., Lawson, K., et al. 2025, *ApJL*, **987**, L41
 Currie, T., Biller, B., Lagrange, A., et al. 2023a, *ASPC*, **534**, 799
 Currie, T., Brandt, G. M., Brandt, T. D., et al. 2023b, *Sci*, **380**, 198
 Currie, T., Burrows, A., Madhusudhan, N., et al. 2013, *ApJ*, **776**, 15
 Currie, T., Lada, C. J., Plavchan, P., et al. 2009, *ApJ*, **698**, 1
 Currie, T., Thalmann, C., Matsumura, S., et al. 2011, *ApJL*, **736**, L33
 Davies, R., Absil, O., Agapito, G., et al. 2023, *A&A*, **674**, A207
 De Rosa, R. J., Nielsen, E. L., Wahhaj, Z., et al. 2023, *A&A*, **672**, A94
 Denis, A., Vigan, A., Chauvin, G., et al. 2026, *A&A*, **707**, L13
 Dent, W. R. F., Wyatt, M. C., Roberge, A., et al. 2014, *Sci*, **343**, 1490
 Doelman, D. S., Snik, F., Por, E. H., et al. 2021, *ApOpt*, **60**, D52
 Dohlen, K., Langlois, M., Saisse, M., et al. 2008, *SPIE*, **7014**, 70143L
 Dupuy, T. J., Liu, M. C., Allers, K. N., et al. 2018, *AJ*, **156**, 57
 Dupuy, T. J., Brandt, G. M., & Brandt, T. D. 2022, *MNRAS*, **509**, 4411
 Dupuy, T. J., Liu, M. C., Evans, E. L., et al. 2023, *MNRAS*, **519**, 1688
 Durisen, R. H., Boss, A. P., Mayer, L., et al. 2007, in *Protostars and Planets V*, ed. B. Reipurth, D. Jewitt, & K. Keil (Univ. of Arizona Press), **607**
 ESO CPL Development Team 2015, EsoRex: ESO Recipe Execution Tool,, Astrophysics Source Code Library, ascl:1504.003
 Faramaz, V., Beust, H., Thébault, P., et al. 2014, *A&A*, **563**, A72
 Ferlet, R., Hobbs, L. M., & Vidal-Madjar, A. 1987, *A&A*, **185**, 267
 Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. 2013, *PASP*, **125**, 306
 Franson, K., Bowler, B. P., Zhou, Y., et al. 2023, *ApJL*, **950**, L19
 Franson, K., Balmer, W. O., Bowler, B. P., et al. 2024, *ApJL*, **974**, L11
 Gagné, J., Mamajek, E. E., Malo, L., et al. 2018, *ApJ*, **856**, 23
 Gagné, J., Faherty, J. K., Ruiz Diaz, A., et al. 2026, arXiv:2604.22012
 Gaia Collaboration, Prusti, T., de Bruijne, J. H. J., et al. 2016, *A&A*, **595**, A1
 Gaia Collaboration, Vallenari, A., Brown, A. G. A., et al. 2023, *A&A*, **674**, A1
 Gardner, J. P., Mather, J. C., Abbott, R., et al. 2023, *PASP*, **135**, 068001
 Gardner, J. P., Mather, J. C., Clampin, M., et al. 2006, *SSRv*, **123**, 485
 Gebhard, T. D., Bonse, M. J., Quanz, S. P., & Schölkopf, B. 2022, *A&A*, **666**, A9
 Gibbs, A., Ruffio, J.-B., Bidot, A., et al. 2026, arXiv:2606.23789
 Ginsburg, A., Sipőcz, B. M., Brasseur, C. E., et al. 2019, *AJ*, **157**, 98
 Girard, J. H., Leisenring, J., Kammerer, J., et al. 2022, *SPIE*, **12180**, 121803Q
 Girard, J. H. V., O'Neal, J., Mawet, D., et al. 2012, *SPIE*, **8447**, 84470L
 Golimowski, D. A., Ardila, D. R., Krist, J. E., et al. 2006, *AJ*, **131**, 3109
 Gomez Gonzalez, C. A., Wertz, O., Absil, O., et al. 2017, *AJ*, **154**, 7
 GRAVITY Collaboration, Nowak, M., Lacour, S., et al. 2020, *A&A*, **633**, A110
 Haffert, S. Y., Bohn, A. J., de Boer, J., et al. 2019, *NatAs*, **3**, 749
 Han, Y., Mansell, E., Jennings, J., et al. 2026, *A&A*, **705**, A196
 Han, Y., Wyatt, M. C., & Dent, W. R. F. 2023, *MNRAS*, **519**, 3257
 Harris, C. R., Millman, K. J., van der Walt, S. J., et al. 2020, *Nature*, **585**, 357
 Heap, S. R., Lindler, D. J., Lanz, T. M., et al. 2000, *ApJ*, **539**, 435
 Hinkley, S., Lacour, S., Marleau, G.-D., et al. 2023, *A&A*, **671**, L5
 Houllé, M., Millour, F., Berio, P., et al. 2025, *A&A*, **704**, A182
 Hunter, J. D. 2007, *CSE*, **9**, 90
 Jaworska, K. P., & Hoeijmakers, H. J. 2026, *A&A*, **708**, A59
 Joye, W. A., & Mandel, E. 2003, *ASPC*, **295**, 489
 Kammerer, J., Girard, J., Carter, A. L., et al. 2022, *SPIE*, **12180**, 121803N
 Kammerer, J., Lawson, K., Perrin, M. D., et al. 2024, *AJ*, **168**, 51
 Kammerer, J., Winterhalder, T. O., Lacour, S., et al. 2025, *A&A*, **704**, A318
 Kenworthy, M., Stolker, T., Kammerer, J., et al. 2025, *A&A*, **701**, A104
 Kenworthy, M. A., Dannert, F. A., Hayoz, J., et al. 2026, *A&A*, **708**, A239
 Kenworthy, M. A., Mellon, S. N., Bailey, J. I., et al. 2021, *A&A*, **648**, A15
 Keppler, M., Benisty, M., Müller, A., et al. 2018, *A&A*, **617**, A44
 Kirkpatrick, J. D., Gelino, C. R., Faherty, J. K., et al. 2021, *ApJS*, **253**, 7
 Krist, J. E., Golimowski, D. A., Schroeder, D. J., & Henry, T. J. 1998, *PASP*, **110**, 1046
 Lacour, S., Wang, J. J., Rodet, L., et al. 2021, *A&A*, **654**, L2
 Lacquement, A., Beust, H., Faramaz-Gorka, V., & Duchêne, G. 2025, *A&A*, **694**, A236
 Lagrange, A.-M., Boccaletti, A., Langlois, M., et al. 2019b, *A&A*, **621**, L8
 Lagrange, A.-M., Bonnefoy, M., Chauvin, G., et al. 2010, *Sci*, **329**, 57
 Lagrange, A.-M., Gratadour, D., Chauvin, G., et al. 2009, *A&A*, **493**, L21
 Lagrange, A.-M., Meunier, N., Rubini, P., et al. 2019a, *NatAs*, **3**, 1135
 Lagrange, A. M., Rubini, P., Nowak, M., et al. 2020, *A&A*, **642**, A18
 Lagrange, A.-M., Wilkinson, C., Málin, M., et al. 2025, *Nature*, **642**, 905
 Landman, R., Stolker, T., Snellen, I. A. G., et al. 2024, *A&A*, **682**, A48
 Launhardt, R., Henning, T., Quirrenbach, A., et al. 2020, *A&A*, **635**, A162
 Lawlor, C., Capelleveen, R. F., Bourdarot, G., et al. 2026, *ApJL*, **1000**, L38

- Lazzaro, D., Sicardy, B., Roques, F., & Greenberg, R. 1994, *Icar*, **108**, 59
- Lebigot, E. O., Gerber, J., Newville, M., et al. 2025, *Imfit/uncertainties: v3.2.3*, Zenodo, doi:10.5281/zenodo.15257927
- Lecavelier des Etangs, A., Cros, L., Hébrard, G., et al. 2022, *NatSR*, **12**, 5855
- Lee, R. A., Gaidos, E., van Saders, J., Feiden, G. A., & Gagné, J. 2024, *MNRAS*, **528**, 4760
- Lenzen, R., Hartung, M., Brandner, W., et al. 2003, *SPIE*, **4841**, 944
- Liu, M. C., Magnier, E. A., Deacon, N. R., et al. 2013, *ApJL*, **777**, L20
- Lovell, J. B., Hales, A. S., Kennedy, G. M., et al. 2026, *A&A*, **705**, A200
- Macias, I., Jenkins, S. A., & Vanderburg, A. 2026, *AJ*, **171**, 197
- Macintosh, B., Graham, J. R., Barman, T., et al. 2015, *Sci*, **350**, 64
- Madurowicz, A., Ruffio, J.-B., Macintosh, B., et al. 2025, *AJ*, **170**, 326
- Maire, A.-L., Langlois, M., Delorme, P., et al. 2021, *JATIS*, **7**, 035004
- Maire, A.-L., Langlois, M., Dohlen, K., et al. 2016, *SPIE*, **9908**, 990834
- Marleau, G.-D., & Cumming, A. 2014, *MNRAS*, **437**, 1378
- Marley, M. S., Fortney, J. J., Hubickyj, O., Bodenheimer, P., & Lissauer, J. J. 2007, *ApJ*, **655**, 541
- Marois, C., Lafrenière, D., Doyon, R., Macintosh, B., & Nadeau, D. 2006, *ApJ*, **641**, 556
- Marois, C., Macintosh, B., Barman, T., et al. 2008, *Sci*, **322**, 1348
- Marois, C., Zuckerman, B., Konopacky, Q. M., Macintosh, B., & Barman, T. 2010, *Natur*, **468**, 1080
- Matthews, E. C., Carter, A. L., Pathak, P., et al. 2024, *Natur*, **633**, 789
- Mawet, D., Milli, J., Wahhaj, Z., et al. 2014, *ApJ*, **792**, 97
- Mesa, D., Gratton, R., Kervella, P., et al. 2023, *A&A*, **672**, A93
- Miles, B. E., Biller, B. A., Patapis, P., et al. 2023, *ApJL*, **946**, L6
- Milli, J., Hibon, P., Christiaens, V., et al. 2017, *A&A*, **597**, L2
- Mollière, P., Kühnle, H., Matthews, E. C., et al. 2025, *A&A*, **703**, A79
- Mollière, P., Molyarova, T., Bitsch, B., et al. 2022, *ApJ*, **934**, 74
- Morzinski, K. M., Males, J. R., Skemer, A. J., et al. 2015, *ApJ*, **815**, 108
- Mouillet, D., Larwood, J. D., Papaloizou, J. C. B., & Lagrange, A. M. 1997, *MNRAS*, **292**, 896
- Mukherjee, S., Fortney, J. J., Morley, C. V., et al. 2024, *ApJ*, **963**, 73
- Mustill, A. J., Pearce, T., Bannister, M., et al. 2026, arXiv:2604.08190
- Nasedkin, E., Mollière, P., Lacour, S., et al. 2024, *A&A*, **687**, A298
- Nielsen, E. L., De Rosa, R. J., Macintosh, B., et al. 2019, *AJ*, **158**, 13
- Nielsen, E. L., De Rosa, R. J., Wang, J. J., et al. 2020, *AJ*, **159**, 71
- Nielsen, E. L., Liu, M. C., Wahhaj, Z., et al. 2014, *ApJ*, **794**, 158
- Nielsen, E. L., Rosa, R. J. D., Rameau, J., et al. 2017, *AJ*, **154**, 218
- Nowak, M., Lacour, S., Lagrange, A.-M., et al. 2020, *A&A*, **642**, L2
- Öberg, K. I., Murray-Clay, R., & Bergin, E. A. 2011, *ApJL*, **743**, L16
- Orban de Xivry, G., Absil, O., De Rosa, R. J., et al. 2024, *SPIE*, **13097**, 1309715
- Parker, L. T., Birkby, J. L., Landman, R., et al. 2024, *MNRAS*, **531**, 2356
- Pearce, T. D., Launhardt, R., Ostermann, R., et al. 2022, *A&A*, **659**, A135
- Pearce, T. D., & Wyatt, M. C. 2014, *MNRAS*, **443**, 2541
- Pearson, D., Taylor, W., Davies, R., et al. 2016, *SPIE*, **9908**, 99083F
- Perrin, M. D., Sivaramakrishnan, A., Lajoie, C. P., et al. 2014, *SPIE*, **9143**, 91433X
- Perrin, M. D., Soummer, R., Elliott, E. M., Lallo, M. D., & Sivaramakrishnan, A. 2012, *SPIE*, **8442**, 84423D
- Phillips, M. W., Tremblin, P., Baraffe, I., et al. 2020, *A&A*, **637**, A38
- Pollack, J. B., Hubickyj, O., Bodenheimer, P., et al. 1996, *Icar*, **124**, 62
- Pueyo, L. 2016, *ApJ*, **824**, 117
- Rajan, A., Rameau, J., De Rosa, R. J., et al. 2017, *AJ*, **154**, 10
- Ravet, M., Bonnefoy, M., Chauvin, G., et al. 2025, *A&A*, **704**, A325
- Rebollido, I., Stark, C. C., Kammerer, J., et al. 2024, *AJ*, **167**, 69
- Rieke, M. J., Baum, S. A., Beichman, C. A., et al. 2003, *SPIE*, **4850**, 478
- Rieke, M. J., Kelly, D. M., Misselt, K., et al. 2023, *PASP*, **135**, 028001
- Rodet, L., & Lai, D. 2024, *MNRAS*, **527**, 11664
- Rodrigo, C., & Solano, E. 2020, in XIV.0 Scientific Meeting (Spanish Astronomical Society), **182**
- Rousset, G., Lacombe, F., Puget, P., et al. 2003, *SPIE*, **4839**, 140
- Ruffio, J.-B., Xuan, J. W., Chachan, Y., et al. 2026, *NatAs*, **10**, 511
- Samland, M., Mollière, P., Bonnefoy, M., et al. 2017, *A&A*, **603**, A57
- Sanghi, A., Liu, M. C., Best, W. M. J., et al. 2023, *ApJ*, **959**, 63
- Scholl, H., Roques, F., & Sicardy, B. 1993, *CeMDA*, **56**, 381
- Skaf, N., Boccaletti, A., Pantin, E., et al. 2023, *A&A*, **675**, A35
- Smallwood, J. L. 2023, *MNRAS*, **523**, 3526
- Smith, B. A., & Terrile, R. J. 1984, *Sci*, **226**, 1421
- Soummer, R., Pueyo, L., & Larkin, J. 2012, *ApJL*, **755**, L28
- Sparks, W., & Ford, H. 2002, *ApJ*, **578**, 543
- Stolker, T., Bonse, M. J., Quanz, S. P., et al. 2019, *A&A*, **621**, A59
- Stolker, T., Quanz, S. P., Todorov, K. O., et al. 2020, *A&A*, **635**, A182
- Stone, J. M., Skemer, A. J., Hinz, P. M., et al. 2018, *AJ*, **156**, 286
- Thompson, W., Lawrence, J., Blakely, D., et al. 2023, *AJ*, **166**, 164
- van Capelleveen, R. F., Ginski, C., Kenworthy, M. A., et al. 2025a, *ApJL*, **990**, L8
- van Capelleveen, R. F., Kenworthy, M. A., Ginski, C., et al. 2025b, *A&A*, **704**, A221
- van Sluijs, L., de Mooij, E., Kenworthy, M., et al. 2019, *A&A*, **626**, A97
- Vandal, T., Rameau, J., & Doyon, R. 2020, *AJ*, **160**, 243
- Virtanen, P., Gommers, R., Oliphant, T. E., et al. 2020, *NatMe*, **17**, 261
- Wahhaj, Z., Koerner, D. W., Ressler, M. E., et al. 2003, *ApJL*, **584**, L27
- Wang, J. 2025, *ApJ*, **981**, 138
- Wang, J. J., Graham, J. R., Pueyo, L., et al. 2016, *AJ*, **152**, 97
- Wang, J. J., Kulikauskas, M., & Blunt, S. 2021, whereistheplanet: Predicting positions of directly imaged companions., Astrophysics Source Code Library, ascl:2101.003
- Wang, J. J., Ruffio, J.-B., De Rosa, R. J., et al. 2015, pyKLIP: PSF Subtraction for Exoplanets and Disks., Astrophysics Source Code Library, ascl:1506.001
- Wenger, M., Ochsenbein, F., Egret, D., et al. 2000, *A&AS*, **143**, 9
- Wertz, O., Absil, O., Gómez González, C. A., et al. 2017, *A&A*, **598**, A83
- Whiteford, N., Glasse, A., Chubb, K. L., et al. 2023, *MNRAS*, **525**, 1375
- Whitmire, D. P., Matese, J. J., Criswell, L., & Mikkola, S. 1998, *Icar*, **132**, 196
- Xuan, J. W., Ruffio, J.-B., Chachan, Y., et al. 2026, *ApJ*, **1000**, 27
- Zhang, Z., Liu, M. C., Best, W. M. J., Dupuy, T. J., & Siverd, R. J. 2021, *ApJ*, **911**, 7
- Zhou, Y., Biller, B. A., Carter, A. L., et al. 2026, *AJ*, Submitted
- Zuckerman, B., Song, I., Bessell, M. S., & Webb, R. A. 2001, *ApJL*, **562**, L87